

Adolescent Behaviors and Experiences Survey				
Sexual Behaviors	Opposite Sex Only		Same Sex Only or Both Sexes	
	Percentage	Confidence Interval	Percentage	Confidence Interval
Ever had sexual intercourse	71.3	68.5 - 74.0	60.9	54.2 - 67.1
Had sexual intercourse for the first time before age 13 years	4.3	3.3 - 5.7	10.4	7.7 - 14.0
Had sexual intercourse with four or more persons during their life	13.6	11.4 - 16.2	16.6	11.9 - 22.8
Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	48.2	44.9 - 51.6	34.0	28.1 - 40.5
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	15.1	12.1 - 18.6	28.5	19.1 - 40.3
Used a condom during last sexual intercourse (among students who were currently sexually active)	57.0	53.9 - 60.2	30.0	20.4 - 41.8
Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active)	22.2	19.0 - 25.9	20.7	10.1 - 37.7
Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)	9.4	8.1 - 10.8	17.7	12.7 - 24.2
Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	8.8	7.3 - 10.5	12.9	8.8 - 18.5
Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	5.8	4.3 - 7.8	6.8	3.2 - 13.7
Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	34.1	30.4 - 38.1	29.8	18.0 - 45.2
Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	12.9	10.9 - 15.1	3.4	1.5 - 7.7
Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active)	13.0	10.8 - 15.6	17.7	10.6 - 28.2