

Adolescent Behaviors and Experiences Survey

Alcohol and Other Drug Use	Percentage	Confidence Interval
Had their first drink of alcohol before age 13 years (other than a few sips)	11.1	9.7 - 12.6
Currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	19.5	16.8 - 22.4
Currently were binge drinking (had four or more drinks of alcohol in a row if they were female or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey)	7.7	5.9 - 10.1
Reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)	2.2	1.6 - 3.1
Usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	38.4	34.3 - 42.7
Ever used marijuana (one or more times during their life)	24.9	22.8 - 27.2
Tried marijuana for the first time before age 13 years	3.4	2.7 - 4.4
Currently used marijuana (one or more times during the 30 days before the survey)	12.8	11.1 - 14.7
Ever used synthetic marijuana (one or more times during their life)	5.8	4.8 - 6.9
Ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	9.6	8.5 - 10.9
Currently took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (one or more times during the 30 days before the survey)	4.3	3.6 - 5.2
Ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	1.4	1.0 - 1.9
Ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	5.8	4.8 - 6.9
Ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	0.7	0.4 - 1.0
Ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	0.8	0.6 - 1.2
Ever used ecstasy (also called "MDMA" or "Molly," one or more times during their life)	1.9	1.4 - 2.7
Ever used hallucinogenic drugs (such as LSD, acid, PCP, angel dust, mescaline, or mushrooms, one or more times during their life)	3.8	2.9 - 4.9
Ever used select illicit drugs (counting cocaine, inhalants, heroin, methamphetamines, ecstasy, or hallucinogens, one or more times during their life)	9.1	7.7 - 10.7
Ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	0.5	0.4 - 0.8
Were offered, sold, or given an illegal drug on school property (during the 12 months before the survey, among students who attended school in-person during the 12 months before the survey)	9.4	7.4 - 11.8