

## Adolescent Behaviors and Experiences Survey

Sexual Behaviors	Percentage	Confidence Interval
Ever had sexual intercourse	28.5	25.9 - 31.3
Had sexual intercourse for the first time before age 13 years	2.2	1.7 - 2.8
Had sexual intercourse with four or more persons during their life	5.7	4.6 - 7.2
Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	18.8	16.6 - 21.2
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	16.8	14.3 - 19.6
Used a condom during last sexual intercourse (among students who were currently sexually active)	54.0	50.5 - 57.4
Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active)	21.9	18.8 - 25.5
Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)	5.8	5.0 - 6.9
Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	4.6	3.8 - 5.6
Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	5.8	4.4 - 7.7
Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	33.7	30.0 - 37.5
Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	11.7	9.8 - 13.8
Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active)	13.4	11.2 - 15.8