



**Magnetic Resonance Imaging:**

30. Was MRI of spinal cord performed?  yes  no  unknown    31. If yes, date of spine MRI: \_\_/\_\_/\_\_\_\_  
32. Did the spinal MRI show a lesion in at least some spinal cord gray matter?  yes  no  unknown  
33. Was MRI of brain performed?  yes  no  unknown    34. If yes, date of brain MRI: \_\_/\_\_/\_\_\_\_

**CSF examination:** 35. Was a lumbar puncture performed?  yes  no  unknown  
If yes, complete 35 (a,b) (If more than 2 CSF examinations, list the first 2 performed)

	Date of lumbar puncture	WBC/mm <sup>3</sup>	% neutrophils	% lymphocytes	% monocytes	% eosinophils	RBC/mm <sup>3</sup>	Glucose mg/dl	Protein mg/dl
35a. CSF from LP1									
35b. CSF from LP2									

**At time of 60 day follow-up please collect and send the following information: for confirmed and probable cases**

- Discharge summary    History and physical (H&P)    Neurology consult notes    EMG report (if done)  
 Infectious disease consult notes (if available)    Vaccine registry record    Diagnostic laboratory reports

**Acute Flaccid Myelitis Outcome – complete follow-up for all reported persons under investigation (PUI) at 60 days after onset of limb weakness.**

36. Date of follow-up: \_\_\_/\_\_\_/\_\_\_\_ (mm/dd/yyyy)

37. Impairment:    None    Minor (any minor involvement)    Significant ( $\leq 2$  extremities, major involvement)  
                           Severe ( $\geq 3$  extremities and respiratory involvement)    Death    Unknown

37a. Date of death: \_\_\_/\_\_\_/\_\_\_\_ (mm/dd/yyyy)

38. **Physical condition** (includes cardiovascular, gastrointestinal, urologic, endocrine as well as neurologic disorders):

- i. Medical problems sufficiently stable that medical or nursing monitoring is not required more often than 3-month intervals
- ii. Medical or nurse monitoring is needed more often than 3-month intervals but not each week.
- iii. Medical problems are sufficiently unstable as to require medical and/or nursing attention at least weekly.
- iv. Medical problems require intensive medical and/or nursing attention at least daily (excluding personal care assistance)

39. **Upper limb functions:** Self-care activities (drink/feed, dress upper/lower, brace/prosthesis, groom, wash, perineal care) dependent mainly upon upper limb function:

- i. Age-appropriate independence in self-care without impairment of upper limbs
- ii. Age-appropriate independence in self-care with some impairment of upper limbs
- iii. Dependent upon assistance in self-care with or without impairment of upper limbs.
- iv. Dependent totally in self-care with marked impairment of upper limbs.

40. **Lower limb functions:** Mobility (walk, stairs, wheelchair, transfer chair/toilet/tub or shower) dependent mainly upon lower limb function:

- i. Independent in mobility without impairment of lower limbs
- ii. Independent of mobility with some impairment of lower limbs, such as needing ambulatory aids, a brace or prosthesis
- iii. Dependent upon assistance or supervision in mobility with or without impairment of lower limbs.
- iv. Dependant totally in mobility with marked impairment of lower limbs.

41. **Sensory components:** Relating to communication (speech and hearing) and vision:

- i. Age-appropriate independence in communication and vision without impairment
- ii. Age-appropriate independence in communication and vision with some impairment such as mild dysarthria, mild aphasia or need for eyeglasses or hearing aid.
- iii. Dependent upon assistance, an interpreter, or supervision in communication or vision
- iv. Dependent totally in communication or vision

42. **Excretory functions** (bladder and bowel control, age-appropriate):

- i. Complete voluntary control of bladder and bowel sphincters
- ii. Control of sphincters allows normal social activities despite urgency or need for catheter, appliance, suppositories, etc.
- iii. Dependent upon assistance in sphincter management
- iv. Frequent wetting or soiling from bowel or bladder incontinence

43. **Support factors:**

- i. Able to fulfil usual age-appropriate roles and perform customary tasks
- ii. Must make some modifications in usual age-appropriate roles and performance of customary tasks
- iii. Dependent upon assistance, supervision, and encouragement from an adult due to any of the above considerations
- iv. Dependent upon long-term institutional care (chronic hospitalization, residential rehabilitation, etc. Excluding time-limited hospitalization for specific evaluation or treatment)

---

## Acute Flaccid Myelitis case definition

([https://cdn.ymaws.com/www.cste.org/resource/resmgr/ps/ps2021/21-ID-02\\_AFM.pdf](https://cdn.ymaws.com/www.cste.org/resource/resmgr/ps/ps2021/21-ID-02_AFM.pdf))

### Clinical Criteria

An illness with onset of acute flaccid\* limb weakness **AND**

Absence of a clear alternative diagnosis attributable to a nationally notifiable condition.

*\* Low muscle tone, limp, hanging loosely, not spastic or contracted.*

### Confirmatory laboratory/imaging evidence:

MRI showing spinal cord lesion with predominant gray matter involvement\* and spanning one or more vertebral segments, **AND**  
Excluding persons with gray matter lesions in the spinal cord resulting from physician diagnosed malignancy, vascular disease, or anatomic abnormalities.

### Presumptive laboratory/imaging evidence:

MRI showing spinal cord lesion where gray matter involvement\* is present but predominance cannot be determined, **AND**  
Excluding persons with gray matter lesions in the spinal cord resulting from physician diagnosed malignancy, vascular disease, or anatomic abnormalities.

### Supportive laboratory/imaging evidence:

MRI showing a spinal cord lesion in at least some gray matter\* and spanning one or more vertebral segments, **AND**  
Excluding persons with gray matter lesions in the spinal cord resulting from physician diagnosed malignancy, vascular disease, or anatomic abnormalities.

*\* Spinal cord lesions may not be present on initial MRI; a negative or normal MRI performed within the first 72 hours after onset of limb weakness does not rule out AFM. Terms in the spinal cord MRI report such as "affecting mostly gray matter," "affecting the anterior horn or anterior horn cells," "affecting the central cord," "anterior myelitis," or "poliomyelitis" would all be consistent with this terminology.*

### Other classification criteria

Autopsy findings that include histopathologic evidence of inflammation largely involving the anterior horn of the spinal cord spanning one or more vertebral segments.

### Vital Records Criteria

Any person whose death certificate lists acute flaccid myelitis as a cause of death or a condition contributing to death.

### Case Classification

#### **Confirmed:**

Meets clinical criteria with confirmatory laboratory/imaging evidence, **OR**

Meets other classification criteria.

#### **Probable:**

Meets clinical criteria with presumptive laboratory/imaging evidence.

#### **Suspect:**

Meets clinical criteria with supportive laboratory/imaging evidence, **AND**

Available information is insufficient to classify case as probable or confirmed.

### Acute Flaccid Myelitis specimen collection information

(<https://www.cdc.gov/acute-flaccid-myelitis/hcp/specimen-collection.html>)

### Acute Flaccid Myelitis job aid

(<https://www.cdc.gov/acute-flaccid-myelitis/downloads/job-aid-for-clinicians-508.pdf>)