

# INTRODUCTION TO FERTILITY CLINIC TABLES

---

The first table in this section is the 2012 National Summary of combined data from all clinics. Individual clinic tables follow, with each clinic's data presented in a one-page table that includes the types of assisted reproductive technology (ART) used, patient diagnoses, success rates, and individual clinic characteristics. Clinics reporting their data to the Centers for Disease Control and Prevention (CDC) are listed in alphabetical order by state, city, and clinic name. Each known nonreporting clinic is also included in alphabetical order, although no data are presented for these clinics.

Many people considering ART will want to use this report to find the “best” clinic. However, comparisons between clinics must be made with caution. Many factors contribute to the success of an ART procedure. Some factors are related to the training and experience of the ART clinic and laboratory professionals and the quality of services they provide. Other factors are related to the patients themselves, such as their age, quality of their eggs and sperm, cause of their infertility, genetic factors, and diagnosis. Some clinics may be more willing than others to accept patients with low chances of success or may specialize in ART treatments that attract particular types of patients.

We encourage consumers considering ART to contact clinics to discuss their specific medical situations and their potential for success using ART. Because clinics did not have the opportunity to provide narratives to explain their data in this report, such conversations could provide additional information to help consumers decide whether to use ART.

Although ART offers important options for the treatment of infertility, the decision to use ART involves many factors in addition to success rates. Therefore, consumers should carefully examine all related financial, psychological, and medical issues before beginning treatment. They also will want to consider the location of the clinic, the counseling and support services available, and the rapport that staff members have with their patients.

## Important Factors to Consider When Using These Tables to Assess a Clinic

- ***ART statistics are from 2012***

Data for cycles started in 2012 could not be published until 2014 because the final outcomes of pregnancies conceived in December 2012 were not known until October 2013. Additional time was then required to collect and analyze the data and prepare the report. Many factors that contribute to a clinic's success rate may have changed in the 2 years since these cycles were performed. Personnel may be different. Equipment and training may or may not have been updated. As a result, success rates for 2012 may not necessarily represent current rates.

- **Success rates may vary**

A clinic's success rates may vary from year to year even if all determining factors remain the same. The more cycles that a clinic carries out, the less the rate is likely to vary. Conversely, clinics that perform fewer cycles are likely to have more variability in success rates from year to year. As an extreme example, if a clinic reports only one ART cycle in a given category, as is sometimes the case in the data presented here, the clinic's success rate in that category would be either 0% or 100%. Because success rates may be misleading if they are based on a small number, the rates are shown as fractions rather than percentages when fewer than 20 cycles or outcomes (pregnancies, transfers, births) are reported in a given category.

- **Some clinics see more than the average number of patients with difficult infertility problems**

Some clinics are willing to offer ART to most potential patients, even those who have a low probability of success. Others discourage such patients or encourage them to use donor eggs, a practice that results in higher success rates among older women. Clinics that accept a higher percentage of women who previously have had multiple unsuccessful ART cycles will generally have lower success rates. In contrast, clinics that offer ART procedures to women who might have become pregnant with less technologically advanced treatment will generally have higher success rates.

- **The percentage of cycles that are canceled varies**

Percentages of canceled cycles using fresh nondonor eggs or embryos vary among clinics from less than 1% to, in a few cases, more than 25%. A high percentage of cancellations tends to lower the percentage of cycles resulting in live births but may increase the percentage of embryo transfers resulting in live births.

- **Percentages of unstimulated (or "natural") cycles are included with those for stimulated cycles**

In an unstimulated cycle, the woman ovulates naturally rather than as the result of the daily injections used in stimulated cycles. Unstimulated cycles are less expensive because they require no daily injections and fewer ultrasounds and blood tests. However, women who use natural or mild stimulation produce only one or two follicles, thus reducing the potential number of embryos for transfer. As a result, clinics that perform a relatively high percentage of unstimulated cycles may have lower success rates. Nationally, 1% of ART cycles using fresh nondonor eggs or embryos in 2012 were unstimulated.

- **Success rates are calculated per cycle rather than per patient**

Success rates shown in this report are presented in terms of cycles, as required by law, rather than in terms of patients. As a result, patients who had more than one ART cycle in 2012 are represented in multiple cycles that are not linked. In addition, for patients who undergo both fresh and frozen cycles, success rates are calculated separately for each cycle. Clinics that have a very high percentage of cycles resulting in live births with frozen embryos would have higher ART success rates if these births were included as successes from the original stimulated cycle. Consumers should look at both rates (for cycles using fresh embryos and for those using frozen embryos) when assessing a clinic's success rates.

- ***The number of embryos transferred varies from clinic to clinic***

In 2012, the average number of embryos that nearly all clinics transferred to women younger than age 35 ranged from 1 to 3 for fresh nondonor cycles. The American Society for Reproductive Medicine (ASRM) and the Society for Assisted Reproductive Technology (SART) discourage the transfer of a large number of embryos because of the increased likelihood of multiple-fetus pregnancies. Multiple-fetus pregnancies, in turn, increase the probability of premature births and related health problems.

An explanation of how to read a fertility clinic table begins on page 15.