State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)	II II	Weighted Number	95% CI¶ (Weighted Number)
U.S. Total**	18-24	25,187	10.0	0.30	(9.4–10.6)	Ш	3,115,838	(2,926,471–3,305,205)
U.S. Total**	25-34	45,248	8.4	0.22	(8.0–8.9)	Ш	3,704,589	(3,507,626–3,901,552)
U.S. Total**	35-44	50,447	8.9	0.24	(8.4–9.3)	Ш	3,578,001	(3,382,160–3,773,842)
U.S. Total**	45-54	62,372	9.9	0.23	(9.5–10.4)	Ш	3,913,368	(3,727,519–4,099,216)
U.S. Total**	55-64	87,328	10.2	0.21	(9.8–10.6)	II	4,260,075	(4,079,502–4,440,648)
U.S. Total**	65+	148,882	8.6	0.17	(8.3–8.9)	Ш	4,511,074	(4,329,139–4,693,008)
AL	18-24	302	13.9	2.48	(9.0–18.8)	Ш	65,073	(40,387–89,759)
AL	25-34	604	7.8	1.25	(5.3–10.2)	Ш	48,845	(32,871–64,818)
AL	35-44	760	10.1	1.29	(7.5–12.6)	Ш	59,269	(43,590–74,948)
AL	45-54	979	11.0	1.20	(8.6–13.3)	II	63,626	(49,316–77,936)
AL	55-64	1,429	11.9	1.06	(9.8–14.0)	II	75,645	(61,761–89,529)
AL	65+	2,398	9.7	0.77	(8.2–11.2)	II	80,903	(67,950–93,856)
AK	18-24	173	13.5	2.92	(7.8–19.3)	II	9,891	(5,421–14,361)
AK	25-34	302	7.3	1.68	(4.0–10.6)	II	8,402	(4,516–12,287)
AK	35-44	355	5.3	1.53	(2.3–8.4)	Ш	4,914	(2,076–7,753)
AK	45-54	383	12.2	2.16	(7.9–16.4)	II	10,392	(6,550–14,235)
AK	55-64	618	10.6	1.93	(6.8–14.4)	II	9,750	(6,046–13,454)
AK	65+	868	7.6	1.17	(5.3–9.9)	Ш	6,582	(4,539–8,624)
AZ	18-24	411	8.9	1.80	(5.4–12.5)	Ш	61,042	(35,785–86,299)
AZ	25-34	795	8.1	1.31	(5.5–10.6)	II	75,745	(50,873–100,617)
AZ	35-44	843	10.6	1.61	(7.4–13.8)	II	92,095	(63,047–121,144)
AZ	45-54	1,046	11.9	1.45	(9.1–14.8)	II	93,337	(70,008–116,665)
AZ	55-64	1,568	12.7	1.46	(9.8–15.5)	Ш	104,355	(79,284–129,426)
AZ	65+	3,161	9.0	0.73	(7.5–10.4)	Ш	110,521	(92,496–128,547)
AR	18-24	166	13.4	3.09	(7.3–19.5)	Ш	38,555	(19,619–57,491)
AR	25-34	367	9.0	1.61	(5.9–12.2)	Ш	34,484	(21,977–46,990)
AR	35-44	429	8.8	1.56	(5.8–11.9)	Ш	32,324	(20,674–43,975)
AR	45-54	637	11.3	1.43	(8.4–14.1)	Ш	39,279	(29,054–49,505)
AR	55-64	1,030	10.1	1.14	(7.9–12.3)	Ш	37,942	(29,194–46,690)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)	II II II	Weighted Number	95% CI¶ (Weighted Number)
AR	65+	2,609	8.1	0.67	(6.8–9.4)	Ш	41,147	(34,261–48,033)
CA	18-24	1,125	8.4	1.05	(6.3–10.5)	II	316,882	(235,964–397,799)
CA	25-34	1,965	6.5	0.69	(5.2–7.9)	II	382,745	(300,585–464,906)
CA	35-44	1,960	8.9	0.90	(7.1–10.6)	II	459,253	(363,640–554,866)
CA	45-54	1,883	8.9	0.82	(7.3–10.5)	Ш	444,722	(360,997–528,448)
CA	55-64	2,115	9.2	0.81	(7.6–10.8)	Ш	444,490	(365,555–523,425)
CA	65+	2,703	9.5	0.82	(7.9–11.1)	Ш	541,656	(446,370–636,941)
СО	18-24	546	10.5	1.42	(7.7–13.3)	Ш	56,340	(40,709–71,972)
СО	25-34	1,048	7.8	0.90	(6.0–9.6)	Ш	65,052	(49,748–80,355)
СО	35-44	1,109	10.8	1.02	(8.8–12.8)	Ш	80,418	(64,836–96,000)
СО	45-54	1,277	7.6	0.82	(5.9–9.2)	Ш	51,786	(40,325–63,248)
СО	55-64	1,778	9.4	0.76	(7.9–10.9)	Ш	67,685	(56,488–78,881)
СО	65+	2,910	8.4	0.59	(7.3–9.6)	Ш	68,346	(58,649–78,044)
СТ	18-24	438	9.1	1.45	(6.2–11.9)	Ш	31,653	(21,380–41,926)
СТ	25-34	864	9.1	1.12	(6.9–11.3)	Ш	39,431	(29,545–49,317)
СТ	35-44	1,022	9.9	1.07	(7.8–12.0)	Ш	40,769	(31,725–49,812)
СТ	45-54	1,705	13.6	1.06	(11.5–15.7)	Ш	61,861	(51,720–72,002)
СТ	55-64	2,360	10.8	0.82	(9.2–12.4)	Ш	53,878	(45,486–62,271)
СТ	65+	3,832	9.5	0.72	(8.1–10.9)	Ш	55,603	(46,886–64,320)
DE	18-24	311	13.4	2.63	(8.2–18.6)	Ш	11,732	(6,864–16,601)
DE	25-34	520	11.7	2.02	(7.7–15.6)	Ш	14,725	(9,362–20,089)
DE	35-44	617	9.9	1.57	(6.9–13.0)	Ш	11,060	(7,464–14,657)
DE	45-54	767	10.6	1.51	(7.6–13.6)	Ш	12,191	(8,582–15,800)
DE	55-64	1,078	10.2	1.20	(7.8–12.6)	Ш	13,303	(10,066–16,540)
DE	65+	1,814	7.2	0.69	(5.9–8.6)	Ш	12,999	(10,545–15,454)
DC	18-24	182	11.1	2.82	(5.5–16.7)	Ш	7,761	(3,587–11,935)
DC	25-34	463	10.3	1.53	(7.3–13.3)	Ш	16,213	(11,255–21,172)
DC	35-44	552	12.4	1.70	(9.1–15.8)	II	12,845	(9,132–16,559)
DC	45-54	657	14.4	1.56	(11.3–17.4)	II	10,673	(8,243–13,102)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
DC	55-64	811	14.4	1.54	(11.4–17.4)	Ш	10,392	(8,029–12,755)
DC	65+	1,567	9.2	0.88	(7.5–10.9)	Ш	7,752	(6,245–9,260)
FL	18-24	874	8.5	1.68	(5.2–11.8)	Ш	149,779	(89,471–210,087)
FL	25-34	1,597	8.5	1.27	(6.1–11.0)	Ш	232,184	(161,752–302,615)
FL	35-44	1,580	7.6	1.10	(5.5–9.8)	Ш	191,689	(136,602–246,776)
FL	45-54	2,151	7.0	1.00	(5.1–9.0)	Ш	185,303	(132,204–238,402)
FL	55-64	2,807	12.9	1.40	(10.1–15.6)	Ш	361,925	(278,910–444,940)
FL	65+	5,946	8.1	0.85	(6.4–9.8)	Ш	359,224	(281,668–436,780)
GA	18-24	676	8.9	1.22	(6.5–11.3)	Ш	92,277	(66,266–118,288)
GA	25-34	1,114	7.7	0.96	(5.8–9.5)	Ш	107,254	(79,917–134,591)
GA	35-44	1,241	8.3	0.93	(6.5–10.2)	II	112,518	(86,887–138,149)
GA	45-54	1,406	10.0	0.95	(8.2–11.9)	II	132,317	(106,736–157,898)
GA	55-64	1,786	10.8	0.93	(8.9–12.6)	Ш	136,368	(111,837–160,898)
GA	65+	2,833	8.3	0.65	(7.0–9.6)	Ш	122,787	(103,324–142,251)
н	18-24	519	12.8	2.01	(8.8–16.7)	Ш	15,918	(10,560–21,276)
н	25-34	925	9.8	1.27	(7.3–12.3)	Ш	19,538	(14,320–24,756)
н	35-44	1,047	7.3	1.10	(5.2–9.5)	Ш	13,398	(9,294–17,501)
н	45-54	1,178	9.3	1.00	(7.3–11.2)	Ш	14,484	(11,307–17,661)
н	55-64	1,552	10.2	1.01	(8.2–12.1)	Ш	18,281	(14,545–22,016)
н	65+	2,521	8.0	0.71	(6.6–9.3)	Ш	20,332	(16,742–23,922)
ID	18-24	266	8.1	2.04	(4.1–12.1)	II	13,841	(6,730–20,952)
ID	25-34	413	10.6	1.91	(6.8–14.3)	Ш	23,252	(14,531–31,974)
ID	35-44	422	7.0	1.60	(3.9–10.2)	II	15,188	(8,206–22,169)
ID	45-54	438	7.6	1.46	(4.7–10.5)	II	13,846	(8,516–19,177)
ID	55-64	663	10.7	1.52	(7.8–13.7)	II	22,209	(15,766–28,652)
ID	65+	1,390	7.4	0.88	(5.7–9.2)	Ш	20,548	(15,687–25,410)
IL	18-24	420	10.0	1.85	(6.3–13.6)	Ш	120,886	(74,271–167,501)
IL	25-34	705	9.1	1.19	(6.7–11.4)	Ш	154,746	(113,338–196,153)
IL	35-44	736	8.1	1.19	(5.7–10.4)	Ш	130,908	(91,584–170,232)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>Cl denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
IL	45-54	857	8.8	1.09	(6.7–11.0)	II	140,552	(105,247–175,857)
IL	55-64	1,015	9.0	0.99	(7.0–10.9)	Ш	146,950	(113,824–180,076)
IL	65+	1,496	7.9	0.80	(6.3–9.4)	Ш	156,740	(124,973–188,506)
IN	18-24	363	10.5	1.75	(7.1–14.0)	Ш	71,522	(47,071–95,973)
IN	25-34	624	8.5	1.26	(6.0–11.0)	Ш	71,690	(49,937–93,443)
IN	35-44	728	8.9	1.11	(6.7–11.1)	Ш	72,694	(54,321–91,066)
IN	45-54	1,114	12.3	1.11	(10.1–14.4)	Ш	94,500	(76,891–112,108)
IN	55-64	1,583	10.3	0.86	(8.6–12.0)	Ш	88,391	(73,230–103,551)
IN	65+	2,921	9.3	0.63	(8.0–10.5)	Ш	95,692	(82,481–108,903)
IA	18-24	641	9.8	1.31	(7.2–12.3)	Ш	32,105	(23,212–40,999)
IA	25-34	992	7.5	0.89	(5.7–9.2)	II	28,706	(21,741–35,672)
IA	35-44	1,219	7.2	0.78	(5.7–8.7)	II	27,013	(21,097–32,930)
IA	45-54	1,348	9.4	0.91	(7.6–11.2)	Ш	32,661	(26,171–39,152)
IA	55-64	1,845	7.6	0.68	(6.2–8.9)	Ш	30,951	(25,290–36,613)
IA	65+	2,900	7.0	0.52	(5.9–8.0)	Ш	37,607	(31,915–43,298)
KS	18-24	652	13.4	1.54	(10.4–16.4)	Ш	40,366	(30,554–50,178)
KS	25-34	1,078	9.7	1.06	(7.7–11.8)	Ш	35,878	(27,817–43,939)
KS	35-44	1,188	9.6	0.95	(7.7–11.4)	Ш	33,635	(26,845–40,426)
KS	45-54	1,520	9.1	0.80	(7.5–10.6)	Ш	28,784	(23,658–33,910)
KS	55-64	2,183	9.5	0.73	(8.1–11.0)	Ш	34,883	(29,382–40,384)
KS	65+	3,937	8.6	0.53	(7.6–9.7)	II	39,026	(34,142–43,911)
KY	18-24	555	11.7	2.16	(7.5–16.0)	II	49,687	(30,483–68,891)
KY	25-34	837	10.9	1.53	(7.8–13.9)	Ш	61,670	(43,696–79,644)
KY	35-44	944	9.3	1.41	(6.5–12.0)	Ш	50,669	(34,948–66,389)
KY	45-54	1,389	13.4	1.54	(10.3–16.4)	Ш	74,065	(56,100–92,029)
KY	55-64	1,880	12.5	1.26	(10.1–15.0)	Ш	73,585	(58,250–88,921)
KY	65+	2,770	10.7	1.03	(8.7–12.7)	Ш	78,739	(63,029–94,449)
LA	18-24	359	9.9	1.86	(6.2–13.5)	Ш	43,664	(26,702–60,626)
LA	25-34	646	10.8	2.41	(6.0–15.5)	Ш	69,598	(36,235–102,961)

### Notes:

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)	II II	Weighted Number	95% CI¶ (Weighted Number)
LA	35-44	673	7.0	1.32	(4.4–9.6)	Ш	40,708	(25,096–56,320)
LA	45-54	833	9.5	1.21	(7.1–11.9)	Ш	49,652	(36,782–62,522)
LA	55-64	1,103	10.7	1.34	(8.1–13.3)	Ш	63,103	(46,753–79,452)
LA	65+	1,392	6.6	0.89	(4.9–8.4)	Ш	47,282	(34,534–60,029)
ME	18-24	313	19.5	3.40	(12.8–26.2)	Ш	22,038	(13,506–30,571)
ME	25-34	711	11.4	1.55	(8.3–14.4)	Ш	17,091	(12,287–21,895)
ME	35-44	890	13.6	1.52	(10.6–16.5)	Ш	20,215	(15,501–24,930)
ME	45-54	1,415	13.0	1.25	(10.5–15.4)	Ш	22,050	(17,623–26,477)
ME	55-64	2,403	10.8	0.85	(9.1–12.4)	Ш	22,895	(19,185–26,604)
ME	65+	4,951	9.7	0.59	(8.5–10.8)	Ш	25,926	(22,704–29,148)
MD	18-24	633	13.1	1.76	(9.6–16.5)	Ш	72,320	(51,701–92,938)
MD	25-34	1,278	9.5	1.10	(7.3–11.6)	II	76,758	(58,466–95,050)
MD	35-44	1,653	8.3	0.86	(6.6–10.0)	Ш	62,868	(49,708–76,027)
MD	45-54	2,618	9.5	0.76	(8.1–11.0)	Ш	72,008	(60,328–83,687)
MD	55-64	3,678	10.3	0.64	(9.0–11.5)	Ш	80,663	(70,395–90,931)
MD	65+	6,932	7.3	0.42	(6.4–8.1)	Ш	66,681	(58,936–74,427)
MA	18-24	497	13.7	1.94	(9.9–17.5)	Ш	97,034	(67,905–126,163)
MA	25-34	794	9.8	1.38	(7.1–12.6)	Ш	88,434	(62,827–114,041)
MA	35-44	731	10.2	1.29	(7.6–12.7)	Ш	82,711	(61,397–104,026)
MA	45-54	987	10.7	1.19	(8.4–13.0)	Ш	89,952	(69,379–110,526)
MA	55-64	1,310	8.2	0.84	(6.6–9.9)	Ш	76,729	(61,035–92,424)
MA	65+	2,082	9.7	0.90	(8.0–11.5)	Ш	108,966	(88,372–129,559)
МІ	18-24	810	14.0	1.49	(11.1–16.9)	Ш	137,830	(106,926–168,734)
MI	25-34	1,217	10.5	1.12	(8.3–12.7)	Ш	131,916	(102,952–160,879)
MI	35-44	1,226	11.7	1.15	(9.5–14.0)	Ш	134,535	(107,214–161,856)
MI	45-54	1,538	12.5	1.06	(10.4–14.6)	Ш	149,815	(123,103–176,528)
MI	55-64	2,112	11.1	0.84	(9.4–12.7)	Ш	153,000	(129,146–176,853)
MI	65+	3,187	8.6	0.66	(7.3–9.9)	Ш	145,302	(122,570–168,034)
MN	18-24	1,201	9.4	0.99	(7.4–11.3)	П	48,840	(38,259–59,421)

### Notes:

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>Cl denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
MN	25-34	1,991	8.7	0.75	(7.2–10.2)	II	63,402	(52,282–74,521)
MN	35-44	2,352	7.1	0.60	(5.9–8.3)	II	50,008	(41,434–58,582)
MN	45-54	2,651	8.9	0.66	(7.6–10.2)	II	58,824	(49,828–67,821)
MN	55-64	3,539	9.0	0.58	(7.8–10.1)	II	67,396	(58,521–76,270)
MN	65+	4,966	7.1	0.45	(6.2–8.0)	II	63,476	(55,392–71,561)
MS	18-24	336	8.6	1.84	(5.0–12.2)	II	25,727	(14,359–37,095)
MS	25-34	586	7.2	1.23	(4.8–9.6)	II	27,577	(18,042–37,113)
MS	35-44	734	10.5	1.33	(7.9–13.1)	II	38,110	(28,261–47,960)
MS	45-54	919	9.7	1.16	(7.4–12.0)	II	33,695	(25,475–41,915)
MS	55-64	1,267	10.5	1.07	(8.4–12.6)	II	38,619	(30,604–46,634)
MS	65+	1,884	10.9	0.98	(9.0–12.8)	II	51,688	(42,134–61,242)
МО	18-24	276	7.9	1.74	(4.5–11.3)	II	45,491	(25,142–65,841)
МО	25-34	553	8.6	1.53	(5.6–11.6)	II	67,398	(42,637–92,159)
МО	35-44	593	10.2	1.66	(6.9–13.5)	Ш	75,549	(50,033–101,065)
МО	45-54	777	10.9	1.35	(8.2–13.5)	Ш	78,212	(58,402–98,021)
МО	55-64	1,221	11.1	1.19	(8.8–13.5)	Ш	90,482	(70,510–110,453)
МО	65+	2,670	7.6	0.63	(6.4–8.9)	Ш	77,789	(65,024–90,553)
MT	18-24	286	11.6	2.16	(7.4–15.9)	Ш	11,615	(7,157–16,074)
MT	25-34	491	10.6	1.74	(7.2–14.1)	Ш	14,064	(9,289–18,838)
MT	35-44	607	10.0	1.51	(7.1–13.0)	Ш	12,516	(8,662–16,371)
MT	45-54	651	12.5	1.62	(9.3–15.7)	Ш	14,212	(10,358–18,065)
MT	55-64	1,124	10.1	1.20	(7.7–12.4)	Ш	15,261	(11,533–18,989)
МТ	65+	1,925	7.2	0.83	(5.6–8.8)	II	14,196	(10,909–17,482)
NE	18-24	919	10.5	1.45	(7.7–13.4)	II	20,701	(14,729–26,674)
NE	25-34	1,531	8.6	1.02	(6.6–10.6)	II	21,113	(15,985–26,241)
NE	35-44	1,806	7.4	0.88	(5.6–9.1)	II	17,521	(13,278–21,763)
NE	45-54	1,910	9.9	1.00	(7.9–11.8)	II	21,214	(16,806–25,622)
NE	55-64	2,899	9.6	0.81	(8.0–11.1)	II	22,920	(18,955–26,885)
NE	65+	5,385	8.2	0.57	(7.1–9.3)	II	24,939	(21,392–28,487)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
NV	18-24	248	10.0	2.45	(5.2–14.9)	Ш	25,718	(12,913–38,523)
NV	25-34	409	6.2	1.45	(3.3–9.0)	Ш	25,906	(13,648–38,164)
NV	35-44	392	8.3	2.42	(3.5–13.0)	II	32,856	(12,860–52,852)
NV	45-54	465	7.1	1.53	(4.1–10.1)	II	26,189	(14,900–37,479)
NV	55-64	587	11.3	2.49	(6.4–16.2)	Ш	42,833	(22,787–62,879)
NV	65+	1,041	6.2	0.93	(4.3–8.0)	Ш	29,838	(20,930–38,746)
NH	18-24	170	14.9	3.14	(8.7–21.1)	Ш	15,297	(8,376–22,218)
NH	25-34	346	12.4	2.25	(8.0–16.8)	II	22,915	(14,123–31,706)
NH	35-44	515	15.6	1.95	(11.8–19.4)	Ш	23,764	(17,474–30,054)
NH	45-54	841	13.3	1.39	(10.6–16.0)	Ш	23,770	(18,591–28,948)
NH	55-64	1,351	10.4	0.98	(8.5–12.4)	Ш	21,915	(17,723–26,107)
NH	65+	2,355	8.1	0.68	(6.8–9.5)	Ш	19,591	(16,312–22,869)
NJ	18-24	205	6.4	1.77	(2.9–9.9)	Ш	51,446	(23,692–79,200)
NJ	25-34	410	6.4	1.65	(3.2–9.7)	Ш	73,117	(35,643–110,590)
NJ	35-44	443	10.2	2.21	(5.9–14.6)	Ш	113,313	(62,850–163,776)
NJ	45-54	485	9.9	2.02	(5.9–13.9)	Ш	117,608	(68,412–166,804)
NJ	55-64	600	9.8	1.61	(6.6–12.9)	Ш	119,919	(80,621–159,217)
NJ	65+	878	7.4	1.37	(4.7–10.1)	Ш	108,411	(68,543–148,279)
NM	18-24	397	10.8	1.81	(7.3–14.4)	Ш	21,686	(14,189–29,184)
NM	25-34	721	6.1	0.98	(4.2–8.1)	Ш	16,630	(11,326–21,935)
NM	35-44	805	7.5	1.09	(5.4–9.7)	II	18,775	(13,271–24,278)
NM	45-54	941	12.3	1.62	(9.1–15.5)	II	28,578	(20,609–36,547)
NM	55-64	1,445	11.2	1.15	(8.9–13.5)	Ш	29,515	(23,268–35,762)
NM	65+	2,309	11.4	1.00	(9.5–13.4)	II	41,328	(33,892–48,765)
NY	18-24	2,185	11.5	1.00	(9.5–13.4)	II	212,061	(173,719–250,404)
NY	25-34	4,074	11.1	0.77	(9.6–12.6)	Ш	314,262	(269,647–358,876)
NY	35-44	4,232	8.4	0.67	(7.1–9.7)	Ш	192,822	(161,674–223,970)
NY	45-54	5,459	11.1	0.77	(9.6–12.6)	Ш	260,522	(222,946–298,097)
NY	55-64	7,236	9.8	0.61	(8.6–11.0)	Ш	247,866	(216,656–279,076)

### Notes:

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)	II II	Weighted Number	95% CI¶ (Weighted Number)
NY	65+	10,912	9.6	0.66	(8.3–10.9)	П	303,437	(260,489–346,386)
NC	18-24	373	12.9	2.12	(8.8–17.1)	II	129,871	(84,876–174,865)
NC	25-34	624	5.9	1.08	(3.8–8.0)	II	78,976	(49,913–108,039)
NC	35-44	616	10.5	1.73	(7.1–13.9)	II	132,217	(86,190–178,244)
NC	45-54	767	9.2	1.30	(6.7–11.8)	Ш	120,189	(85,416–154,963)
NC	55-64	828	11.0	1.41	(8.2–13.8)	Ш	147,667	(108,338–186,996)
NC	65+	1,406	8.4	1.07	(6.3–10.5)	Ш	141,674	(104,934–178,414)
ND	18-24	250	8.8	2.20	(4.5–13.2)	Ш	7,558	(3,651–11,465)
ND	25-34	467	6.5	1.26	(4.0–9.0)	Ш	7,117	(4,322–9,912)
ND	35-44	536	9.6	1.47	(6.7–12.5)	Ш	8,749	(5,991–11,507)
ND	45-54	689	7.9	1.15	(5.6–10.1)	Ш	6,046	(4,269–7,824)
ND	55-64	1,242	8.6	0.98	(6.7–10.6)	Ш	7,934	(6,094–9,774)
ND	65+	2,316	8.1	0.68	(6.7–9.4)	Ш	9,184	(7,630–10,739)
ОН	18-24	649	11.3	1.54	(8.3–14.3)	Ш	124,328	(89,418–159,237)
ОН	25-34	1,143	7.1	0.93	(5.3–9.0)	Ш	106,312	(78,512–134,113)
ОН	35-44	1,255	11.3	1.39	(8.5–14.0)	Ш	155,612	(115,230–195,994)
ОН	45-54	1,847	9.7	0.93	(7.9–11.5)	Ш	136,362	(109,631–163,094)
ОН	55-64	2,887	8.8	0.68	(7.4–10.1)	Ш	138,808	(116,860–160,756)
ОН	65+	4,778	9.1	0.63	(7.9–10.4)	Ш	182,263	(156,604–207,922)
OK	18-24	274	11.1	2.17	(6.8–15.3)	Ш	43,352	(25,597–61,107)
OK	25-34	548	10.2	1.44	(7.4–13.0)	Ш	53,566	(37,972–69,160)
OK	35-44	617	10.1	1.31	(7.5–12.7)	Ш	49,687	(36,531–62,842)
OK	45-54	682	12.4	1.50	(9.5–15.4)	Ш	54,866	(41,074–68,658)
ОК	55-64	1,064	10.4	1.21	(8.1–12.8)	Ш	49,918	(37,904–61,932)
ОК	65+	1,985	8.6	0.72	(7.2–10.0)	П	53,523	(44,523–62,524)
OR	18-24	417	11.2	1.79	(7.7–14.7)	II	41,272	(27,628–54,916)
OR	25-34	777	12.5	1.44	(9.6–15.3)	II	70,959	(53,592–88,326)
OR	35-44	822	11.9	1.23	(9.5–14.3)	II	63,967	(50,295–77,639)
OR	45-54	902	14.6	1.57	(11.5–17.7)	II	68,592	(52,784–84,399)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
OR	55-64	1,085	11.0	1.09	(8.9–13.2)	II	60,402	(48,234–72,570)
OR	65+	1,753	9.9	1.06	(7.8–11.9)	II	72,839	(56,725–88,952)
PA	18-24	469	10.7	1.85	(7.0–14.3)	II	126,874	(81,384–172,363)
PA	25-34	824	9.3	1.19	(7.0–11.7)	II	152,969	(113,533–192,405)
PA	35-44	812	9.9	1.29	(7.4–12.4)	II	146,290	(107,490–185,091)
PA	45-54	972	10.7	1.19	(8.3–13.0)	II	164,976	(127,420–202,532)
PA	55-64	1,225	11.5	1.11	(9.4–13.7)	II	208,037	(166,956–249,118)
PA	65+	1,772	8.9	0.95	(7.0–10.8)	II	206,140	(161,220–251,060)
RI	18-24	247	10.6	2.53	(5.6–15.6)	II	11,688	(5,876–17,501)
RI	25-34	486	12.8	1.99	(8.9–16.8)	II	18,303	(12,362–24,243)
RI	35-44	619	11.5	1.50	(8.5–14.4)	II	14,356	(10,507–18,205)
RI	45-54	832	12.7	1.44	(9.9–15.5)	II	16,121	(12,334–19,908)
RI	55-64	1,262	11.9	1.18	(9.6–14.2)	II	17,581	(13,954–21,208)
RI	65+	1,965	11.4	0.97	(9.5–13.3)	II	20,337	(16,745–23,928)
SC	18-24	441	9.2	1.71	(5.8–12.6)	II	43,995	(27,109–60,882)
SC	25-34	788	9.4	1.19	(7.1–11.7)	II	62,152	(46,075–78,229)
SC	35-44	1,030	8.5	1.02	(6.5–10.5)	II	52,062	(39,436–64,688)
SC	45-54	1,383	9.4	0.94	(7.5–11.2)	II	56,863	(45,284–68,442)
SC	55-64	2,136	10.8	0.89	(9.1–12.6)	Ш	71,890	(59,598–84,183)
SC	65+	4,820	7.6	0.46	(6.7–8.5)	II	68,136	(59,899–76,373)
SD	18-24	363	9.2	2.44	(4.4–14.0)	II	7,978	(3,622–12,333)
SD	25-34	688	8.7	1.76	(5.2–12.1)	II	9,329	(5,478–13,180)
SD	35-44	748	6.7	1.42	(3.9–9.5)	Ш	6,934	(4,004–9,865)
SD	45-54	950	8.5	1.58	(5.4–11.6)	II	7,804	(4,823–10,786)
SD	55-64	1,486	7.5	1.13	(5.3–9.8)	Ш	8,583	(6,001–11,165)
SD	65+	2,688	7.1	0.88	(5.4–8.8)	II	10,238	(7,678–12,799)
TN	18-24	296	7.0	1.95	(3.1–10.8)	Ш	42,885	(18,488–67,283)
TN	25-34	600	9.1	1.51	(6.2–12.1)	Ш	81,240	(53,783–108,697)
TN	35-44	672	8.4	1.37	(5.7–11.0)	Ш	69,396	(46,365–92,426)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
TN	45-54	785	10.6	1.55	(7.6–13.7)	II	87,254	(60,998–113,511)
TN	55-64	1,072	13.2	1.34	(10.6–15.8)	II	115,966	(91,495–140,436)
TN	65+	1,618	9.7	0.97	(7.8–11.6)	II	109,560	(87,463–131,658)
TX	18-24	723	6.8	1.28	(4.3–9.3)	II	196,883	(123,445–270,321)
TX	25-34	1,439	7.1	1.12	(4.9–9.3)	II	285,244	(194,973–375,515)
TX	35-44	1,389	6.8	1.21	(4.4–9.2)	II	249,515	(160,384–338,646)
TX	45-54	1,520	9.0	1.40	(6.3–11.8)	II	304,536	(208,681–400,390)
TX	55-64	1,970	7.5	1.15	(5.3–9.8)	II	241,666	(167,432–315,899)
TX	65+	3,824	7.6	0.93	(5.8–9.4)	II	277,970	(210,795–345,144)
UT	18-24	1,042	11.1	1.23	(8.7–13.6)	II	39,793	(30,619–48,966)
UT	25-34	1,625	8.3	0.84	(6.7–10.0)	II	37,108	(29,462–44,753)
UT	35-44	1,858	7.2	0.71	(5.8–8.6)	II	30,370	(24,344–36,396)
UT	45-54	1,544	10.0	0.93	(8.2–11.8)	II	30,844	(24,945–36,743)
UT	55-64	1,663	10.6	0.96	(8.7–12.5)	II	30,843	(25,067–36,620)
UT	65+	2,561	9.7	0.75	(8.2–11.2)	II	34,223	(28,776–39,670)
VT	18-24	224	15.5	2.76	(10.0–20.9)	II	10,196	(6,357–14,036)
VT	25-34	492	12.2	1.89	(8.5–15.9)	II	8,711	(5,902–11,520)
VT	35-44	700	11.7	1.50	(8.7–14.6)	II	8,067	(5,927–10,207)
VT	45-54	997	12.5	1.32	(9.9–15.1)	II	9,377	(7,320–11,434)
VT	55-64	1,496	12.5	1.14	(10.3–14.7)	II	11,557	(9,356–13,759)
VT	65+	2,405	9.9	0.84	(8.3–11.6)	II	11,710	(9,680–13,741)
VA	18-24	574	10.4	1.55	(7.4–13.5)	II	86,295	(59,895–112,695)
VA	25-34	1,088	8.8	1.11	(6.6–11.0)	II	98,913	(73,464–124,362)
VA	35-44	1,225	7.3	0.91	(5.6–9.1)	II	79,512	(59,800–99,224)
VA	45-54	1,643	9.1	0.85	(7.4–10.8)	II	97,732	(79,349–116,115)
VA	55-64	2,185	9.3	0.88	(7.6–11.0)	II	101,019	(81,382–120,655)
VA	65+	3,381	7.1	0.57	(6.0–8.2)	II	93,724	(78,697–108,752)
WA	18-24	670	12.3	1.48	(9.4–15.2)	II	82,575	(61,849–103,301)
WA	25-34	1,421	8.0	0.87	(6.3–9.7)	Ш	85,246	(66,366–104,125)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>Cl denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)		Weighted Number	95% CI¶ (Weighted Number)
WA	35-44	1,538	6.7	0.69	(5.3–8.0)	II	63,649	(50,412–76,886)
WA	45-54	1,648	11.1	0.96	(9.2–13.0)	II	95,561	(78,491–112,631)
WA	55-64	2,534	10.8	0.78	(9.2–12.3)	II	102,173	(86,948–117,399)
WA	65+	4,870	9.8	0.57	(8.7–10.9)	II	114,587	(100,953–128,221)
WV	18-24	218	12.2	2.42	(7.4–16.9)	II	19,751	(11,549–27,953)
WV	25-34	394	8.9	1.49	(5.9–11.8)	II	18,483	(12,152–24,813)
WV	35-44	557	11.9	1.55	(8.9–14.9)	II	25,414	(18,557–32,272)
WV	45-54	749	14.9	1.55	(11.8–17.9)	II	32,715	(25,455–39,975)
WV	55-64	1,083	15.0	1.33	(12.3–17.6)	II	38,087	(30,857–45,318)
WV	65+	1,823	11.1	0.92	(9.3–12.9)	II	39,771	(33,010–46,533)
WI	18-24	298	10.3	1.92	(6.5–14.1)	II	58,057	(36,142–79,973)
WI	25-34	518	10.3	1.80	(6.8–13.9)	II	73,938	(47,076–100,801)
WI	35-44	574	8.5	1.30	(5.9–11.0)	II	58,817	(40,675–76,959)
WI	45-54	739	9.9	1.42	(7.1–12.7)	II	67,466	(47,594–87,339)
WI	55-64	1,092	9.5	1.16	(7.2–11.7)	II	76,844	(57,485–96,203)
WI	65+	1,653	7.3	0.95	(5.4–9.1)	II	72,064	(52,967–91,161)
WY	18-24	204	10.3	2.12	(6.1–14.5)	II	5,705	(3,281–8,129)
WY	25-34	355	7.0	1.40	(4.2–9.7)	II	5,284	(3,129–7,440)
WY	35-44	475	8.9	1.47	(6.0–11.8)	II	6,456	(4,269–8,643)
WY	45-54	498	8.5	1.42	(5.8–11.3)	II	5,384	(3,551–7,217)
WY	55-64	1,042	9.1	1.01	(7.1–11.0)	II	7,001	(5,420–8,583)
WY	65+	1,819	8.8	0.80	(7.2–10.3)	II	8,074	(6,569–9,579)
Territories								
GU	18-24	151	3.4	1.30	(0.9–6.0)	II	611	(159–1,063)
GU	25-34	222	4.3	1.46	(1.5–7.2)	II	899	(297–1,502)
GU	35-44	232	3.7	1.49	(0.8–6.7)	II	836	(171–1,502)
GU	45-54	321	7.4	1.83	(3.8–11.1)	II	1,442	(729–2,154)
GU	55-64	331	3.0	1.01	(1.0–5.0)	II	424	(144–703)
GU	65+	354	9.8	2.81	(4.3–15.3)	II	1,047	(412–1,681)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>Cl denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question

State	Age Group	Sample Size§	Prevalence (percent)	Standard Error	95% CI¶ (percent)	===	Weighted Number	95% CI¶ (Weighted Number)
PR	18-24	350	12.2	2.08	(8.1–16.3)	Ш	40,853	(26,276–55,429)
PR	25-34	659	9.6	1.33	(6.9–12.2)	Ш	42,507	(30,557–54,458)
PR	35-44	720	10.8	1.42	(8.0–13.6)	II	47,499	(34,665–60,332)
PR	45-54	829	12.1	1.48	(9.2–15.0)	Ш	52,383	(38,941–65,826)
PR	55-64	923	13.0	1.34	(10.4–15.7)	Ш	55,988	(44,034–67,942)
PR	65+	1,309	9.9	0.96	(8.0–11.8)	Ш	62,033	(49,703–74,364)

### Notes:

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

<sup>\*</sup>Aged 18+ years

<sup>\*\*</sup>U.S. Total includes 50 states plus the District of Columbia and excludes the three territories.

<sup>¶</sup>CI denotes confidence interval.

<sup>§</sup>Sample size is the number of respondents answered current asthma question