

Table L1
Child[§] Lifetime Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2019

State	Sample Size	Prevalence (Percent)	Standard Error	95% CI [¶] (Percent)	 	Weighted Number	95% CI [¶] (Number)
Total *	47,981	11.3	0.31	(10.7–11.9)		5,580,439	(5,274,187–5,886,690)
CA	2,687	12.1	0.79	(10.7–13.8)		1,069,360	(927,951–1,210,769)
CT	1,500	14.2	1.22	(12.0–16.8)		99,012	(81,897–116,128)
DC	445	15.8	2.23	(11.9–20.7)		20,131	(14,387–25,875)
FL	2,583	11.1	1.05	(9.2–13.3)		448,509	(364,750–532,267)
GA	1,346	14.6	1.65	(11.7–18.1)		348,357	(268,254–428,460)
HI	1,692	13.9	1.03	(12.0–16.0)		39,971	(34,004–45,937)
IL	1,270	9.8	1.08	(7.9–12.1)		267,477	(208,192–326,762)
IN	1,653	9.7	0.79	(8.2–11.3)		143,627	(120,615–166,640)
KS	2,554	11.7	0.85	(10.1–13.5)		79,758	(68,047–91,470)
ME	1,498	13.1	1.29	(10.8–15.9)		29,870	(23,944–35,797)
MA	668	13.2	1.80	(10.1–17.2)		165,104	(118,879–211,330)
MI	2,133	11.0	0.88	(9.4–12.8)		227,053	(190,779–263,327)
MN	3,328	8.1	0.61	(7.0–9.4)		101,350	(86,365–116,335)
MS	1,090	11.1	1.29	(8.8–13.9)		73,553	(56,370–90,736)
MO	1,321	10.5	1.09	(8.6–12.9)		138,585	(110,328–166,842)
MT	1,428	8.7	0.90	(7.1–10.7)		19,597	(15,611–23,584)
NE	3,507	8.6	0.71	(7.3–10.1)		40,472	(33,837–47,107)
NV	667	11.2	1.88	(8.0–15.5)		72,250	(47,722–96,778)
NH	887	11.4	1.33	(9.0–14.2)		28,032	(21,587–34,476)
NM	1,230	8.5	1.24	(6.4–11.3)		40,377	(28,636–52,118)
NY	804	13.1	1.64	(10.2–16.7)		510,668	(379,706–641,631)
OH	2,162	11.5	1.08	(9.6–13.8)		285,782	(232,047–339,517)
OR	1,201	11.4	1.69	(8.5–15.2)		91,233	(62,972–119,493)
PA	1,382	12.3	1.07	(10.3–14.5)		306,930	(254,373–359,487)
RI	893	13.0	1.55	(10.2–16.3)		25,485	(19,409–31,561)
TX	2,382	9.9	1.12	(7.9–12.3)		709,036	(549,209–868,863)
UT	3,796	8.2	0.52	(7.3–9.3)		75,955	(66,596–85,314)
VT	1,067	12.8	1.42	(10.3–15.9)		14,471	(11,281–17,661)

Notes:

[§]Aged 0-17

[¶]CI denotes confidence interval.

* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention**

Table L1
Child[§] Lifetime Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2019

State	Sample Size	Prevalence (Percent)	Standard Error	95% CI [*] (Percent)	 	Weighted Number	95% CI [*] (Number)
WI	807	8.8	1.25	(6.6–11.6)		108,435	(78,337–138,533)
Territory							
PR	1,419	23.4	1.26	(21.0–26.0)		162,368	(144,711–180,025)

Notes:

[§]Aged 0-17

^{*}CI denotes confidence interval.

* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health
Centers for Disease Control and Prevention**