

CDC Survey of Maternity Practices in Infant Nutrition and Care

2013 Survey

Island Territories Results Report

mPINC Survey?

What is the The Maternity Practices in Infant Nutrition and Care (mPINC) Survey is a national survey from the Centers for Disease Control and Prevention (CDC) that assesses infant feeding care processes, policies, and staffing expectations in maternity care settings.

this report?

This report summarizes results from all Island Territories facilities that participated in the 2013 mPINC Survey and identifies opportunities to improve mother-baby care at hospitals and birth centers and related health outcomes throughout Island Territories.

Who participates in the mPINC survey?

All hospitals with maternity services and all free-standing birth centers in the United States are invited to participate in CDC's mPINC survey every two years.

Island Territories

Island Territories Highlights: Strengths



Documentation of Mothers' Feeding Decisions

Staff at all (100%) Island Territories facilities consistently ask about and record mothers' infant feeding decisions.

Standard documentation of infant feeding decisions is important to adequately support maternal choice.



Availability of Prenatal Breastfeeding Instruction

Most (83%) Island Territories facilities include breastfeeding education as a routine element of their prenatal classes.

Prenatal education about breastfeeding is important because it provides mothers with a better understanding of the benefits and requirements of breastfeeding, resulting in improved breastfeeding rates.

Island Territories Highlights: Opportunities for Improvement



Appropriate Use of Breastfeeding Supplements

None (0%) of Island Territories facilities adhere to standard clinical practice guidelines against routine supplementation with formula, glucose water, or water.

The American Academy of Pediatrics (AAP) and the American College of Obstetricians and Gynecologists (ACOG) Guidelines for Perinatal Care recommend against routine supplementation because supplementation with formula and/or water makes infants more likely to receive formula at home and stop breastfeeding prematurely.



Inclusion of Model Breastfeeding Policy Elements

Only 17% of Island Territories facilities have comprehensive breastfeeding policies including all model breastfeeding policy components recommended by the Academy of Breastfeeding Medicine (ABM).

The ABM model breastfeeding policy elements are the result of extensive research on best practices to improve breastfeeding outcomes. Facility policies determine the nature of care that is available to patients. Facilities with comprehensive policies consistently have the highest rates of exclusive breastfeeding regardless of patient population characteristics such as ethnicity, income, and payer status.



Adequate Assessment of Staff Competency

Only 33% of Island Territories facilities annually assess staff competency for basic breastfeeding management and support.

Implementing comprehensive assessment of staff training and skills for basic breastfeeding management and support establishes the foundation for quality infant feeding care. Adequate training and skills assessment are critical to ensure that mothers and infants receive care that is consistent, evidence-based, and appropriate.



Initiation of Mother and Infant Skin-to-Skin Care

Only 50% of Island Territories facilities initiate skin-to-skin care for at least 30 minutes upon delivery of the newborn. Upon delivery, the newborn should be placed skin-to-skin with the mother and allowed uninterrupted time to initiate and establish breastfeeding in order to improve infant health outcomes and reduce the risk of impairment of the neonatal immune system from unnecessary non-breast milk feeds.



Breastfeeding is a public health priority.



Breastfeeding is associated with decreased risk for infant morbidity and mortality as well as maternal morbidity,¹ and provides optimal infant nutrition. *Healthy People 2020*² establishes breastfeeding initiation, continuation, and exclusivity as national priorities.

Changes in maternity care practices improve breastfeeding rates.

There are many opportunities to protect, promote, and support breastfeeding in Island Territories. Opportunities such as those listed below can help bring ideal maternity care practices to all Island Territories hospitals.

Change opportunities:

- Examine Island Territories regulations for maternity facilities and evaluate their evidence base.
- Sponsor an Island Territories-wide summit of key decision-making staff at maternity facilities to highlight the importance of evidence-based practices for breastfeeding.
- Encourage and support hospital staff across Island Territories to be trained in providing care that supports mothers to breastfeed.
- Establish links among maternity facilities and community breastfeeding support networks in Island Territories.
- Implement evidence-based practices in medical care settings across Island Territories that support mothers' efforts to breastfeed.
- Integrate maternity care into related hospital-wide Quality Improvement efforts across Island Territories.
- Promote utilization of the Joint Commission's Perinatal Care Core Measure Set including exclusive breast milk feeding at hospital discharge in Island Territories hospital data collection systems.

Questions about the mPINC survey?

Information about the mPINC survey, results, reports, scoring, and history is at: www.cdc.gov/mpinc

For more information:

Centers for Disease Control and Prevention Division of Nutrition, Physical Activity, and Obesity Atlanta, GA USA

November 2014

Island Territories' 2013 Survey Results

73	Island Territories' State mPINC Score	Island Territories State mPINC Ran

Dimension Subscore* Ideal Response to mPINC Survey Question Response Response		(out of 53)'	(out of 100)*
Initial skin-to-skin contact is at least 30 min w/in 2 hours (cesarean births) Initial breastfeeding opportunity is w/in 1 hour (vaginal births) Initial breastfeeding opportunity is w/in 2 hours (cesarean births) Routine procedures are performed skin-to-skin Initial feeding is breast milk (vaginal births) Initial feeding is breast milk (vaginal births) Initial feeding is breast milk (cesarean births) Supplemental feedings to breastfeeding infants are rare Water and glucose water are not used Infant feeding decision is documented in the patient chart Staff provide breastfeeding advice & instructions to patients Staff teach breastfeeding cues to patients Staff teach patients not to limit suckling time Staff directly observe & assess breastfeeding 50	em ank [†]	Facilities with Ideal	mPINC Care Dimension
Tabor and Delivery Care	50	50	Initial skin-to-skin contact is at least 30 min w/in 1 hour (vaginal births)
Third breastfeeding opportunity is w/in 2 hours (cesarean births) 20	52	20	Initial skin-to-skin contact is at least 30 min w/in 2 hours (cesarean births)
Initial breastfeeding opportunity is w/in 2 hours (cesarean births) 20	24	67	
Feeding of Breastfed Infants Breastfed Infants Breastfed Infants Initial feeding is breast milk (vaginal births) Supplemental feedings to breastfeeding infants are rare Water and glucose water are not used Infant feeding decision is documented in the patient chart Staff provide breastfeeding advice & instructions to patients Staff teach breastfeeding cues to patients Staff teach patients not to limit suckling time Staff directly observe & assess breastfeeding 50	51	20	
Feeding of Breastfed Infants Breastfed Infants Initial feeding is breast milk (cesarean births) Supplemental feedings to breastfeeding infants are rare Water and glucose water are not used Infant feeding decision is documented in the patient chart Staff provide breastfeeding advice & instructions to patients Staff teach breastfeeding cues to patients 67 Staff teach patients not to limit suckling time Staff directly observe & assess breastfeeding 50	15	50	Routine procedures are performed skin-to-skin
Breast-fed Infants Supplemental feedings to breastfeeding infants are rare O Water and glucose water are not used Infant feeding decision is documented in the patient chart Staff provide breastfeeding advice & instructions to patients Staff teach breastfeeding cues to patients Staff teach patients not to limit suckling time Staff directly observe & assess breastfeeding 50	19	83	Initial feeding is breast milk (vaginal births)
Water and glucose water are not used 100 Infant feeding decision is documented in the patient chart 100 Staff provide breastfeeding advice & instructions to patients 50 Staff teach breastfeeding cues to patients 67 Staff teach patients not to limit suckling time 17 Staff directly observe & assess breastfeeding 50	53	20	Feeding of Initial feeding is breast milk (cesarean births)
Infant feeding decision is documented in the patient chart 100 Staff provide breastfeeding advice & instructions to patients 50 Staff teach breastfeeding cues to patients 67 Staff teach patients not to limit suckling time 17 Staff directly observe & assess breastfeeding 50	53	0	Infants Supplemental feedings to breastfeeding infants are rare
Staff provide breastfeeding advice & instructions to patients 50 Staff teach breastfeeding cues to patients 67 Staff teach patients not to limit suckling time 17 Staff directly observe & assess breastfeeding 50		100	Water and glucose water are not used
Breast-feeding Assistance Staff teach breastfeeding cues to patients Staff teach patients not to limit suckling time Staff directly observe & assess breastfeeding 50		100	Infant feeding decision is documented in the patient chart
Freast-feeding Assistance Staff teach patients not to limit suckling time 17 Staff directly observe & assess breastfeeding 50	53	50	Staff provide breastfeeding advice & instructions to patients
feeding Assistance Staff teach patients not to limit suckling time 17 Staff directly observe & assess breastfeeding 50	53	67	Staff teach breastfeeding cues to patients
Staff directly observe & assess breastfeeding 50	51	17	feeding Staff teach patients not to limit suckling time
Staff use a standard feeding assessment tool 33	53	50	Staff directly observe & assess breastfeeding
	52	33	Staff use a standard feeding assessment tool
Staff rarely provide pacifiers to breastfeeding infants 83	3	83	Staff rarely provide pacifiers to breastfeeding infants
Mother-infant pairs are not separated for postpartum transition 33	52	33	Mother-infant pairs are not separated for postpartum transition
Contact Mother-infant pairs room-in at night 100		100	Contact Mother-infant pairs room-in at night
Between Mother and Mother infant pairs are not separated during the hospital stay 100		100	
Infant Infant procedures, assessment, and care are in the patient room 50	1	50	
Non-rooming-in infants are brought to mothers at night for feeding 100		100	Non-rooming-in infants are brought to mothers at night for feeding
Facility Discharge Staff provide appropriate discharge planning (referrals & other multi-modal support) 83	1	83	Facility Staff provide appropriate discharge planning (referrals & other multi-modal support)
	12	83	Discharge packs containing infant formula samples and marketing
New staff receive appropriate breastfeeding education 0	53	0	New staff receive appropriate breastfeeding education
Staff Current staff receive appropriate breastfeeding education 17	40	17	Staff Current staff receive appropriate breastfeeding education
	53	0	8 1 7
Assessment of staff competency in breastfeeding management & support is at least annual	52	33	
Breastfeeding policy includes all 10 model policy elements 17	39	17	Breastfeeding policy includes all 10 model policy elements
Breastfeeding policy is effectively communicated 83	15	83	Breastfeeding policy is effectively communicated
Structural &	53	50	Structural &
Organizational Aspects of Facility provides breastfeeding support to employees 33	52	33	
C 'D !!	16	33	
	43	83	
Facility has a designated staff member responsible for coordination of lactation care	50	50	

^{*} Quality Practice scores range from 0 to 100 for each question, dimension of care, facility, and state. The highest, best possible score for each is 100. Each facility and state's "Total Score" is made up of subscores for practices in each of 7 dimensions of care.

References

[†] Ranks range from 1 to 53, with 1 being the highest rank. In case of a tie, both are given the same rank. State ranks are not shown for survey questions with 90% or more facilities reporting ideal responses.

¹ Ip S, Chung M, Raman G, et al. Breastfeeding and maternal and infant health outcomes in developed countries. Rockville, MD: US Dept of Health and Human Services, Agency for Healthcare Research and Quality; 2007.
² US Dept of Health and Human Services. Healthy People 2020 Summary of Objectives: Maternal, Infant, and Child Health. Available at http://www.healthypeople.gov/2020/topicsobjectives2020/pdfs/MaternalChildHealth.pdf

³DiGirolamo AM, Grummer-Strawn LM, Fein S. Effect of maternity care practices on breastfeeding. Pediatrics 2008;122, Supp 2:S43-9

Fairbank L, O'Meara S, Renfrew MJ, Woolridge M, Snowden AJ, Lister-Sharp D. A systematic review to evaluate the effectiveness of interventions to promote the initiation of breastfeeding. Health Technology Assessment 2000;4:1-171.