



# Guide to Hand Hygiene Opportunities in Hemodialysis

Hand hygiene opportunity category	Specific examples
1. Prior to touching a patient	<ul style="list-style-type: none"> <li>• Prior to entering station to provide care to patient</li> <li>• Prior to contact with vascular access site</li> <li>• Prior to adjusting or removing cannulation needles</li> </ul>
2. Prior to aseptic procedures	<ul style="list-style-type: none"> <li>• Prior to cannulation or accessing catheter</li> <li>• Prior to performing catheter site care</li> <li>• Prior to parenteral medication preparation</li> <li>• Prior to administering IV medications or infusions</li> </ul>
3. After body fluid exposure risk	<ul style="list-style-type: none"> <li>• After exposure to any blood or body fluids</li> <li>• After contact with other contaminated fluids (e.g., spent dialysate)</li> <li>• After handling used dialyzers, blood tubing, or prime buckets</li> <li>• After performing wound care or dressing changes</li> </ul>
4. After touching a patient	<ul style="list-style-type: none"> <li>• When leaving station after performing patient care</li> <li>• After removing gloves</li> </ul>
5. After touching patient surroundings	<ul style="list-style-type: none"> <li>• After touching dialysis machine</li> <li>• After touching other items within dialysis station</li> <li>• After using chairside computers for charting</li> <li>• When leaving station</li> <li>• After removing gloves</li> </ul>

Please make note of the following during this session.			
	Yes	No	Comments
There is a sufficient supply of alcohol-based hand sanitizer			
There is a sufficient supply of soap at handwashing stations			
There is a sufficient supply of paper towels at handwashing stations			
There is visible and easy access to hand washing sinks or hand sanitizer			

