Facility N	lame:					Obse	erver:				
Date(s):_				Location	of Medicat	ion Prepara	tion:				_
						lication p		ON a "Φ" if not pe	rformed/pe	erformed inco	orrectly. If
				•		•	• •	procedure to b	• •		•
Day (i.e., M, Tu, W)	Shift (i.e., 1 - 4)	Discipline	Med prep done in designated area	Med prep area is clean *	All vial(s) are inspected **	Hand hygiene performed	all vial(s)	All vials entered with new needle and new syringe	Med prep done aseptically	All single dose vial(s) discarded	All multi dose vial(s) discarded or stored properly
·			urse, T =tech BSERVATION		udent, O =			ions performe ber of sessions			

^{**}Vial should be discarded if sterility is questionable, or expiration date or beyond-use date has been exceeded. If a multi-dose vial will not be immediately discarded after use, the vial should be labeled upon opening to indicate the beyond-use date.



^{*}Preparation of injectable medications must be performed in a designated clean area that is free of obvious contamination sources (e.g., blood, body fluids, contaminated equipment, tap water).

Facility Na	me:		Observer:								
Date:		Da	y: M W F 7	Tu Th Sa	Shift: 1 st 2 nd 3 rd 4	th Start tii	Start time:				
(Use a "√" i	f action perform	ed correctly, a	"Ф" if not perf	edication adr formed/performed to be counted as s	ed incorrectly. If n	ot observed, le	ave blank.	All applicable			
Discipline	Medication properly transported to patient station*	Hand hygiene performed	Clean gloves worn	Injection port disinfected with antiseptic**	Medication administered aseptically	Syringe discarded at point of use	Gloves removed	Hand hygiene performed			
•	P=physician, N =nu observation perio			Numbe	er of procedures pe of procedures obs						
ADDITIONA	L COMMENTS/OF	BSERVATIONS:									



^{*} Medications should be transported directly from medication preparation area to individual patient. Medications should be prepared as close as possible to the time of medication administration. Medications that are not immediately administered by the person who prepared the medication must be labeled appropriately.

**Appropriate antiseptics are chlorhexidine, povidone-iodine, tincture of iodine, and 70% alcohol.