

Large-Scale Survey for Tickborne Bacteria, Khammouan Province, Laos

Andrew J. Taylor, Khamsing Vongphayloth, Malavanh Vongsouvath, Marc Grandadam, Paul T. Brey, Paul N. Newton, Ian W. Sutherland,¹ Sabine Dittrich¹

We screened 768 tick pools containing 6,962 ticks from Khammouan Province, Laos, by using quantitative real-time PCR and identified *Rickettsia* spp., *Ehrlichia* spp., and *Borrelia* spp. Sequencing of *Rickettsia* spp.–positive and *Borrelia* spp.–positive pools provided evidence for distinct genotypes. Our results identified bacteria with human disease potential in ticks in Laos.

Rickettsia, *Borrelia*, *Ehrlichia*, *Anaplasma*, and *Coxiella* spp. are tick-associated bacteria and well-described human pathogens. All of these bacteria, except *Coxiella* spp., are primarily transmitted through tick bites and cause febrile disease with a wide spectrum of severity. Tickborne bacterial pathogens are believed to be an underrecognized cause of acute febrile illness in Southeast Asia (1).

In Laos, spotted fever group *Rickettsia* have been shown to cause undifferentiated fever in 2% of febrile hospitalized adult patients (2). However, data on bacteria in ticks in Laos are sparse. To date, 1 *Rickettsia* sp. has been identified in a *Boophilus* sp. tick from Luang Namtha Province; this species showed 99.8% similarity with the *Rickettsia* sp. FUJ98 *ompA* gene (3). No other tickborne bacteria have been reported from Laos. Therefore, we investigated *Rickettsia*, *Borrelia*, *Ehrlichia*, *Anaplasma*, and *Coxiella* spp. in ticks from Khammouan Province, Laos.

The Study

We collected ticks in Nakai District, Khammouan Province, during the dry seasons (December–April) during 2012–2014, as previously described (4) (online Technical Appendix Figures 1, 2, <http://wwwnc.cdc.gov/EID/article/22/9/15-1969-Techapp1.pdf>). A total of 6,692 ticks were pooled ($n = 768$ pools, 1–10 ticks/pool) according to genus, sex, developmental stage, collection period, and

site. One *Amblyomma testudinarium* nymph that contained a blood meal was processed separately.

We extracted DNA by using the NucleoSpin 8 Virus Extraction Kit (Macherey-Nagel, Düren, Germany). Pools were screened by using single quantitative real-time PCRs specific for *Rickettsia* spp. (17-kDa gene), *Borrelia* spp. (23S rRNA gene), *Anaplasma* spp. (major surface protein 2 gene), *Ehrlichia* spp. (16S rRNA gene), and *Coxiella* spp. (IS1111) (5–8) (online Technical Appendix Table 1). Five microliters of diluted (1:10) template containing 1× Platinum Supermix-UDG (Invitrogen, Carlsbad, CA, USA) and bovine serum albumin (40 mg/mL) were used for each assay. Positive and nontemplate controls were included in each run. Screening by PCR was performed once per sample. In concordance with published guidelines, results were considered positive if they had a cycle quantitation (C_q) value <40 and likely positive if they had a C_q value 40–45 (9).

Sequencing was attempted for pools with C_q values <40 (online Technical Appendix Table 2) and performed by Macrogen (Seoul, South Korea). Consensus sequences were analyzed by using CLC Main Workbench 7 (<http://www.clcbio.com/products/clc-main-workbench/>) and BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and submitted to GenBank. Phylogenetic trees were constructed by using the Kimura 2-parameter model and the neighbor-joining method. Bootstrap values were determined by using 1,000 replications.

A total of 768 tick pools containing 6,692 ticks were screened. Pools contained 3 genera of ticks: 59.9% (460/768) *Haemaphysalis* spp., 36.3% (279/768) *A. testudinarium*, and 3.8% (29/768) *Dermacentor auratus*. Of the pools, 3% (23/768) contained adults, 36.5% (280/768) contained larvae, and 60.5% (465/768) contained nymphs (Table 1).

Rickettsia spp. were identified in 5.7% (44/768) of pools, and an additional 2.3% (18/768) of pools were likely positive for *Rickettsia* spp. Sequences consistent with 5 described *Rickettsia* species or genotypes were identified: *R. tamurae*, *R. japonica*, *Rickettsia* sp. ATT, *Rickettsia* sp. Kagoshima6, and *Rickettsia* sp. TwKM01 (Table 2; Figure 1).

Three novel genotypes (Table 2) were identified that might be new species. *Candidatus Rickettsia laoensis* (pool 447) was identified in 1 *Haemaphysalis* sp. pool. Phylogenetic analysis of 2845–2920-bp concatenated sequences of *gltA*, *sca4*, and *ompB* genes suggested that this bacteria

Author affiliations: Mahosot Hospital, Vientiane, Laos (A.J. Taylor, M. Vongsouvath, P.N. Newton, S. Dittrich); University of Oxford, Oxford, UK (A.J. Taylor, P.N. Newton, S. Dittrich); Institut Pasteur du Laos, Vientiane (K. Vongphayloth, M. Grandadam, P.T. Brey, I.W. Sutherland); US Naval Medical Research Center–Asia, Sembawang, Singapore (I.W. Sutherland)

DOI: <http://dx.doi.org/10.3201/eid2208.151969>

¹These senior authors contributed equally to this article.

Table 1. Tick pools tested for bacteria after screening by quantitative PCR, Khammouan Province, Laos*

Bacteria and tick species	No. positive pools/no. tested (%)				
	Total	Larvae	Nymphs	Adult males	Adult females
<i>Rickettsia</i> spp.					
All	44/768 (5.7)	6/280 (2.1)	37/465 (8.2)	0/12 (0)	1/11 (9.1)
<i>Amblyomma testudinarium</i>	27/279 (10.0)	0/61 (0)	27/217 (12.9)	0/1 (0)	0/1 (0)
<i>Haemaphysalis G1</i>	5/398 (3.8)	6/194 (3.1)	9/200 (4.5)	0/3 (0)	0/1 (0)
<i>H. hystricis</i>	1/6 (16.7)	NS	NS	0/3 (0)	1/3 (33.3)
<i>Dermacentor auratus</i>	1/29 (3.4)	0/0 (0)	1/26 (3.8)	0/2 (0)	0/1 (0)
<i>Ehrlichia</i> spp.					
All	12/768 (1.6)	4/280 (1.4)	6/465 (1.3)	1/12 (8.3)	1/11 (9.1)
<i>A. testudinarium</i>	2/279 (0.7)	0/61 (0)	2/217 (0.9)	0/1 (0)	0/1 (0)
<i>Haemaphysalis G1</i>	8/398 (2.0)	4/194 (2.1)	4/200 (2.0)	0/3 (0)	0/1 (0)
<i>H. aborensis</i>	2/6 (33.3)	NS	NS	1/3 (33.3)	1/3 (33.3)
<i>Borrelia</i> spp.					
All	12/768 (1.6)	2/280 (0.7)	8/465 (1.7)	2/12 (16.7)	NS
<i>A. testudinarium</i>	2/279 (0.7)	1/61 (1.6)	1/217 (0.5)	0/1 (0)	0/1 (0)
<i>Haemaphysalis G1</i>	6/398 (1.5)	1/194 (0.5)	5/200 (2.5)	0/3 (0)	0/1 (0)
<i>Haemaphysalis G1.2</i>	1/13 (7.7)	NS	1/13 (7.7)	NS	NS
<i>H. aborensis</i>	2/6 (33.3)	NS	NS	2/3 (66.7)	0/3 (0)
<i>D. auratus</i>	1/29 (3.4)	0/0 (0)	1/26 (3.8)	0/2 (0)	0/1 (0)
<i>Coxiella</i> spp.					
All	5/511 (1.0)†	4/187 (2.1)†	1/310 (0.3)	0/8 (0)	0/6 (0)
<i>Haemaphysalis G1</i>	5/279 (1.8)†	4/162 (2.5)†	1/117 (0.9)	NS	NS
<i>Anaplasma</i> spp.					
All	2/768 (0.3)†	0/280 (0)†	0/465 (0)†	0/12 (0)	0/11 (0)
<i>A. testudinarium</i>	1/279 (0.4)†	0/61 (0)	1/217 (0.5)†	0/1 (0)	0/1 (0)
<i>Haemaphysalis G1</i>	1/398 (0.3)†	1/194 (0.5)†	0/200 (0)	0/3 (0)	0/1 (0)

*NS, no samples were available for screening.

†Includes samples with cycle quantitation values <40 and 40–45.

belonged to the *R. massiliae* group of rickettsiae (online Technical Appendix Figure 3). *Candidatus* Rickettsia mahosotii (pools 81 and 372) was identified in *Haemaphysalis* spp. and *A. testudinarium* pools. Phylogenetic analysis of *gltA*, *sca4*, and *ompB* genes suggested that this bacteria belonged to the *R. rickettsii* group (online Technical Appendix Figure 3). *Candidatus* Rickettsia khammouanensis was identified in 1 *Haemaphysalis* sp. nymph pool (pool 120). Phylogenetic analysis of *gltA*, 17-kDa, and *ompB* genes suggested a relationship with the *R. helvetica* group (online Technical Appendix Figure 4).

In addition, 15 *A. testudinarium* pools showed dual peaks for 17-kDa gene sequences, which suggested the presence of *R. tamurae* and *Rickettsia* sp. ATT. Sequencing of *sca4*, *ompA*, and *ompB* genes from 1 of these pools (pool 239) identified unique sequences (Table 2; online Technical Appendix Figure 4).

Borrelia spp. were identified in 1.6% (12/768) of pools (Table 1). Two unique sequences obtained from *Haemaphysalis* spp. pools showed 99.3% (298/300) (GenBank accession no. KR733069) and 98.7% (296/300) (accession no. KR733068) identity with Shiretoko *Haemaphysalis* *Borrelia* sp. (AB897888). Phylogenetic analysis confirmed that both bacteria were closely related to Shiretoko *Haemaphysalis* *Borrelia* sp. (accession no. B897888) and belong to the relapsing fever group of *Borrelia* (Figure 2).

Twelve (1.6%) of 768 pools were positive for *Ehrlichia* spp. (Table 1); an additional 6 pools (0.8%)

were likely positive. One short sequence from a *Haemaphysalis* sp. nymph pool (pool 357) was obtained, and this sequence showed 100% identity (116/116 bases) with the genus *Ehrlichia*.

No pools were positive for *Anaplasma* spp., but 2 were likely positive (Table 1). Although not all pools were tested for *Coxiella* spp. (n = 511), 1 pool (0.2%) was positive, and 4 pools were likely positive for *C. burnetti*. No confirmatory sequences were obtained from these pools. The 1 tick that contained a blood meal (*A. testudinarium* nymph) showed negative results by screening PCRs.

Conclusions

This study provides evidence that *Rickettsia* spp., *Borrelia* spp., and *Ehrlichia* spp. are present in ticks in Laos. Several *Rickettsia* spp. identified in this study are human pathogens. Infections with *R. tamurae* (2) and *R. japonica* are well described in Southeast Asia (10). However, the pathogenicity of *Rickettsia* sp. TwkM01 (11), *Rickettsia* sp. ATT (12), *Rickettsia* sp. kagoshima6 genotypes (13) and potential novel *Candidatus* Rickettsia laoensis, *Candidatus* Rickettsia mahosotii, and *Candidatus* Rickettsia khammouanensis is unknown. *Candidatus* Rickettsia khammouanensis is phylogenetically related to *R. helvetica*, for which there is serologic evidence for its role as a human pathogen in Laos (2). Unique *ompA*, *ompB*, and *sca4* sequences identified in this study (Table 2) might indicate the presence of

Table 2. Sequence data for *Rickettsia* species isolated from ticks, Khammouan Province, Laos*

Tick pool	Tick species and stage	<i>Rickettsia</i> spp. gene, GenBank accession no., and % similarity (no. matching nucleotides/total)				
		17-kDa	<i>gltA</i>	<i>sca4</i>	<i>ompA</i>	<i>ompB</i>
110	<i>Amblyomma testudinarium</i> nymph	Unclear sequence	NS	Unclear sequence	KT753264, 100.0 (529/529) with <i>Rickettsia</i> sp. TwKM01 EF219467	NS
177, 180, 216, 220	<i>A. testudinarium</i> nymph	KR733070, 100.0 (355/355) with <i>R. tamurae</i> AB114825	KT753265, 99.8 (1,096/1,098) with <i>R. tamurae</i> AB812551	KT753266, 99.7 (607/609) with <i>R. tamurae</i> DQ113911	NS	NS
315	<i>A. testudinarium</i> nymph	KT753267, 98.8 (407/412) with <i>R. raoultii</i> JX885457	KT753268, 99.9 (1,036/1,037) with <i>Rickettsia</i> kagoshima6 JQ697956	KT753269, 96.8 (795/821) with <i>Rickettsia</i> sp. AUS 118, KF666473	Could not be amplified	KT753270, 95.0 (1,073/1,129) with <i>R. massiliae</i> CP003319
239	<i>A. testudinarium</i> nymph	KT753271, 99.7 (360/361) with <i>Rickettsia</i> sp. ATT AF483196	KT753272, 99.7 (1,048/1,051) with <i>R. tamurae</i> (AB812551)/KT753273; 99.2 (367/370) with <i>Rickettsia</i> sp. hhmj7 KC566999	KT753274, 97.1 (759/782) with <i>Rickettsia</i> sp. AUS 118 KF666473	KT753275, 87.2 (530/602) with <i>R. raoultii</i> JQ792137	KT753276, 97.5 (1,052/1,079) with <i>R. massiliae</i> CP003319
76, 337, 450, 453	<i>Haemaphysalis</i> G1 nymphs (3), <i>A. testudinarium</i> nymph (1)	KT753277, 98.4 (417/423) with <i>R. raoultii</i> JX885457	KT753278, 99.9 (1,037/1,038) with <i>Rickettsia</i> sp. kagoshima6 JQ697956	KT753279, 98.4 (794/807) with <i>R. japonica</i> AF155055	Could not be amplified	KT753280, 96.0 (410/427) with <i>R. raoultii</i> EU036984
81, 372	<i>Haemaphysalis</i> G1 nymphs, <i>A. testudinarium</i> nymph (17 kDa only)	KT753283, 99.0 (408/412) with <i>R. raoultii</i> JX885457	KT753284, 99.5 (1,090/1,096) with <i>R. sibirica</i> U59734	KT753285, 98.5 (838/851) with <i>R. japonica</i> AF155055	Could not be amplified	KT753286, 97.7 (1,118/1,144) with <i>R. massiliae</i> CP003319
120	<i>Haemaphysalis</i> G1 nymph	KT753287, 96.1 (391/407) with <i>R. helvetica</i> GU827073	KT753288, 97.1 (370/381) with <i>Candidatus</i> <i>Rickettsia</i> rara DQ365805	Could not be amplified (x2)	Could not be amplified (x2)	KT753289, 86.4 (362/419), <i>R. aeschlimannii</i> AF123705
407	<i>Haemaphysalis hysticis</i> adult	KR733074, 100.0 (413/413), <i>R. japonica</i> AP011533	KT753281, 100.0 (1,063/1,063), <i>R. japonica</i> AP011533	NS	NS	KT753282, 100.0 (1,191/1,191) with <i>R. japonica</i> AP011533
447	<i>Haemaphysalis</i> G1 nymph	KT753291, 98.6 (407/413) with <i>R. massiliae</i> CP000683	KT753290, 99.6 (961/965) with <i>R. raoultii</i> JX885455	KT753292, 97.5 (809/830) with <i>Rickettsia</i> sp. AUS118 KF66473	KT753293, 97.5 (591/606) with <i>Rickettsia</i> sp. JL-02 AY093696	KT753294, 98.4 (1,137/1,156), with <i>R. massiliae</i> CP003319

*New sequences were compared with reference sequences. NS, not sequenced.

Rickettsia sp. ATT (12), which was previously believed to be identical to *R. tamurae* (14), and suggests that it might be a distinct species. Further studies, including whole-genome sequencing, are required to identify and confirm these novel genotypes and understand their role in human disease.

Borrelia spp. sequences identified in *Haemaphysalis* spp. pools were shown to have high concordance with the Shiretoko *Haemaphysalis* *Borrelia* isolated from *Haemaphysalis* spp. ticks and deer in Japan (15). The species belongs to the relapsing fever group of *Borrelia* and is related to *B. lonestari*.

Sequence data for *Ehrlichia* spp. indicated the presence of these bacteria but were not sufficient to identify

them to the species level. The C_q values were high (40–45) for *Anaplasma* spp., but no sequence data were obtained. *Coxiella* spp. were screened by using primers for IS1111, which are not specific for *C. burnetii*, and no confirmatory sequence data were obtained. Because of limited reagents, screening of all 768 pools for *Coxiella* spp. was not completed. Further work is required to investigate the presence of these bacteria in Laos.

Our study had several limitations. First, pooling of ticks precludes an accurate assessment of prevalence of bacterial pathogens. Second, sequences obtained from some *A. testudinarium* pools had dual peaks, suggestive of multiple infections, and could therefore not be interpreted. Third, ticks were collected only from 1 area in Laos




Figure 1. Phylogenetic analysis of *Rickettsia* spp. in ticks, Khammouan Province, Laos. The tree was constructed by using partial nucleotide sequences (350 bp) of the 17-kDa gene, the Kimura 2-parameter model, and the neighbor-joining method. Analyses were supported by bootstrap analysis with 1,000 replications. Numbers along branches are bootstrap values. GenBank accession numbers are shown for reference sequences. Sample numbers for each tick are shown in parentheses. Scale bar indicates nucleotide substitutions per site.




Figure 2. Phylogenetic analysis of *Borrelia* spp. in ticks, Khammouan Province, Laos. The tree was constructed by using partial nucleotide sequences (299–323 bp) of the *flaB* gene, the Kimura 2-parameter model, and the neighbor-joining method. Analyses were supported by bootstrap analysis with 1,000 replications. Numbers along branches are bootstrap values. GenBank accession numbers are shown for reference sequences. Sample numbers for each tick are shown in parentheses. Scale bar indicates nucleotide substitutions per site.

(Khammouan Province); thus, extrapolating findings to the entire country must be done cautiously.

Our results highlight the frequency of tickborne bacterial infections in Laos. These findings emphasize the need for further research of tick-associated bacteria and their role in human disease.

Acknowledgments

We thank the staff of Mahosot Hospital, especially Soulignasack Thongpaseuth, for providing technical assistance, and Al Richards and Ju Jiang for fruitful discussions.

This study was supported by the US Naval Medical Research Center–Asia in support of the Department of Defense Global

Emerging Infections Surveillance Program, the Institut Pasteur du Laos, and the Wellcome Trust of Great Britain.









Dr. Taylor is a research physician at the Centre for Tropical Medicine and Global Health, University of Oxford, Oxford, UK. His primary research interest is infectious diseases.

References

1. Blacksell SD, Kantipong P, Watthanaworawit W, Turner C, Tanganuchitcharnchai A, Jintawon S, et al. Underrecognized arthropod-borne and zoonotic pathogens in northern and northwestern Thailand: serological evidence and opportunities for awareness. *Vector Borne Zoonotic Dis.* 2015;15:285–90. <http://dx.doi.org/10.1089/vbz.2015.1776>
2. Phongmany S, Rolain JM, Phetsouvanh R, Blacksell SD, Soukkhaseum V, Rasachack B, et al. Rickettsial infections and fever, Vientiane, Laos. *Emerg Infect Dis.* 2006;12:256–62. <http://dx.doi.org/10.3201/eid1202.050900>
3. Kernif T, Socolovschi C, Wells K, Lakim MB, Inthalad S, Slesak G, et al. *Bartonella* and *Rickettsia* in arthropods from the Lao PDR and from Borneo, Malaysia. *Comp Immunol Microbiol Infect Dis.* 2012;35:51–7. <http://dx.doi.org/10.1016/j.cimid.2011.10.003>
4. Vongphayloth K, Brey PT, Robbins RG, Sutherland IW. First survey of the hard tick (*Acari: Ixodidae*) fauna of Nakai District, Khammouane Province, Laos, and an updated checklist of the ticks of Laos. *Systematic and Applied Acarology.* 2016;21:166–80. <http://dx.doi.org/10.11158/saa.21.2.2>
5. Wright CL, Nadolny RM, Jiang J, Richards AL, Sonenshine DE, Gaff HD, et al. *Rickettsia parkeri* in gulf coast ticks, southeastern Virginia, USA. *Emerg Infect Dis.* 2011;17:896–8. <http://dx.doi.org/10.3201/eid1705.101836>
6. Courtney JW, Kostelnik LM, Zeidner NS, Massung RF. Multiplex real-time PCR for detection of *Anaplasma phagocytophilum* and *Borrelia burgdorferi*. *J Clin Microbiol.* 2004;42:3164–8. <http://dx.doi.org/10.1128/JCM.42.7.3164-3168.2004>
7. Fournier PE, Thuny F, Richet H, Lepidi H, Casalta JP, Arzouni JP, et al. Comprehensive diagnostic strategy for blood culture-negative endocarditis: a prospective study of 819 new cases. *Clin Infect Dis.* 2010;51:131–40. <http://dx.doi.org/10.1086/653675>
8. Loftis AD, Massung RF, Levin ML. Quantitative real-time PCR assay for detection of *Ehrlichia chaffeensis*. *J Clin Microbiol.* 2003;41:3870–2. <http://dx.doi.org/10.1128/JCM.41.8.3870-3872.2003>
9. Bustin SA, Benes V, Garson JA, Hellems J, Huggett J, Kubista M, et al. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem.* 2009;55:611–22. <http://dx.doi.org/10.1373/clinchem.2008.112797>
10. Takada N, Fujita H, Kawabata H, Ando S, Sakata A, Takano A, et al. Spotted fever group *Rickettsia* sp. closely related to *Rickettsia japonica*, Thailand. *Emerg Infect Dis.* 2009;15:610–1.
11. Tsui PY, Tsai KH, Weng MH, Hung YW, Liu YT, Hu KY, et al. Molecular detection and characterization of spotted fever group rickettsiae in Taiwan. *Am J Trop Med Hyg.* 2007;77:883–90.
12. Hirunkanokpun S, Kittayapong P, Cornet J-P, Gonzalez J-P. Molecular evidence for novel tick-associated spotted fever group rickettsiae from Thailand. *J Med Entomol.* 2003;40:230–7. <http://dx.doi.org/10.1603/0022-2585-40.2.230>
13. Gaowa, Ohashi N, Aochi M, Wurutu D, Wu, Yoshikawa Y, et al. Rickettsiae in ticks, Japan, 2007–2011. *Emerg Infect Dis.* 2013;19:338–40. <http://dx.doi.org/10.3201/eid1902.120856>
14. Fournier PE, Takada N, Fujita H, Raoult D. *Rickettsia tamurae* sp. nov., isolated from *Amblyomma testudinarium* ticks. *Int J Syst Evol Microbiol.* 2006;56:1673–5. <http://dx.doi.org/10.1099/ijs.0.64134-0>
15. Lee K, Takano A, Taylor K, Sashika M, Shimozuru M, Konnai S, et al. A relapsing fever group *Borrelia* sp. similar to *Borrelia lonestari* found among wild sika deer (*Cervus nippon yesoensis*) and *Haemaphysalis* spp. ticks in Hokkaido, Japan. *Ticks Tick Borne Dis.* 2014;5:841–7. <http://dx.doi.org/10.1016/j.tbd.2014.06.006>

Address for correspondence: Andrew J. Taylor; Center for Tropical Medicine and Global Health, Nuffield Department of Medicine, Research Building, University of Oxford, Oxford OX3 7FZ, UK; email: andrewtaylor9@gmail.com

WORLD HEALTH DAYS

 world AIDS day december 1	 HOW SAFE IS YOUR FOOD? world health day	 WORLD RABIES DAY SEPTEMBER 28 world rabies day september 28
 WORLD HEPATITIS DAY world hepatitis day july 28	 WORLD IMMUNIZATION WEEK	 Find TB. Treat TB. Working together to eliminate TB. world TB day march 24
 WORLD MALARIA DAY world malaria day april 25	 world pneumonia day november 12	Visit the World Health Days section on our website for the latest articles and information on emerging infectious diseases in our global community.
http://wwwnc.cdc.gov/eid/page/world-health-days		

Large-Scale Survey for Tickborne Bacteria, Khammouan Province, Laos

Technical Appendix

Technical Appendix Table 1. Primers and probes used for screening quantitative PCR in large-scale survey for tickborne bacteria, Khammouan Province, Laos*

Bacteria	Gene	Primers and probes used to screen samples			Reference†
		Primer or probe	Sequence, 5'→3'		
<i>Rickettsia</i> spp.	17 kDa	R17K128F2	F-GGGCGGTATGAAYAAACAAG	(5)	
Spotted fever group	17 kDa	R17K238R	R-CCTACACCTACTCCVACAAG	(5)	
Spotted fever group	17 kDa	R17K202TAQP	P-CCGAATTGAGAACCAAGTAATGC	(5)	
<i>Borrelia</i> spp.	23S rRNA	Bb23Sf	F-CGAGTCTTAAAAGGGCGATTTAGT	(6)	
<i>Borrelia</i> spp.	23S rRNA	Bb23Sr	R-TATTTATGGCCAGGCTGAAGC	(6)	
<i>Borrelia</i> spp.	23S rRNA	Bb23Sp	P-AGATGTGGTAGACCCGAAGCCGAGTG	(6)	
<i>Coxiella</i> spp.	IS1111	IS1111f	F-CAAGAAACGTATCGCTGTGGC	(7)	
<i>Coxiella</i> spp.	IS1111	IS1111R	R-CACAGAGCCACCGTATGAATC	(7)	
<i>Coxiella</i> spp.	IS1111	IS1111 probe	P-CCGAGTTCGAAACAATGAGGGCTG	(7)	
<i>Ehrlichia</i> spp.	16S rRNA	EHR16S-17	F-GCGGCAAGCCTAACACAT	(8)	
<i>Ehrlichia</i> spp.	16S rRNA	EHR16S-97	R-CCCGTCTGCCACTAACCAATTATT	(8)	
<i>Ehrlichia</i> spp.	16S rRNA	EHR16S-38	P-AGTCGAACGGACAATTGCTTATAACCTTTTGGT	(8)	
<i>Anaplasma</i> spp.	msp2	ApMSP2f	F-ATGGAAGGTAGTGTGGTTATGGTATT	(6)	
<i>Anaplasma</i> spp.	msp2	APMSP2r	R-TTGGTCTTGAAGCGCTCGTA	(6)	
<i>Anaplasma</i> spp.	msp2	ApMSP2p	P-TGGTGCCAGGGTTGAGCTTGAGATTG	(6)	

*F, forward; IS, insertion sequence; msp2, major surface protein 2; P, probe; R, reverse.

†References are in the text of the article.

Technical Appendix Table 2. Primers used for sequencing in large-scale survey for tickborne bacteria, Khammouan Province, Laos*

Bacteria	Gene	Primers used to sequence bacteria-positive samples			
		Primer	Sequence, 5'→3'	Size, bp	Reference
<i>Rickettsia</i> spp.	17 kDa	R17kM61F†	F-ACTTTACAAAATTCTAAAAACCATATACT	524	(1)
Spotted fever group	17 kDa	R17K31F‡	F-CATTGTCCGTCAGGTTGGCG	524	(1)
Spotted fever group	17 kDa	Rr2608Rnew†‡	R-CATTGTCCGTCAGGTTGGCG	434	(1)
Spotted fever group	<i>gltA</i>	Cs1dF†‡	F-ATGACTAATGGCAATAATAA	1,237	(2)
Spotted fever group	<i>gltA</i>	RpE CS877p‡	F-CATAACCAAGTGTAAAGCTG	1,237	(2)
Spotted fever group	<i>gltA</i>	CS1273R†	R-GGGGGCCTGCTCACGGCGG	382	(2)
Spotted fever group	<i>gltA</i>	RpE CS1258n‡	R-ATTGCAAAAAGTACAGTGAACA	382	(2)
Spotted fever group	<i>sca4</i>	RrD749F†‡	F-TGGTAGCATTAAAAGCTGATGG	1,078	(3)
Spotted fever group	<i>sca4</i>	RrD928F‡	F-ATTTATACACTTGCGGTAACAC	1,078	(3)
Spotted fever group	<i>sca4</i>	RrD1826R†‡	R-TCTAAATKCTGCTGMATCAAT	899	(3)
Spotted fever group	<i>ompA</i>	RompAM50F†	R-TTGCGTTATAACACTTTTTAAGTGA	692	(4)
Spotted fever group	<i>ompA</i>	190–70F‡	F-ATGGCGAATATTTCTCCAAAA	692	(4)
Spotted fever group	<i>ompA</i>	RompA642R†	R-ATTACCTATTGTTCCGTTAATGGCA	632	(4)
Spotted fever group	<i>ompA</i>	190–701R‡	R-GTTCCGTTAATGGCAGCATCT	632	(4)
Spotted fever group	<i>ompB</i>	RompB11F†‡	F-ACCATAGTAGCMAGTTTTGCAG	1,902	(5)
Spotted fever group	<i>ompB</i>	120–607F‡	F-AATATCGCTGACGGTCAAGGT	1,902	(5)
Spotted fever group	<i>ompB</i>	RompB1902R†‡	R-CCGTCATTTCCAATAACTA	1,265	(5)
Spotted fever group	<i>ompB</i>	RAK1452R‡	R-SGTTAACTTKACCGYTTATAACTGT	1,452	(5)
<i>Borrelia</i> spp.	<i>fla B</i>	280F†	F-GCAGTTCARTCAGGTAACGG	1,452	(6)
<i>Borrelia</i> spp.	<i>fla B</i>	754R†	R-TAGCAAGTGATGTATTRGCATCAAC	475	(6)
<i>Borrelia</i> spp.	<i>fla B</i>	301F‡	F-ACATATTCAGATGCAGACAGAGG	475	(6)
<i>Borrelia</i> spp.	<i>fla B</i>	737R‡	R-GCATCAACTGRGTTGTAAACATTAACAGG	437	(6)
<i>Ehrlichia</i> spp.	16 S rRNA	Ehr16SF	F-GTACCYACAGAAGAAGTCC	437	(7)
<i>Ehrlichia</i> spp.	16 S rRNA	Ehr16SR	R-GCTGTAAACGATGAGTGCTAA	345	(7)


*F, forward; R, reverse.

†First reaction.


‡Second reaction.

References


1. Maina A. Sero-epidemiology and molecular characterization of Rickettsiae infecting humans, selected animals and arthropod vectors in Asembo, western Kenya, 2007–2010 [Doctoral dissertation]. Nairobi (Kenya): Jomo Kenyatta University of Agriculture and Technology; 2012.
2. Roux V, Rydkina E, Ereemeeva M, Raoult D. Citrate synthase gene comparison, a new tool for phylogenetic analysis, and its application for the rickettsiae. *Int J Syst Bacteriol.* 1997;47:252–61. [PubMed](#)
<http://dx.doi.org/10.1099/00207713-47-2-252>
3. Jiang J, Sangkasuwan V, Lerdthusnee K, Sukwit S, Chuenchitra T, Rozmajzl P, et al. Human infection with *Rickettsia honei* Thailand. *Emerg Infect Dis.* 2005;11:1473–5. [PubMed](#)
<http://dx.doi.org/10.3201/eid1109.050011>
4. Fournier PE, Roux V, Raoult D. Phylogenetic analysis of spotted fever group rickettsiae by study of the outer surface protein rOmpA. *Int J Syst Bacteriol.* 1998;48:839–49. [PubMed](#)
<http://dx.doi.org/10.1099/00207713-48-3-839>
5. Roux V, Raoult D. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (*ompB*). *Int J Syst Evol Microbiol.* 2000;50:1449–55. [PubMed](#)
<http://dx.doi.org/10.1099/00207713-50-4-1449>
6. Clark KL, Leydet B, Hartman S. Lyme borreliosis in human patients in Florida and Georgia, USA. *Int J Med Sci.* 2013;10:915–31. [PubMed](#) <http://dx.doi.org/10.7150/ijms.6273>
7. Dittrich S, Phuklia W, Turner GD, Rattanavong S, Chansamouth V, Dumler SJ, et al. *Neorickettsia sennetsu* as a neglected cause of fever in South-East Asia. *PLoS Negl Trop Dis.* 2015;9:e0003908. [PubMed](#)
<http://dx.doi.org/10.1371/journal.pntd.0003908>
8. Phongmany S, Rolain JM, Phetsouvanh R, Blacksell SD, Soukkhaseum V, Rasachack B, et al. Rickettsial infections and fever, Vientiane, Laos. *Emerg Infect Dis.* 2006;12:256–62. [PubMed](#)
<http://dx.doi.org/10.3201/eid1202.050900>
9. Kernif T, Socolovschi C, Wells K, Lakim MB, Inthalad S, Slesak G, et al. *Bartonella* and *Rickettsia* in arthropods from the Lao PDR and from Borneo, Malaysia. *Comp Immunol Microbiol Infect Dis.* 2012;35:51–7. [PubMed](#) <http://dx.doi.org/10.1016/j.cimid.2011.10.003>




Technical Appendix Figure 1. Khammouan Province (red star), Laos, where ticks were collected in this study (see Technical Appendix Figure 2). Locations of previous studies investigating *Rickettsia* spp. in Laos are shown by the green star (8) and blue star (9).



Technical Appendix Figure 2. Location of tick collections sites (red stars) in Khammouan Province, Laos.



Technical Appendix Figure 3. Phylogenetic analysis of *gltA*, *sca4*, and *ompB* genes of candidate novel *Rickettsia* spp., Kammouan Province, Laos. The tree was constructed by using concatenated partial nucleotide sequences (2,845–2,920 bp) of *gltA*, *sca4*, and *ompB* genes; the Kimura-80 model; and the neighbor-joining method. Analyses were supported by using bootstrap analysis with 1,000 replications. Numbers along branches are bootstrap values. Sample numbers identifying each tick pool from this study are shown in parentheses after the sequence name. Scale bar indicates nucleotide substitutions per site.



Technical Appendix Figure 4. Phylogenetic analysis of *gltA*, 17 kDa, and *ompB* genes of *Rickettsia* spp., Kammouan Province, Laos. The tree was constructed by using concatenated partial nucleotide sequences (1,114–1,117 bp) of concatenated sequences of *gltA*, 17-kDa, and *ompB* genes; the Kimura-80 model; and the neighbor-joining method. Analyses were supported by using bootstrap analysis with 1,000 replications. Numbers along branches are bootstrap values. Sample numbers identifying each tick pool from this study are shown in parentheses after the sequence name. Scale bar indicates nucleotide substitutions per site.