

About the Author

Dr. Farrington recently completed her pediatrics residency at the Children's Hospital Medical Center of Akron in 2022. She is currently working as a pediatric hospitalist at the St. Louis Children's Hospital-Washington University.

References

1. Kemenesi G, Bányai K. Tick-borne flaviviruses, with a focus on Powassan virus. *Clin Microbiol Rev.* 2018;32:e00106-17. <https://doi.org/10.1128/CMR.00106-17>
2. Centers for Disease Control and Prevention. Powassan virus [cited 2022 May 3]. <https://www.cdc.gov/powassan/statistics.html>
3. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215:403-10. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
4. Glaser CA, Honarmand S, Anderson LJ, Schnurr DP, Forghani B, Cossen CK, et al. Beyond viruses: clinical profiles and etiologies associated with encephalitis. *Clin Infect Dis.* 2006;43:1565-77. <https://doi.org/10.1086/509330>
5. Miller S, Naccache SN, Samayoa E, Messacar K, Arevalo S, Federman S, et al. Laboratory validation of a clinical metagenomic sequencing assay for pathogen detection in cerebrospinal fluid. *Genome Res.* 2019;29:831-42. <https://doi.org/10.1101/gr.238170.118>
6. Piantadosi A, Kanjilal S, Ganesh V, Khanna A, Hyle EP, Rosand J, et al. Rapid detection of Powassan virus in a patient with encephalitis by metagenomic sequencing. *Clin Infect Dis.* 2018;66:789-92. <https://doi.org/10.1093/cid/cix792>
7. Hermance ME, Thangamani S. Powassan virus: an emerging arbovirus of public health concern in North America. *Vector Borne Zoonotic Dis.* 2017;17:453-62. <https://doi.org/10.1089/vbz.2017.2110>
8. Eisen RJ, Eisen L, Beard CB. County-scale distribution of *Ixodes scapularis* and *Ixodes pacificus* (Acari: Ixodidae) in the continental United States. *J Med Entomol.* 2016;53:349-86. <https://doi.org/10.1093/jme/tjv237>
9. Bouchard C, Dibbernardo A, Koffi J, Wood H, Leighton PA, Lindsay LR. N increased risk of tick-borne diseases with climate and environmental changes. *Can Commun Dis Rep.* 2019;45:83-9 <https://doi.org/10.14745/ccdr.v45i04a02>
10. Diuk-Wasser MA, VanAcker MC, Fernandez MP. Impact of land use changes and habitat fragmentation on the eco-epidemiology of tick-borne diseases. *J Med Entomol.* 2021;58:1546-64. <https://doi.org/10.1093/jme/tjaa209>

Address for correspondence: Scott F Pangonis, Children's Medical Center of Akron, 1 Perkins Sq, Akron, OH 44321, USA; email: spangonis@akronchildrens.org

Rickettsia conorii Subspecies *israelensis* in Captive Baboons

Giovanni Sgroi, Roberta Iatta, Grazia Carelli, Annamaria Uva, Maria Alfonsa Cavalera, Piero Laricchiuta, Domenico Otranto

Author affiliations: Experimental Zooprophyllactic Institute of Southern Italy, Portici, Italy (G. Sgroi); University of Bari Aldo Moro, Bari, Italy (G. Sgroi, R. Iatta, G. Carelli, A. Uva, M.A. Cavalera, D. Otranto); Zoosafari Park Fasano, Brindisi, Italy (P. Laricchiuta); Bu-Ali Sina University, Hamedan, Iran (D. Otranto)

DOI: <https://doi.org/10.3201/eid2904.221176>

Hamadryas baboons (*Papio hamadryas*) may transmit zoonotic vector-borne pathogens to visitors and workers frequenting zoological parks. We molecularly screened 33 baboons for vector-borne pathogens. Three (9.1%) of 33 animals tested positive for *Rickettsia conorii* subspecies *israelensis*. Clinicians should be aware of potential health risks from spatial overlapping between baboons and humans.

Papio hamadryas baboons (order Primates, family Cercopithecidae) are frequently hosted in zoological gardens worldwide. The natural susceptibility of baboons to many zoonotic agents (1) may present a potential risk for transmission of emerging infectious diseases to humans. Nevertheless, few data are available on vector-borne pathogens of human concern that are hosted by baboons (e.g., *Rickettsia africae*, *Babesia microti*-like parasites, and *Anaplasma phagocytophilum*) (1). Data are likewise scarce on the role of *P. hamadryas* baboons in circulating arthropod vectors in zoological gardens and the resulting risk for transmitting vector-borne pathogens to persons frequenting such areas. We aimed to determine the occurrence of zoonotic vector-borne pathogens in a zoopark in the Apulia region of southern Italy and assess baboons' potential roles as reservoirs of emerging pathogens. Our study was approved by the University of Bari Aldo Moro ethics committee (Prot. Uniba 176/19).

During February–December 2020, we anesthetized baboons in the zoopark and housed them in cages for blood sampling. For each baboon, we recorded age, sex, weight, and body condition score (1–5); we obtained peripheral blood samples by cephalic vein puncture. To determine complete blood count and for molecular analysis, we collected 2 mL blood

samples in Vacutainer K3-EDTA tubes. For biochemical analysis, we collected an additional 5 mL blood in Vacutainer clot activator serum tubes and centrifuged (15 min at 1,500 × g at room temperature), then delivered it to the University of Bari Department of Veterinary Medicine (Bari, Italy). We extracted DNA using QIAGEN QIAamp DNA Blood and Tissue kits (<https://www.qiagen.com>) and molecularly tested for vector-borne pathogens (Table) (2–4). We purified and sequenced amplicons in both directions using a Big Dye Terminator v3.1 Cycle Sequencing Kit in an Applied Biosystems 3130 Genetic Analyzer (ThermoFisher, <https://www.thermofisher.com>), then edited and analyzed them using Geneious version 9.0 (<https://www.geneious.com>). We then compared resulting sequences with those in GenBank. We performed complete blood counts using CELL-DYN 3700 Hematology Analyzer (Abbott, <https://www.abbott.com>), biochemical profile using a KPM Analytics SAT 450 random access analyzer (<https://www.kpmanalytics.com>), and protein electrophoresis analyses using Sebia Hydrasys 2 Scan Focusing (<https://www.sebia.com>). We calculated 95% CIs for proportions and χ^2 and odds ratios (OR) to assess differences in prevalence and infection risk stratified by age and sex. We used *t*-tests to compare mean laboratory values between baboons positive and negative for vector-borne pathogens. We considered *p* values <0.05 statistically significant.

We included 33 baboons: 21 male, 12 female; 13 juvenile, 16 adult, and 4 elderly. Blood samples from 3/33 (9.1%, 95% CI 3.1%–23.4%; 1 adult male, 1 adult female, 1 juvenile male) were positive for *R. conorii* subsp. *israelensis* by the *gltA* gene; all samples were negative by *ompA* and *ompB* genes. The only sequence type we identified showed 99%–100% nucleotide identity with *R. conorii* subsp. *israelensis* from GenBank; we deposited our sequence in GenBank (accession no. OQ360110). All baboons tested negative for other vector-borne pathogens.

Although we found adult and male baboons at higher risk for infection (OR 2.6), we found no significant difference by age or sex (*p* = 0.439). No baboon showed ectoparasitic infestation or clinical signs of vector-borne diseases, and all displayed good physical status (mean complete blood count 3, average bodyweight 17.5 kg). Hematologic and serum chemistry values were within normal ranges (Appendix Tables 1, 2, <https://wwwnc.cdc.gov/EID/article/29/4/22-1176-App1.pdf>) for both *R. conorii*-negative and -positive baboons (*p* >0.05).

Our study revealed a nonnegligible prevalence (9.1%, 3/33) of *R. conorii* subsp. *israelensis* in *P. hamadryas* baboons, representing a pathogen–host association previously demonstrated only among asymptomatic dogs and cats from Portugal (5) and in severe cases among symptomatic humans from Italy (6). This survey confirms circulation of rickettsiae among baboons, also reported in 1 study of *R. africae* in *P. cynocephalus* yellow baboons from Zambia (1).

Despite routine treatment of baboons (orally administering 0.4 mg/kg ivermectin every 15 days by ground bait), presence of ticks in the zoopark was supported by a previous finding of tickborne pathogens (*A. phagocytophilum*, *Coxiella burnetii*, and *Rickettsia* spp.) in a lion (7). Given the baboon grooming behavior of removing ectoparasites from their bodies, lack of *Rhipicephalus sanguineus* sensu lato ticks, a vector of rickettsiae (8), was not surprising (9). However, association between zoopark-dwelling baboons and *Rhipicephalus* spp. ticks, including *R. sanguineus* s.l., is well known (9). Because this tick species is prevalent in the study area in all developmental stages, exposure very likely occurs (10).

Taken together, the high density of *P. hamadryas* baboons, their close proximity to the zoopark, and the anthrophilic behavior of *R. sanguineus* s.l. ticks (10) highlight the threat to park visitors and workers from *R. conorii* subsp. *israelensis* infection. Absence of clinical signs in positive baboons and lack of

Table. PCR protocols used in study of vector-borne pathogens among baboons, Italy, 2020

Pathogen	Target gene	Primer	Sequence, 5' → 3'	Fragment length, bp	Reference
<i>Babesia/Theileria</i> spp.	18S rRNA	RLB-F RLB-R	GAGGTAGTGACAAGAAATAACAATA TCTTCGATCCCCTAACTTTC	460–520	(2)
<i>Ehrlichia/Anaplasma</i> spp.	16S rRNA	EHR-16SD HER-16SR	GGTACCYACAGAAGAAGTCC TAGCACTCATCGTTTACAGC	345	(2)
<i>Rickettsia</i> spp.	<i>gltA</i>	CS-78F CS-323R	GCAAGTATCGGTGAGGATGTAAT GCTTCCTTAAAATTCAATAAATCAGGAT	401	(2)
Spotted fever group Rickettsiae	<i>ompA</i>	Rr190.70F Rr190.701R	ATGGCGAATATTTCTCCAAA GTTCCGTTAATGGCAGCATCT	632	(2)
	<i>ompB</i>	120–2788 120–3599	AAACAATAATCAAGGTAAGT TACTTCCGGTTACAGCAAAGT	600	(3)
<i>Leishmania infantum</i>	kDNA minicircle	Leish-1 Leish-2	AACTTTTCTGGTCTCTCCG GGTAG ACCCCCAGTTTCCCGCC	120	(4)

differences in hematological and biochemical parameters between negative and positive animals indicate the asymptomatic features of infection and make clarifying the baboons' role as a potential reservoir more urgent. Measures to control tick circulation should be established to reduce risk for transmission of *R. conorii* subsp. *israelensis* to zoopark visitors and workers.

Acknowledgments

The authors are grateful to the staff of Fasano Zoopark involved in field activities.

R.I. and D.O. were partially supported by EU funding within the NextGeneration EU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (Project no. PE00000007, INF-ACT).

About the Author

Mr. Sgroi has a PhD in animal health and zoonosis from the University of Bari Aldo Moro. His main research activities focus on biology, epidemiology, and control of vector-borne pathogens of zoonotic concern.

References

- Nakayama J, Hayashida K, Nakao R, Ishii A, Ogawa H, Nakamura I, et al. Detection and characterization of zoonotic pathogens of free-ranging non-human primates from Zambia. *Parasit Vectors*. 2014;7:490. <https://doi.org/10.1186/s13071-014-0490-x>
- Sgroi G, Iatta R, Lia RP, D'Alessio N, Manoj RRS, Veneziano V, et al. Spotted fever group rickettsiae in *Dermacentor marginatus* from wild boars in Italy. *Transbound Emerg Dis*. 2021;68:2111–20. <https://doi.org/10.1111/tbed.13859>
- Roux V, Raoult D. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (ompB). *Int J Syst Evol Microbiol*. 2000;50:1449–55. <https://doi.org/10.1099/00207713-50-4-1449>
- Sgroi G, Iatta R, Veneziano V, Bezerra-Santos MA, Lesiczka P, Hrazdilová K, et al. Molecular survey on tick-borne pathogens and *Leishmania infantum* in red foxes (*Vulpes vulpes*) from southern Italy. *Ticks Tick Borne Dis*. 2021;12:101669. <https://doi.org/10.1016/j.ttbdis.2021.101669>
- Maia C, Cristóvão JM, Pereira A, Parreira R, Campino L. Detection of *Rickettsia conorii israelensis* DNA in the blood of a cat and a dog from southern Portugal. *Top Companion Anim Med*. 2019;36:12–5. <https://doi.org/10.1053/j.tcam.2019.06.001>
- Guccione C, Colomba C, Rubino R, Bonura C, Anastasia A, Agrenzano S, et al. A severe case of Israeli spotted fever with pleural effusion in Italy. *Infection*. 2022;50:269–72. <https://doi.org/10.1007/s15010-021-01693-8>
- Torina A, Naranjo V, Pennisi MG, Patania T, Vitale F, Laricchiuta P, et al. Serologic and molecular characterization of tickborne pathogens in lions (*Panthera leo*) from the Fasano Safari Park, Italy. *J Zoo Wildl Med*. 2007;38:591–3. <https://doi.org/10.1638/2007-0043R1.1>
- Rovero C, Brouqui P, Raoult D. Questions on Mediterranean spotted fever a century after its discovery. *Emerg Infect Dis*. 2008;14:1360–7. <https://doi.org/10.3201/eid1409.071133>
- Akinyi MY, Tung J, Jeneby M, Patel NB, Altmann J, Alberts SC. Role of grooming in reducing tick load in wild baboons (*Papio cynocephalus*). *Anim Behav*. 2013;85:559–68. <https://doi.org/10.1016/j.anbehav.2012.12.012>
- Otranto D, Dantas-Torres F, Giannelli A, Latrofa MS, Cascio A, Cazzin S, et al. Ticks infesting humans in Italy and associated pathogens. *Parasit Vectors*. 2014;7:328. <https://doi.org/10.1186/1756-3305-7-328>

Address for correspondence: Domenico Otranto, University of Bari Aldo Moro, strada provinciale per Casamassima km3, 70010 Valenzano, Italy; email: domenico.otranto@uniba.it

Prevention of *Thelazia callipaeda* Reinfection among Humans

Marija Trenkić, Suzana Tasić-Otašević, Marcos Antonio Bezerra-Santos, Marko Stalević, Aleksandar Petrović, Domenico Otranto

Author affiliations: Ophthalmology Clinic University Clinical Center, Niš, Serbia (M. Trenkić); University of Niš, Niš (M. Trenkić, S. Tasić-Otašević, M. Stalević, A. Petrović); Public Health Institute, Niš (S. Tasić-Otašević); University of Bari Aldo Moro, Bari, Italy (M.A. Bezerra-Santos, D. Otranto); Bu-Ali Sina University, Hamedan, Iran (D. Otranto)

DOI: <https://doi.org/10.3201/eid2904.221610>

Thelazia callipaeda is a zoonotic vector-borne nematode that infects and causes eye disease among a wide range of domestic and wild mammals, including humans. We describe an unusual case of reinfection by this nematode in Serbia and call for a focus on preventive measures in endemic areas.

The genus *Thelazia* (order Spirurida, family Thelaziidae) comprises several species of nematode that cause ocular infections in different host mammals, including humans (1). Over the past 20 years, the *T. callipaeda* eyeworm has gained interest among

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Rickettsia conorii subspecies *israelensis* in Captive Baboons

Appendix

Appendix Table 1. Complete blood count and biochemical analyses results from baboons testing negative (n = 30) and positive (n = 3) to *Rickettsia conorii* subspecies *israelensis* DNA, during 2020, in Italy*

Parameter	Units	Mean BabR- †	Mean BabR+ ‡	Standard deviation	Standard error	t-test value	p-value	Range
Complete blood count	g/dL	12.7	12.2	1.7	1.0	0.5	0.632	12.6 ± 0.9
Hgb	K/μL	442.8	438.7	67.9	41.1	0.1	0.921	316 ± 83
Plt	Femtoliters	7.3	7.9	0.7	0.4	1.4	0.167	8.3 ± 1.0
MPV	K/μL	11.7	9.4	3.3	2.0	1.1	0.314	9.6 ± 2.9
WBC	M/μL	5.2	4.8	0.46	0.3	1.5	0.133	4.95 ± 0.32
RBC	%	40.7	36.6	3.6	2.2	1.8	0.069	38.2 ± 2.5
Hct	Femtoliters	77.6	78.6	2.6	1.6	0.6	0.530	77 ± 2.9
MCV	pg/dL	21.7	22.3	0.9	0.5	1.1	0.279	25.3 ± 0.9
MCH	g/dL	32.5	33.5	1.8	1.1	0.9	0.366	32.9 ± 0.7
MCHC	g/dL	31.7	32.2	1.9	1.1	0.5	0.650	NA
CHCM	pg/dL	3.7	3.4	0.4	0.2	1.2	0.224	NA
CHDW	%	14.0	14.5	0.7	0.4	1.2	0.247	NA
RDW	%	2.1	1.9	0.2	0.1	1.6	0.108	NA
HDW	K/μL	9.1	7.8	3.2	1.9	0.7	0.501	3.3 ± 1.9
Neu	K/μL	1.8	1.2	0.6	0.4	1.6	0.108	2.1 ± 1.3
Lym	K/μL	0.6	0.4	0.2	0.1	1.6	0.108	2.0 ± 2.0
Mon	K/μL	0.03	0.02	0.1	0.06	0.2	0.858	1.0 ± 1.0
Eos	K/μL	0.025	0.03	0.02	0.01	0.2	0.869	0.05 ± 0.05
Bas	%	0.3	0.3	0.07	0.04	0	1.000	NA
‡Biochemical analyses	%	41.2	40.8	3.2	1.9	0.2	0.838	NA
Pct	g/dL	23.1	20.9	3.3	1.9	1.1	0.279	NA
PDW	IU/L	564.0	477.0	622.6	377.0	0.2	0.819	NA
MPC	IU/L	49.6	31.7	16.3	9.8	1.8	0.079	NA
CPK	IU/L	33.4	18.7	32.6	19.7	0.7	0.462	NA
AST	IU/L	642.2	886.0	489.4	296.3	0.8	0.417	NA
ALT	IU/L	22.7	32.7	12.6	7.6	1.3	0.200	NA
ALP	IU/L	5,225.5	5,456.8	1,829.4	1,107.7	0.2	0.836	NA
GGT	mg/dL	0.4	0.3	0.1	0.06	1.6	0.111	NA
Cholinesterase	mEq/L	145.5	142.8	3.5	2.1	1.3	0.212	NA
Total bilirubin	mEq/L	4.2	4.5	0.7	0.4	0.7	0.485	NA
Sodium	mEq/L	22.0	18.7	5.8	3.5	0.9	0.355	NA
Potassium	mEq/L	111.0	107.4	3.5	2.1	1.7	0.099	NA
Sodium/potassium ratio	mmol/L	10.6	13.2	3.7	2.2	1.1	0.255	NA
Chlorine	mg/dL	120.3	105.6	30.5	18.5	0.8	0.432	NA
Anion gap	mg/dL	1.0	0.9	0.3	0.1	0.5	0.587	NA
Glucose	mg/dL	28.1	32.1	10.5	6.3	0.6	0.534	NA
Creatinine	mg/dL	9.0	9.5	0.7	0.4	1.2	0.247	NA
Urea	mg/dL	4.3	3.5	1.5	0.9	0.8	0.385	NA
Calcium	g/dL	6.6	6.8	0.5	0.3	0.7	0.514	NA
Phosphorus	mg/dL	0.18	0.2	0.05	0.03	0.7	0.510	NA
Total proteins	mg/dL	91.7	100.7	18.9	11.4	0.8	0.438	NA
Albumin	mg/dL	50.9	53.4	21.4	12.9	0.1	0.848	NA
Cholesterol	μg/dL	34.7	31.8	5.8	3.5	0.8	0.415	NA
Triglycerides	mmol/L	27.2	25.3	3.2	1.9	0.9	0.334	NA

*BabR- *R. conorii* subsp. *israelensis* DNA-negative baboons, BabR+ *R. conorii* subsp. *israelensis* DNA-positive baboons, HGB (hemoglobin), PLT (platelet), MPV (mean platelet volume), WBC (white blood cell), RBC (red blood cell), Hct (hematocrit), MCV (mean corpuscular volume), MCH (mean corpuscular hemoglobin), MCHC (mean corpuscular hemoglobin concentration), CHCM (cellular hemoglobin concentration mean), CHDW (cell hemoglobin distribution width), RDW (red cell distribution width), HDW (hemoglobin distribution width), Neu (neutrophils), Lym (lymphocytes), Mon (monocytes) Eos (eosinophils), Bas (basophils), Pct (procalcitonin), PDW (platelet distribution width), MPC (mean platelet component), CPK (creatinine phosphokinase), AST (aspartate aminotransferase), ALT (alanine aminotransferase), ALP (alkaline phosphatase), GGT (gamma glutamyl transpeptidase).

†BabR-, Baboons tested negative to *Rickettsia conorii* subspecies *israelensis* DNA.

‡BabR+, Baboons tested positive to *Rickettsia conorii* subspecies *israelensis* DNA.

Appendix Table 2. Serum protein electrophoresis in baboons tested negative (n = 33) and positive to *Rickettsia conorii* subspecies *israelensis* DNA, during 2020, in Italy. The protein concentration values are expressed in g/dl

Parameter	Mean value *BabR ⁻	Mean value †BabR ⁺	Standard deviation	Standard error	t-test value	p-value
Albumin	3.9	4.2	0.6	0.4	0.8	0.415
α-1 globulins	0.2	0.14	0.06	0.04	1.7	0.106
α-2 globulins	0.7	0.5	0.2	0.1	1.6	0.108
β-1 globulins	0.5	0.4	0.1	0.06	1.6	0.111
β-2 globulins	0.5	0.48	0.1	0.06	0.3	0.745
γ-globulins	0.9	0.8	0.3	0.06	1.6	0.111
Total proteins	6.6	6.5	0.5	0.3	0.3	0.744
Albumin/globulins ratio	1.5	1.9	0.5	0.3	1.3	0.196

*BabR⁻–*R. conorii* subsp. *israelensis* DNA–negative baboons, BabR⁺ *R. conorii* subsp. *israelensis* DNA–positive baboons