

Dr. Boostrom is the director of Family Health Services at the Corpus Christi-Nueces County Public Health District. Her research interests include communicable disease surveillance and the epidemiology of typhus in south Texas.

## References

1. Traub R, Wisseman CL Jr, Azad AF. The ecology of murine typhus: a critical review. *Trop Dis Bull* 1978;75:237–317.
2. Azad AF. Epidemiology of murine typhus. *Annu Rev Entomol* 1990;35:553–69.
3. Dumler JS, Taylor JP, Walker DH. Clinical and laboratory features of murine typhus in Texas, 1980 through 1987. *JAMA* 1991;266:1365–70.
4. Fan MY, Walker DH, Yu SR, Liu Q. Epidemiology and ecology of rickettsial diseases in the People's Republic of China. *Rev Infect Dis* 1987;9:823–40.
5. Corwin A, Habib M, Olson J, Scott D, Ksiazek T, Watts DM. The prevalence of arboviral, rickettsial, and hanta-like viral antibody among schoolchildren in the Nile delta of Egypt. *Trans R Soc Trop Med Hyg* 1992;86:677–9.
6. Silpapojakul K, Chupuppakarn S, Yuthasompob S, Varachit B, Chaipak D, Borkerd T, et al. Scrub and endemic murine typhus in children with obscure fever in the tropics. *Pediatr Infect Dis J* 1991;10:200–3.
7. Schriefer ME, Sacci JB Jr, Dumler JS, Bullen MG, Azad AF. Identification of a novel rickettsial infection in a patient diagnosed with murine typhus. *J Clin Microbiol* 1994;32:949–54.
8. Schriefer ME, Sacci JB Jr, Higgins JA, Taylor JP, Azad AF. Murine typhus: updated role of multiple urban components and a second typhus-like rickettsiae. *J Med Entomol* 1994;31:681–5.
9. Adams WH, Emmons RW, Brooks JE. The changing ecology of murine (endemic) typhus in southern California. *Am J Trop Med Hyg* 1970;19:311–8.
10. Williams SG, Sacci JB Jr, Schriefer ME, Anderson EM, Fujioka KK, Sorvillo FJ, et al. Typhus and typhus-like rickettsiae associated with opossums and their fleas in Los Angeles County, California. *J Clin Microbiol* 1992;30:1758–62.
11. Azad AF, Sacci JB Jr, Nelson WM, Dasch GA, Schmidtman ET, Carl M. Genetic characterization and transovarial transmission of a novel typhus-like *Rickettsia* found in cat fleas. *Proc Natl Acad Sci U S A* 1992;89:43–6.
12. Higgins JA, Radulovic S, Schriefer ME, Azad AF. *Rickettsia felis*: a new species of pathogenic rickettsia isolated from cat fleas. *J Clin Microbiol* 1996;34:671–4.
13. Zavala-Velazquez JD, Sosa-Ruiz JA, Zavala-Castro J, Jimenez-Delgado B, Vado-Solis IE, Sanchez-Elias RA, et al. *Rickettsia felis*—the etiologic agent of three cases of rickettsiosis in Yucatan. *Lancet* 2000;356:1079–80.
14. Bouyer DH, Stenos J, Crocquet-Valdes P, Moron CG, Popov VL, Zavala JE, et al. *Rickettsia felis*: the molecular characterization of a new member of the spotted fever group. *Int J Syst Evol Microbiol* 2001;51:339–47.
15. Raoult D, Scola BL, Enea M, Fournier PE, Roux V, Fenollar F, et al. A flea-associated *Rickettsia* pathogenic for humans. *Emerg Infect Dis* 2001;7:73–81.
16. Noden BH, Radulovic S, Higgins JA, Azad AF. Molecular identification of two closely related rickettsial species, *Rickettsia typhi* and *R. felis*, in individual cat fleas, *Ctenocephalides felis* (Siphonaptera: Pulicidae). *J Med Entomol* 1998;35:410–4.
17. Sorvillo FJ, Gondo B, Emmons R, Ryan P, Waterman SH, Tilzer A, et al. A suburban focus of endemic typhus in Los Angeles County: association with seropositive domestic cats and opossums. *Am J Trop Med Hyg* 1993;48:269–73.
18. Higgins JA, Sacci JB Jr, Schriefer ME, Endris RG, Azad AF. Molecular identification of rickettsia-like microorganisms associated with colonized cat fleas (*Ctenocephalides felis*). *Insect Mol Biol* 1994;3:27–33.
19. Azad AF, Radulovic S, Higgins JA, Noden BH, Troyer MJ. Flea-borne rickettsioses: some ecological considerations. *Emerg Infect Dis* 1997;3:319–28.
20. Azad AF, Beard CB. Interactions of rickettsial pathogens with arthropod vectors. *Emerg Infect Dis* 1998;4:179–86.

Address for correspondence: Abdu F. Azad, Department of Microbiology and Immunology, University of Maryland School of Medicine, 655 W. Baltimore Street, Baltimore, MD 21201, USA; fax: 410-706-0282; e-mail: aazad@umaryland.edu

### Emerging Infectious Diseases Policy on Corrections

The Emerging Infectious Diseases journal wishes error-free articles. To that end, we

1) Make corrections as quickly as we become aware of errors

2) Publish corrections online and in print. Online, we correct the error in the article it occurred with a note that the article was corrected and the date of correction. In print, we prominently publish a full correction, printing all needed information, and provide the URL of the corrected online article for reprints.

For additional information on corrections, send e-mail to [eideditor@cdc.gov](mailto:eideditor@cdc.gov).

### Correction, Vol. 8, No. 3

In *Listeria monocytogenes* Infection in Israel and Review of Cases Worldwide, by Y. Siegman-Igra et al., an error appears in the discussion section. The corrected sentence appears below and online at <http://www.cdc.gov/ncidod/EID/vol8no3/01-0195.htm>.

The case-fatality rate in the collected data on nonperinatal infection was 36% (413 of 1,149 patients for whom this information was available).

We regret any confusion this error may have caused