

IMPORTANT UPDATE

Effective November 11, 2009

FDA Expands Use of CSL Limited™ 2009 H1N1 Monovalent Influenza Vaccine to Persons Ages 6 Months and Older

CDC Recommendations for Use

Patient's Age and Vaccine Dose	CSL Limited™ Available Vaccine Formulations	
	0.5 mL Pre-Filled Syringes	5 mL Multi-Dose Vial
6 through 35 months (0.25 mL)		
	<p>Use when other age appropriate formulations are not available</p> <ul style="list-style-type: none"> Administer a half dose (0.25 mL) from syringe contents; unused vaccine must be discarded Using CSL™ pre-filled syringes in children 6 through 35 months of age will result in wasted vaccine 	<p>Recommended for use</p> <ul style="list-style-type: none"> Ancillary supplies (needles, syringes, alcohol prep pads, and vaccination records) provided with each order are sufficient to administer 10 doses per vial Providers who choose to administer more than 10 doses per vial will need to supplement their own ancillary supplies
3 years and older (0.5 mL)	<p>Recommended for use</p> <ul style="list-style-type: none"> No concerns with wasted vaccine Needles, alcohol prep pads, and vaccination records supplied with each order of vaccine will be sufficient 	<p>Recommended for use</p> <ul style="list-style-type: none"> Needles, syringes, alcohol prep pads, and vaccination records supplied with each order of vaccine will be sufficient

