

HPAI Outbreaks in U.S. Birds: Self-Observation Instructions for Demobilizing Bird Flu Responders

You are being given this information and these instructions because you participated in the U.S. response to domestic outbreaks of avian influenza.

As part of your work, you may have been around bird flu viruses. Infected birds shed avian influenza virus in their saliva, mucous and feces. Human infections with bird flu viruses can happen when enough virus gets into a person's eyes, nose or mouth, or is inhaled. This can happen when virus is in the air (in droplets or possibly dust) and a person breathes it in, or when a person touches something that has virus on it then touches their mouth, eyes or nose. Rare human infections with some avian viruses have occurred most often after unprotected contact with infected birds or surfaces contaminated with avian influenza viruses. However, some infections have been identified where direct contact was not known to have occurred.

The Centers for Disease Control and Prevention (CDC) believes the risk of infection with AI viruses is low. Because human infections with these viruses are possible, however, all people participating in avian influenza outbreak response efforts should be monitored for illness for 10 days after their last possible exposure to infected birds or potentially-contaminated environments, even if exposure to the sick birds was minimal or if personal protective equipment (PPE) was worn appropriately. State and local health departments are helping with this monitoring effort and they may contact you while you're observing your health. By following the instructions below, you can help ensure that you receive prompt medical evaluation, possible testing and appropriate treatment if you become ill with signs and symptoms that could be due to AI virus infection. Thank you for your contribution to the AI domestic response effort.

Please follow these instructions carefully:

1. Monitor your health for symptoms of avian influenza virus infection.

Immediately following demobilization, monitor yourself daily for any of these signs and symptoms for 10 days:

Fever (Temperature of 100°F [37.8°C] or greater) or feeling feverish.	Diarrhea
Cough	Nausea
Sore throat	Vomiting
Runny or stuffy nose	Fatigue
Muscle or body aches	Seizures
Headaches	Rash
Eye redness	
Difficulty breathing	
Shortness of breath	

You should observe your health daily even if you carefully followed all guidelines and instructions from your Safety Officer for properly putting on and taking off personal protective equipment (PPE) and maintaining biosecurity precautions while working the response.

HPAI H5 Outbreaks in U.S. Birds: Self-Observation Instructions for Demobilizing H5 Bird Flu Responders

Re-start your 10-day monitoring period from Day 1 if, while you are observing your health for signs and symptoms, you are around sick birds again.

2. Call your state/local public health department immediately if you develop any illness signs or symptoms during the 10-day observation period.

Responders who have demobilized (are no longer working on the response) should report any illness signs or symptoms to the health department of the state they are in at the time. Your health department will help you determine what to do next. (Responders who are still on duty should report any illness to their Safety Officer.)

Remember:

- Most of the signs and symptoms of bird flu overlap with those of other respiratory illnesses (like seasonal flu).
- If you develop any of the signs or symptoms listed in this fact sheet, call the health department of the state you are in immediately. Your health department wants to hear from you, even if it turns out to be a 'false alarm'.
- Your health department may contact you by phone, email or text while you are observing your health.
- If you have symptoms, your health department may give you instructions and ask you to get tested for avian influenza virus infection.
 - To test for avian influenza virus infection, a doctor or nurse will collect a sample from you by swabbing your nose and/or throat.
- If you become sick while you are observing your health after demobilization, a doctor may prescribe you an antiviral medication that is used for treatment of influenza. It's important to take the medication as prescribed. (CDC recommends that clinicians prescribe antiviral medications for treatment of ill persons who had exposure to avian influenza viruses)

Here's an example of a log you can keep as a reminder to check your health each day for illness signs and symptoms.

Day	Date	Temperature (Fahrenheit)	Signs and Symptoms <i>Write 'None' if you aren't experiencing any of the signs and symptoms.</i>
1	Nov. 8	98.5	None
2	Nov. 9	99.1	Cough
3			
4			
5			
6			

Where to Find More Information:

Visit <http://www.cdc.gov/flu/avianflu/h5/index.htm> or <http://espanol.cdc.gov/enes/flu/avianflu/h5/index.htm>

Call 1-800-CDC-INFO (1-800-232-4636); TTY: 888-232-6348