

2008-2009 Influenza Season Week 37 ending September 19, 2009

All data are preliminary and may change as more reports are received.

Synopsis: During week 37 (September 13-19, 2009), influenza activity increased in the U.S.

- 2,326 (23.9%) specimens tested by U.S. World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NREVSS) collaborating laboratories and reported to CDC/Influenza Division were positive for influenza.
- 99% of all subtyped influenza A viruses being reported to CDC were 2009 influenza A (H1N1) viruses.
- The proportion of deaths attributed to pneumonia and influenza (P&I) was below the epidemic threshold.
- Three influenza-associated pediatric deaths were reported and all three were associated with 2009 influenza A (H1N1) virus infection.
- The proportion of outpatient visits for influenza-like illness (ILI) was above the national baseline. Regions 2 through 10 reported ILI above region-specific baseline levels; only Region 1 was below its' region-specific baseline.
- Twenty-six states reported geographically widespread influenza activity, 11 states reported regional influenza activity, 12 states, the District of Columbia, and Puerto Rico reported local influenza activity, one state and Guam reported sporadic influenza activity, and the U.S. Virgin Islands did not report.
- The 2009-10 influenza season officially begins October 4, 2009.

National and Regional Summary of Select Surveillance Components

HHS Surveillance Regions*	Data for current week			Data cumulative for the 2008-09 season						
	Out-patient ILI†	% positive for flu‡	Number of jurisdictions reporting regional or widespread activity§	A (H1)	A (H3)	2009 A (H1N1)	A (unable to sub-type)¶	A (Subtyping not performed)	B	Pediatric Deaths
Nation	Elevated	23.9%	37 of 54	8,240	4,363	46,329	1,325	23,056	10,782	117
Region 1	Normal	4.2%	2 of 6	584	304	3,041	13	1,711	815	4
Region 2	Elevated	3.7%	1 of 4	290	228	1,829	21	2,398	713	20
Region 3	Elevated	15.1%	5 of 6	1,250	222	4,846	20	1,069	1,363	10
Region 4	Elevated	24.2%	8 of 8	1,078	784	8,719	105	4,644	1,318	13
Region 5	Elevated	20.2%	5 of 6	1,593	213	8,903	214	1,009	1,383	18
Region 6	Elevated	25.9%	5 of 5	830	312	4,595	14	6,108	2,669	18
Region 7	Elevated	21.0%	3 of 4	488	84	1,448	570	660	519	1
Region 8	Elevated	17.1%	3 of 6	541	219	1,989	80	2,231	506	9
Region 9	Elevated	37.2%	3 of 5	1,200	1,680	7,937	74	2,687	810	21
Region 10	Elevated	27.6%	2 of 4	386	317	3,022	214	539	686	3

* HHS regions (Region I: CT, ME, MA, NH, RI, VT; Region II: NJ, NY, Puerto Rico, US Virgin Islands; Region III: DE, DC, MD, PA, VA, WV; Region IV: AL, FL, GA, KY, MS, NC, SC, TN; Region V: IL, IN, MI, MN, OH, WI; Region VI: AR, LA, NM, OK, TX; Region VII: IA, KS, MO, NE; Region VIII: CO, MT, ND, SD, UT, WY; Region IX: AZ, CA, Guam, HI, NV; and Region X: AK, ID, OR, WA).

† Elevated means the % of visits for ILI is at or above the national or region-specific baseline.

‡ National data are for current week; regional data are for the most recent three weeks.

§ Includes all 50 states, the District of Columbia, Guam, Puerto Rico, and U.S. Virgin Islands.


¶ The majority of influenza A viruses that cannot be sub-typed as seasonal influenza viruses are 2009 A (H1N1) influenza viruses upon further testing.

U.S. Virologic Surveillance: WHO and NREVSS collaborating laboratories located in all 50 states and Washington D.C. report to CDC the number of respiratory specimens tested for influenza and the number positive by influenza type and subtype. The results of tests performed during the current week are summarized in the table below.

	Week 37
No. of specimens tested	9,744
No. of positive specimens (%)	2,326 (24.9%)
Positive specimens by type/subtype	
Influenza A	2,324 (99.9%)
A (2009 H1N1)	1,395 (60.0%)
A (subtyping not performed)	907 (39.0%)
A (unable to subtype)	17 (0.7%)
A (H3)	1 (0.1%)
A (H1)	4 (0.2%)
Influenza B	2 (0.1%)

During week 37, seasonal influenza A (H1), A (H3) and influenza B viruses co-circulated at low levels with 2009 influenza A (H1N1) viruses. Ninety-nine percent of all subtyped influenza A viruses reported to CDC this week were 2009 influenza A (H1N1) viruses.

Influenza Positive Tests Reported to CDC by U.S. WHO/NREVSS Collaborating Laboratories, National Summary, 2008-09



Pneumonia and Influenza Hospitalization and Death Tracking: This new system was implemented on August 30, 2009, and replaces the weekly report of laboratory confirmed 2009 H1N1-related hospitalizations and deaths. Jurisdictions can now report to CDC either laboratory confirmed or pneumonia and influenza syndromic-based counts of hospitalizations and deaths resulting from all types or subtypes of influenza, not just those from 2009 H1N1 influenza virus. To allow jurisdictions to implement the new case definition, counts were reset to zero on August 30, 2009. From August 30-September 19, 2009, 10,082 hospitalizations and 936 deaths associated with influenza virus infection or based on syndromic surveillance for influenza and pneumonia, were reported to CDC. This is the third week of data from this new system and reflects reports by 52 jurisdictions. CDC will continue to use its traditional surveillance systems to track the progress of the remainder of the 2008-09 season, and for the 2009-10 influenza season, which officially begins October 4, 2009.

Antigenic Characterization: CDC has antigenically characterized 2,191 seasonal human influenza viruses [1,213 influenza A (H1), 264 influenza A (H3) and 714 influenza B viruses] collected by U.S. laboratories since October 1, 2008, and 716 2009 influenza A (H1N1) viruses.

1,186 (98%) of 1,213 seasonal influenza A (H1) viruses tested are related to the influenza A (H1N1) component of the 2009-10 Northern Hemisphere influenza vaccine (A/Brisbane/59/2007) and 27 (2%) showed reduced titers with antisera produced against A/Brisbane/59/2007. Two hundred twenty-nine (87%) of 264 influenza A (H3N2) viruses tested are related to the influenza A (H3N2) 2009-10 vaccine component (A/Brisbane/10/2007) and 35 viruses (13%) tested showed reduced titers with antisera produced against A/Brisbane/10/2007.

All 716 2009 influenza A (H1N1) viruses are related to the A/California/07/2009 (H1N1)pdm reference virus selected by WHO as the component of the 2009 influenza A (H1N1) vaccine.

Influenza B viruses currently circulating can be divided into two distinct lineages represented by the B/Yamagata/16/88 and B/Victoria/02/87 viruses. Six hundred thirty-eight (89%) of 714 influenza B viruses tested belong to the B/Victoria lineage, and of these, 630 (99%) are related to the influenza vaccine component for the 2009-10 Northern Hemisphere influenza vaccine (B/Brisbane/60/2008), and the remaining eight (1%) showed reduced titers with antisera produced against B/Brisbane/60/2008. Seventy-six (11%) influenza B viruses tested belong to the B/Yamagata lineage.

Data on antigenic characterization should be interpreted with caution given that antigenic characterization data are based on hemagglutination inhibition (HI) testing using a panel of reference ferret antisera, and results may not correlate with clinical protection against circulating viruses provided by influenza vaccination.

Annual influenza vaccination is expected to provide the best protection against those virus strains that are related to the vaccine strains, but limited to no protection may be expected when the vaccine and circulating virus strains are so different as to be from different lineages, as is seen with the two lineages of influenza B viruses. Antigenic characterization of 2009 influenza A (H1N1) viruses indicates that these viruses are antigenically and genetically unrelated to seasonal influenza A (H1N1) viruses, suggesting that little to no protection would be expected from vaccination with seasonal influenza vaccine.

Antiviral Resistance: Since October 1, 2008, 1,148 seasonal influenza A (H1N1), 261 influenza A (H3N2), 654 influenza B, and 705 2009 influenza A (H1N1) virus isolates have been tested for resistance to the neuraminidase inhibitors (oseltamivir and zanamivir). In addition, 973 2009 influenza A (H1N1) original clinical samples were tested for a single known mutation in the virus that confers oseltamivir resistance. Also, 1,153 seasonal influenza A (H1N1), 261 influenza A (H3N2), and 526 2009 influenza A (H1N1) viruses have been tested for resistance to the adamantanes (amantadine and rimantadine). The results of antiviral resistance testing performed on these viruses are summarized in the table below. Additional laboratories perform antiviral testing and report their results to CDC. Two additional oseltamivir resistant 2009 influenza A (H1N1) viruses have been identified by these laboratories.

	Viruses tested (n)	Resistant Viruses, Number (%)	Viruses tested (n)	Resistant Viruses, Number (%)	Viruses tested (n)	Resistant Viruses, Number (%)
		Oseltamivir		Zanamivir		Adamantanes
Seasonal Influenza A (H1N1)	1,148	1,143 (99.6%)	1,148	0 (0)	1,153	6 (0.5%)
Influenza A (H3N2)	261	0 (0)	261	0 (0)	261	261 (100%)
Influenza B	654	0 (0)	654	0 (0)	N/A*	N/A*
2009 Influenza A (H1N1)	1,678	9 ^{†‡} (0.6)	705	0 (0)	526	526 (100%)

*The adamantanes (amantadine and rimantadine) are not effective against influenza B viruses.

[†]Two screening tools were used to determine oseltamivir resistance: sequence analysis of viral genes and a neuraminidase inhibition assay.

[‡] Additional laboratories perform antiviral resistance testing and report their results to CDC. Two additional oseltamivir resistant 2009 influenza A (H1N1) viruses have been identified by these laboratories, bringing the total number to 11.

2009 influenza A (H1N1) viruses were tested for oseltamivir resistance by a neuraminidase inhibition assay and/or detection of genetic sequence mutation, depending on the type of specimen tested. Original clinical samples were examined for a single known mutation in the virus that confers oseltamivir resistance in currently circulating seasonal influenza A (H1N1) viruses, while influenza virus isolates were tested using a neuraminidase inhibition assay that determines the presence or absence of neuraminidase inhibitor resistance, followed by the neuraminidase gene sequence analysis of resistant viruses.


The majority of 2009 influenza A (H1N1) viruses are susceptible to the neuraminidase inhibitor antiviral medication oseltamivir; however, rare sporadic cases of oseltamivir resistant 2009 influenza A (H1N1) viruses have been detected worldwide, including 11 cases in the United States. All tested viruses retain their sensitivity to the other neuraminidase inhibitor: zanamivir. Ten patients (including the nine viruses detected at CDC and one identified by an additional laboratory) had documented exposure to oseltamivir through either treatment or chemoprophylaxis, and the remaining patient is currently under investigation to determine exposure to oseltamivir. Occasional development of oseltamivir resistance during treatment or prophylaxis is not unexpected. Enhanced surveillance is expected to detect additional cases of oseltamivir resistant 2009 influenza A (H1N1) viruses, and such cases will be investigated to assess the spread of resistant strains in the community. To prevent the spread of antiviral resistant virus strains, CDC reminds clinicians and the public of the need to continue hand and cough hygiene measures for the duration of any symptoms of influenza (<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5832a3.htm>).

Additional information on antiviral recommendations for treatment and chemoprophylaxis of influenza virus infection is available at <http://www.cdc.gov/h1n1flu/recommendations.htm>. Antiviral treatment with oseltamivir or zanamivir is recommended for all patients with confirmed or suspected influenza virus infection who are hospitalized or who are at higher risk for influenza complications.

Three seasonal influenza A (H1N1) viruses collected between February 8 and May 11, 2009, were found to be resistant to both oseltamivir and the adamantanes (amantadine and rimantadine). All seasonal influenza A (H1N1) viruses tested retain their sensitivity to zanamivir. The three dually-resistant viruses represent less than 1% of all seasonal influenza A (H1N1) viruses tested during the 2008-09 influenza season, and as a result, no changes to the influenza antiviral treatment or prophylaxis recommendations will be made at this time. CDC will continue to monitor trends in antiviral resistance over the summer and throughout the upcoming 2009-10 influenza season.

Pneumonia and Influenza (P&I) Mortality Surveillance: During week 37, 6.1% of all deaths reported through the 122-Cities Mortality Reporting System were due to P&I. This percentage was below the epidemic threshold of 6.3% for week 37.


Pneumonia and Influenza Mortality for 122 U.S. Cities
Week ending 9/19/2009



Influenza-Associated Pediatric Mortality: Three influenza-associated pediatric deaths were reported to CDC during week 37 (Texas [2] and Virginia). These deaths were associated with 2009 influenza A (H1N1) virus infection, and occurred between August 30 and September 19, 2009. Since September 28, 2008, CDC has received 117 reports of influenza-associated pediatric deaths that occurred during the current influenza season (25 deaths in children less than 2 years, 12 deaths in children 2-4 years, 35 deaths in children 5-11 years, and 45 deaths in individuals 12-17 years). Forty-nine of the 117 deaths were due to 2009 influenza A (H1N1) virus infections, and four of these have occurred since August 30, 2009.

Of the 50 children who had specimens collected for bacterial culture from normally sterile sites, 19 (38.0%) were positive; *Staphylococcus aureus* was identified in 13 (68.4%) of the 19 children. Six of the *S. aureus* isolates were sensitive to methicillin and seven were methicillin resistant. Seventeen (89.5%) of the 19 children with bacterial coinfections were five years of age or older and 12 (63.2%) of the 19 children were 12 years of age or older. Nineteen (38.8%) of the 49 children with confirmed 2009 influenza A (H1N1) infection had a specimen collected from a normally sterile site; five (26.3%) of the 19 children had a positive bacterial culture (methicillin sensitive *S. aureus* [2], methicillin resistant *S. aureus* [2], and *Streptococcus constellatus*).

Number of Influenza-Associated Pediatric Deaths by Week of Death: 2005-06 season to present




Influenza-Associated Hospitalizations: Laboratory-confirmed influenza-associated hospitalizations are monitored through the Emerging Infections Program (EIP), a population-based surveillance network.

During April 15, 2009 – September 19, 2009, the following preliminary laboratory-confirmed overall influenza associated hospitalization rates were reported by the EIP (*rates include influenza A, influenza B, and 2009 influenza A (H1N1)*):

Rates for children aged 0-23 months, 2-4 years, and 5-17 years were 2.6, 1.1, and 0.9 per 10,000, respectively. Rates for adults aged 18-49 years, 50-64 years, and ≥ 65 years, the overall flu rates were 0.6, 0.6, and 0.5 per 10,000, respectively.

EIP Influenza Laboratory-Confirmed Cumulative Hospitalization Rates, Spring/Summer 2009




*This value represents an age group-specific average influenza rate from October 1 to April 30 from the 2005-06, 2006-07, and 2007-08 influenza seasons.

**Note: The scales for the 0-23 month and the ≥65 year age groups differ from other age groups.











Outpatient Illness Surveillance: Nationwide during week 37, 4.6% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is above the national baseline of 2.4%.

Percentage of Visits for Influenza-like Illness (ILI) Reported by the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), National Summary, 2008-09 and Previous Two Seasons



*There was no week 53 during the 2006-07 and 2007-08 seasons, therefore the week 53 data point for those seasons is an average of weeks 52 and 1.

On a regional level, the percentage of outpatient visits for ILI ranged from 1.4% to 8.9%, and decreased in four of the 10 surveillance regions. Nine regions (Regions 2 through 10) reported a proportion of outpatient visits for ILI above region-specific baseline levels, while Region 1 reported a percentage of visits for ILI below its' region-specific baseline.




NOTE: Scales differ between regions

NOTE: There was no week 53 during the 2006-07 and 2007-08 seasons, therefore the week 53 data point for those seasons is an average of weeks 52 and 1.

Geographic Spread of Influenza as Assessed by State and Territorial Epidemiologists: The influenza activity reported by state and territorial epidemiologists indicates geographic spread of both seasonal influenza and 2009 influenza A (H1N1) viruses and does not measure the severity of influenza activity.

During week 37, the following influenza activity was reported:

- Widespread influenza activity was reported by 26 states (Alabama, Alaska, Arizona, Arkansas, California, Colorado, Delaware, Florida, Georgia, Illinois, Kansas, Kentucky, Louisiana, Maryland, Minnesota, Mississippi, Nevada, New Mexico, North Carolina, Oklahoma, Pennsylvania, South Carolina, Texas, Tennessee, Virginia, and Washington).
- Regional influenza activity was reported by 11 states (Indiana, Iowa, Maine, Montana, Nebraska, New Hampshire, New Jersey, Ohio, South Dakota, West Virginia, and Wisconsin).
- Local influenza activity was reported by the District of Columbia, Puerto Rico, and 12 states (Connecticut, Hawaii, Idaho, Massachusetts, Michigan, Missouri, New York, North Dakota, Oregon, Rhode Island, Utah, and Wyoming).
- Sporadic activity was reported by Guam and one state (Vermont).
- The U.S. Virgin Islands did not report.



* This map indicates geographic spread & does not measure the severity of influenza activity

A description of surveillance methods is available at: <http://www.cdc.gov/flu/weekly/fluactivity.htm>

Report prepared: September 25, 2009.