

## 2009-2010 Influenza Season Week 14 ending April 10, 2010

All data are preliminary and may change as more reports are received.

**Synopsis:** During week 14 (April 4-10, 2010), influenza activity decreased in the U.S.

- 52 (2.7%) specimens tested by U.S. World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NREVSS) collaborating laboratories and reported to CDC/Influenza Division were positive for influenza.
- Among 34 subtyped influenza A viruses, 33 were 2009 influenza A (H1N1) and one was influenza A (H3).
- The proportion of deaths attributed to pneumonia and influenza (P&I) was below the epidemic threshold.
- Three influenza-associated pediatric deaths were reported. One was associated with 2009 influenza A (H1N1) virus infection, one was associated with an influenza A virus for which the subtype was undetermined, and one death was associated with a seasonal influenza A (H1) virus infection, but occurred during the 2008-09 season.
- The proportion of outpatient visits for influenza-like illness (ILI) was 0.9%, which is below the national baseline of 2.3%. All 10 regions reported ILI below region-specific baseline levels.
- No states reported widespread influenza activity. Two states reported regional influenza activity. Puerto Rico and one state reported local influenza activity. The District of Columbia, Guam and 37 states reported sporadic influenza activity. The U.S. Virgin Islands and nine states reported no influenza activity, and one state did not report.

### National and Regional Summary of Select Surveillance Components

HHS Surveillance Regions**	Data for current week			Data cumulative since August 30, 2009 (Week 35)*						
	Out-patient ILI†	% positive for flu‡	Number of jurisdictions reporting regional or widespread activity§	A (H1)	A (H3)	2009 A (H1N1)	A (unable to subtype)¶	A (Subtyping not performed)	B	Pediatric Deaths
<b>Nation</b>	Normal	2.7%	2 of 54	34	65	66,605	288	22,408	295	272
<b>Region 1</b>	Normal	0.6%	0 of 6	4	5	3,403	14	503	12	6
<b>Region 2</b>	Normal	0.7%	0 of 4	8	6	1,809	0	2,404	14	19
<b>Region 3</b>	Normal	1.9%	0 of 6	3	11	10,714	48	1,453	18	14
<b>Region 4</b>	Normal	9.6%	2 of 8	0	7	9,813	95	4,679	89	52
<b>Region 5</b>	Normal	3.2%	0 of 6	8	25	9,495	46	1,579	19	36
<b>Region 6</b>	Normal	4.2%	0 of 5	2	3	5,185	19	5,408	56	73
<b>Region 7</b>	Normal	5.1%	0 of 4	3	1	3,292	3	847	9	8
<b>Region 8</b>	Normal	1.4%	0 of 6	4	2	9,863	0	3,941	56	16
<b>Region 9</b>	Normal	1.1%	0 of 5	0	4	8,344	49	1,246	18	38
<b>Region 10</b>	Normal	1.3%	0 of 4	2	1	4,687	14	348	4	10

\*Influenza season officially begins each year at week 40. This season data from week 35 will be included to show the trend of influenza activity before the official start of the 2009-10 influenza season.

\*\*HHS regions (Region 1 CT, ME, MA, NH, RI, VT; Region 2: NJ, NY, Puerto Rico, US Virgin Islands; Region 3: DE, DC, MD, PA, VA, WV; Region 4: AL, FL, GA, KY, MS, NC, SC, TN; Region 5: IL, IN, MI, MN, OH, WI; Region 6: AR, LA, NM, OK, TX; Region 7: IA, KS, MO, NE; Region 8: CO, MT, ND, SD, UT, WY; Region 9: AZ, CA, Guam, HI, NV; and Region 10: AK, ID, OR, WA). Use of the national baseline for regional data or regional baselines for state data is not appropriate.

† Elevated means the % of visits for ILI is at or above the national or region-specific baseline.

‡ National data are for current week; regional data are for the most recent three weeks.

§ Includes all 50 states, the District of Columbia, Guam, Puerto Rico, and U.S. Virgin Islands.


¶ Subtyping results for the majority of specimens in this category were inconclusive because of low virus titers.

**U.S. Virologic Surveillance:** WHO and NREVSS collaborating laboratories, located in all 50 states and Washington, D.C., report to CDC the number of respiratory specimens tested for influenza and the number positive by influenza type and subtype. The results of tests performed during the current week are summarized in the table below.

	<b>Week 14</b>
<b>No. of specimens tested</b>	1,939
<b>No. of positive specimens (%)</b>	52 (2.7%)
<b>Positive specimens by type/subtype</b>	
<b>Influenza A</b>	52 (100.0%)
<b>A (2009 H1N1)</b>	33 (63.5%)
<b>A (subtyping not performed)</b>	18 (34.6%)
<b>A (unable to subtype)</b>	0 (0.0%)
<b>A (H3)</b>	1 (1.9%)
<b>A (H1)</b>	0 (0.0%)
<b>Influenza B</b>	0 (0.0%)

During week 14, one influenza A (H3N2) virus was reported, and all other subtyped influenza A viruses reported to CDC this week were 2009 influenza A (H1N1) viruses.


Influenza Positive Tests Reported to CDC by U.S. WHO/NREVSS Collaborating Laboratories, National Summary, August 30, 2009-April 10, 2010



**Pneumonia and Influenza Hospitalization and Death Tracking:** The Aggregate Hospitalization and Death Reporting Activity (AHDRA) system was implemented on August 30, 2009, and replaces the weekly report of laboratory confirmed 2009 H1N1-related hospitalizations and deaths that began in April 2009. Jurisdictions report to CDC the number of hospitalizations and deaths resulting from all types or subtypes of influenza, not just those associated with 2009 H1N1. Counts were reset to zero on August 30, 2009. From August 30, 2009 – April 3, 2010, 41,883 laboratory-confirmed influenza-associated hospitalizations and 2,125 laboratory-confirmed influenza-associated deaths were reported to CDC.

Reporting of influenza-associated hospitalizations and deaths through AHDRA was discontinued during the week ending April 3, 2010 (week 13). CDC will continue to use its traditional influenza surveillance systems to track the remainder of the 2009-10 season.

### Weekly Laboratory-Confirmed Influenza-Associated Hospitalizations and Deaths Reported to AHDRA, National Summary, August 30, 2009 – April 3, 2010



**Antigenic Characterization:** CDC has antigenically characterized two seasonal influenza A (H1N1), 13 influenza A (H3N2), 23 influenza B, and 1,701 2009 influenza A (H1N1) viruses collected since September 1, 2009.

Both seasonal influenza A (H1N1) viruses tested were related to the influenza A (H1N1) component of the 2009-10 Northern Hemisphere influenza vaccine (A/Brisbane/59/2007).

The 13 influenza A (H3N2) viruses tested showed reduced titers with antisera produced against A/Brisbane/10/2007, the 2009-2010 Northern Hemisphere influenza A (H3N2) vaccine component, and were antigenically related to A/Perth/16/2009, the WHO recommended influenza A (H3N2) component of the 2010 Southern Hemisphere and 2010-11 Northern Hemisphere vaccine formulations.

Influenza B viruses currently circulating globally can be divided into two distinct lineages represented by the B/Yamagata/16/88 and B/Victoria/02/87 viruses. The influenza B component of the 2009-10 and 2010-11 Northern Hemisphere vaccines belongs to the B/Victoria lineage. The 23 influenza B viruses tested belong to the B/Victoria lineage and are related to the influenza vaccine component for the 2009-10 and 2010-11 Northern Hemisphere influenza B vaccine strain (B/Brisbane/60/2008).

One thousand six hundred ninety-six (99.7%) of 1,701 2009 influenza A (H1N1) viruses tested are related to the A/California/07/2009 (H1N1) reference virus selected by WHO as the 2009 H1N1 vaccine virus, and as a component in the 2010-11 Northern Hemisphere vaccine. Five viruses (0.3%) tested showed reduced titers with antiserum produced against A/California/07/2009.

Annual influenza vaccination is expected to provide the best protection against those virus strains that are related to the vaccine strains, but limited to no protection may be expected when the vaccine and circulating virus strains are so different as to be from different lineages. Antigenic characterization of 2009 influenza A (H1N1) viruses indicates that these viruses are only distantly related antigenically and genetically to seasonal influenza A (H1N1) viruses, suggesting that little to no protection would be expected from vaccination with seasonal influenza vaccine.

**Composition of the 2010-11 Influenza Vaccine:** WHO has recommended vaccine strains for the 2010-11 Northern Hemisphere trivalent influenza vaccine, and FDA has made the same recommendations for the U.S. influenza vaccine. Both agencies recommend that the vaccine contain A/California/7/2009-like (2009 H1N1), A/Perth/16/2009-like (H3N2), and B/Brisbane/60/2008-like (B/Victoria lineage) viruses. A seasonal influenza A (H1N1) component is not included in the 2010-11 formulation, and the A (H3N2) component has been changed from the 2009-10 Northern Hemisphere vaccine formulation. This recommendation was based on surveillance data related to epidemiology and antigenic characteristics, serological responses to 2009-10 trivalent seasonal and 2009 H1N1 monovalent vaccines, and the availability of candidate strains and reagents.

**Antiviral Resistance:** Since September 1, 2009, one seasonal influenza A (H1N1), 13 influenza A (H3N2), 23 influenza B, and 1,705 2009 influenza A (H1N1) virus isolates have been tested for resistance to the neuraminidase inhibitors (oseltamivir and zanamivir), and 2,748 2009 influenza A (H1N1) original clinical samples were tested for a single known mutation in the virus that confers oseltamivir resistance. In addition, one seasonal influenza A (H1N1), 13 influenza A (H3N2), and 1,705 2009 influenza A (H1N1) virus isolates have been tested for resistance to the adamantanes (amantadine and rimantadine). The results of antiviral resistance testing performed on these viruses are summarized in the table below. Additional laboratories perform antiviral testing and report their results to CDC, and positive results from that testing are included in the footnote.

**Antiviral Resistance Testing Results on Samples Collected Since September 1, 2009.**

	Viruses tested (n)	Resistant Viruses, Number (%)	Viruses tested (n)	Resistant Viruses, Number (%)	Isolates tested (n)	Resistant Viruses, Number (%)
		Oseltamivir		Zanamivir		Adamantanes
Seasonal Influenza A (H1N1)	1	1 (100.0)	0	0 (0)	1	0 (0)
Influenza A (H3N2)	13	0 (0)	0	0 (0)	13	13 (100.0)
Influenza B	23	0 (0)	0	0 (0)	N/A*	N/A*
2009 Influenza A (H1N1)	4,453	52 <sup>†‡</sup> (1.2)	1,617	0 (0)	1,705	1,701 (99.8)

\*The adamantanes (amantadine and rimantadine) are not effective against influenza B viruses.

<sup>†</sup>Two screening tools were used to determine oseltamivir resistance: sequence analysis of viral genes and a neuraminidase inhibition assay.

<sup>‡</sup>Additional laboratories perform antiviral resistance testing and report their results to CDC. Three additional oseltamivir resistant 2009 influenza A (H1N1) virus has been identified by these laboratories since September 1, 2009, bringing the total number to 55.

All but one subtyped influenza A virus reported during week 14 was a 2009 influenza A (H1N1) virus, and nearly all of 2009 influenza A (H1N1) viruses tested since April 2009 have been resistant to the adamantanes (amantadine and rimantadine).

Antiviral treatment with oseltamivir or zanamivir is recommended for all patients with confirmed or suspected influenza virus infection who are hospitalized, are at higher risk for influenza complications, or who have lower respiratory tract or progressive disease. Additional information on antiviral recommendations for treatment and chemoprophylaxis of influenza virus infection is available at <http://www.cdc.gov/H1N1flu/recommendations.htm>.

2009 influenza A (H1N1) viruses were tested for oseltamivir resistance by a neuraminidase inhibition assay and/or detection of genetic sequence mutation, depending on the type of specimen tested. Original clinical samples were examined for a single known mutation in the virus that confers oseltamivir resistance in currently circulating seasonal influenza A (H1N1) viruses, while influenza virus isolates were tested using a neuraminidase inhibition assay that determines the presence or absence of neuraminidase inhibitor resistance, followed by neuraminidase gene sequence analysis of resistant viruses.

The majority of 2009 influenza A (H1N1) viruses are susceptible to the neuraminidase inhibitor antiviral medication oseltamivir; however, rare sporadic cases of oseltamivir resistant 2009 influenza A (H1N1) viruses have been detected worldwide. A total of 64 cases of oseltamivir resistant 2009 influenza A (H1N1) viruses have been identified in the United States since April 2009. No new cases have been identified since last week. Fifty-five of these specimens were collected after September 1, 2009. The proportion of oseltamivir-resistant 2009 H1N1 viruses does



**Influenza-Associated Pediatric Mortality:** Three influenza-associated pediatric deaths were reported to CDC during week 14 (New York City and Texas [2]). One death was associated with a seasonal influenza A (H1) virus infection, one was associated with 2009 influenza A (H1N1) virus infection, and one was associated with an influenza A virus for which the subtype was undetermined. The deaths reported during week 14 occurred between May 3, 2009, and March 27, 2010. The death associated with seasonal influenza A (H1) reported during week 14 occurred during the 2008-09 season, bringing the total number of reported pediatric deaths occurring during that season to 134.


Since August 30, 2009, CDC has received 272 reports of influenza-associated pediatric deaths that occurred during the current influenza season (48 deaths in children less than 2 years old, 30 deaths in children 2-4 years old, 102 deaths in children 5-11 years old, and 92 deaths in children 12-17 years old). Two hundred twenty-one (81%) of the 272 deaths were due to 2009 influenza A (H1N1) virus infections, 50 were associated with an influenza A virus for which the subtype is undetermined, and one was associated with an influenza B virus infection. A total of 281 deaths in children associated with 2009 influenza A (H1N1) virus infection have been reported to CDC.

Among the 272 deaths in children, 145 children had specimens collected for bacterial culture from normally sterile sites and 50 (34.5%) of the 145 were positive; *Streptococcus pneumoniae* was identified in 11 (22.0%) of the 50 children, and *Staphylococcus aureus* was identified in 15 (30.0%) of the 50 children. Four *S. aureus* isolates were sensitive to methicillin, ten were methicillin resistant, and one did not have sensitivity testing performed. Thirty-three (66.0%) of the 50 children with bacterial coinfections were five years of age or older, and 17 (34.0%) of the 50 children were 12 years of age or older.

**Laboratory-Confirmed Influenza-Associated Pediatric Deaths by Date and Type/Subtype of Influenza.**

Date	2009 H1N1 Influenza	Influenza A-Subtype Unknown	Seasonal Influenza	Total
<b>Number of Deaths REPORTED for Current Week – Week 14 (Week ending April 10, 2010)</b>	1	1	1	3
<b>Number of Deaths OCCURRED since August 30, 2009</b>	221	50	1	272
<b>Number of Deaths OCCURRED since April 26, 2009</b>	281	53	3	337

## Number of Influenza-Associated Pediatric Deaths by Week of Death: 2006-07 season to present




**Influenza-Associated Hospitalizations:** Laboratory-confirmed influenza-associated hospitalizations are monitored using a population-based surveillance network that includes the 10 Emerging Infections Program (EIP) sites (CA, CO, CT, GA, MD, MN, NM, NY, OR and TN) and 6 new sites (IA, ID, MI, ND, OK and SD).

During September 1, 2009 – April 10, 2010, the following preliminary laboratory-confirmed overall influenza associated hospitalization rates were reported by EIP and the new sites (*rates include influenza A, influenza B, and 2009 influenza A (H1N1)*):

Rates [EIP (new sites)] for children aged 0-4 years and 5-17 years were 6.7 (10.6) and 2.5 (3.6) per 10,000, respectively. Rates [EIP (new sites)] for adults aged 18-49 years, 50-64 years, and ≥ 65 years were 2.4 (1.7), 3.2 (2.0) and 2.7 (1.8) per 10,000, respectively.


# EIP Influenza Laboratory-Confirmed Cumulative Hospitalization Rates, 2009-10 and Previous Three Seasons\*



\* The 2008-09 EIP rate ended as of April 14, 2009 due to the onset of the 2009 H1N1 season.


**Outpatient Illness Surveillance:** Nationwide during week 14, 0.9% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is below the national baseline of 2.3%.

Percentage of Visits for Influenza-like Illness (ILI) Reported by the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Weekly National Summary, October 1, 2006 – April 10, 2010




On a regional level, the percentage of outpatient visits for ILI ranged from 0.5% to 1.3% during week 14. All 10 regions reported ILI below region-specific baseline levels. (Note: Use of the national baseline for regional ILI data or regional baselines for state-level data is not appropriate.)


Region 1 - CT, ME, MA, NH, RI, VT




Region 6 - AR, LA, NM, OK, TX




Region 2 - NJ, NY, USVI




Region 7 - IA, KS, MO, NE




Region 3 - DE, DC, MD, PA, VA, WV




Region 8 - CO, MT, ND, SD, UT, WY




Region 4 - AL, FL, GA, KY, MS, NC, SC, TN




Region 9 - AZ, CA, HI, NV



Region 5 - IL, IN, MI, MN, OH, WI



Region 10 - AK, ID, OR, WA




NOTE: Scales differ between regions

\*Use of the regional baselines for state data is not appropriate.

**Geographic Spread of Influenza as Assessed by State and Territorial Epidemiologists:** The influenza activity reported by state and territorial epidemiologists indicates geographic spread of both seasonal influenza and 2009 influenza A (H1N1) viruses and does not measure the severity of influenza activity.

During week 14, the following influenza activity was reported:

- No states reported widespread influenza activity.
- Regional influenza activity was reported by two states (Alabama and Georgia).
- Local influenza activity was reported by Puerto Rico and one state (Hawaii).
- Sporadic influenza activity was reported by the District of Columbia, Guam, and 37 states (Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Jersey, New Mexico, New York, North Carolina, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming).
- The U.S. Virgin Islands and nine states (Delaware, Idaho, Nebraska, New Hampshire, North Dakota, Oregon, Rhode Island, South Dakota, and Utah) reported no influenza activity.
- One state (Maryland) did not report.



\* This map indicates geographic spread & does not measure the severity of influenza activity

A description of surveillance methods is available at: <http://www.cdc.gov/flu/weekly/fluactivity.htm>  
Report prepared: April 16, 2010.