

2013-2014 Influenza Season Week 51 ending December 21, 2013

All data are preliminary and may change as more reports are received.

Synopsis: During week 51 (December 15-21, 2013), influenza activity continued to increase in the United States.

- **Viral Surveillance:** Of 6,813 specimens tested and reported by U.S. World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NREVSS) collaborating laboratories during week 51, 1,639 (24.1%) were positive for influenza.
- **Pneumonia and Influenza Mortality:** The proportion of deaths attributed to pneumonia and influenza (P&I) was below the epidemic threshold.
- **Influenza-associated Pediatric Deaths:** One influenza-associated pediatric death that occurred during the 2012-2013 season was reported.
- **Influenza-associated Hospitalizations:** A cumulative rate for the season of 4.3 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.
- **Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 3.0%, above the national baseline of 2.0%. Eight regions reported ILI at or above region-specific baseline levels. Six states experienced high ILI activity; eight states experienced moderate ILI activity; six states experienced low ILI activity; 28 states experienced minimal ILI activity, and the District of Columbia, New York City, and two states had insufficient data.
- **Geographic Spread of Influenza:** The geographic spread of influenza in 10 states was reported as widespread; Guam and 23 states reported regional influenza activity; 12 states reported local influenza activity; the District of Columbia, Puerto Rico, and four states reported sporadic influenza activity, and the U.S. Virgin Islands and one state did not report.

National and Regional Summary of Select Surveillance Components

HHS Surveillance Regions*	Data for current week			Data cumulative since September 29, 2013 (Week 40)				
	Out-patient ILI†	% positive for flu‡	Number of jurisdictions reporting regional or widespread activity§	2009 H1N1	A (H3)	A (Subtyping not performed)	B	Pediatric Deaths
Nation	Elevated	24.1%	34 of 54	4,396	284	3,637	457	4
Region 1	Elevated	10.6%	3 of 6	87	15	16	8	0
Region 2	Normal	8.2%	2 of 4	169	18	86	28	0
Region 3	Elevated	13.0%	2 of 6	343	20	4	14	0
Region 4	Elevated	22.6%	7 of 8	598	6	2,083	272	1
Region 5	Elevated	24.9%	4 of 6	485	27	111	13	1
Region 6	Elevated	23.1%	4 of 5	663	45	945	57	2
Region 7	Elevated	17.1%	3 of 4	281	11	24	11	0
Region 8	Elevated	25.5%	3 of 6	1,027	27	290	27	0
Region 9	Normal	14.0%	2 of 5	318	68	54	18	0
Region 10	Elevated	26.0%	4 of 4	425	47	24	9	0

* <http://www.hhs.gov/about/regionmap.html>

† Elevated means the % of visits for ILI is at or above the national or region-specific baseline.


‡ National data are for current week; regional data are for the most recent three weeks.

§ Includes all 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands.

U.S. Virologic Surveillance: WHO and NREVSS collaborating laboratories located in all 50 states, Puerto Rico, and Washington D.C. report to CDC the number of respiratory specimens tested for influenza and the number positive by influenza virus type and influenza A virus subtype. The results of tests performed during the current week are summarized in the table below.

	Week 51
No. of specimens tested	6,813
No. of positive specimens (%)	1,639 (24.1%)
Positive specimens by type/subtype	
Influenza A	1,610 (98.2%)
2009 H1N1	915 (56.8%)
H3	22 (1.4%)
Subtyping not performed	673 (41.8%)
Influenza B	29 (1.8%)

Influenza Positive Tests Reported to CDC by U.S. WHO/NREVSS Collaborating Laboratories, National Summary, 2013-14 Season



Antigenic Characterization*: CDC has antigenically characterized 408 influenza viruses [354 2009 H1N1 viruses, 46 influenza A (H3N2) viruses, and 8 influenza B viruses] collected by U.S. laboratories since October 1, 2013 by hemagglutination inhibition (HI).

- **2009 H1N1 [354]:** All 354 2009 H1N1 viruses tested were characterized as A/California/7/2009-like, the influenza A (H1N1) component of the 2013-2014 Northern Hemisphere influenza vaccine.
- **Influenza A (H3N2) [46]:** All 46 influenza A (H3N2) viruses tested have been characterized as A/Texas/50/2012-like, the influenza A (H3N2) component of the 2013-2014 Northern Hemisphere influenza vaccine.
- **Influenza B [8]:** Three (38%) of the eight influenza B viruses tested belong to B/Yamagata/16/88-lineage and the remaining five (62%) influenza B viruses tested belong to B/Victoria/02/87 lineage.
- **Yamagata Lineage [3]:** Three influenza B/Yamagata-lineage viruses were characterized as B/ Massachusetts/2/1012-like, which is included as an influenza B component of the 2013-2014 Northern Hemisphere trivalent and quadrivalent influenza vaccines.
- **Victoria Lineage [5]:** Five influenza B/Victoria-lineage viruses were characterized as B/Brisbane/60/2008-like, which is included as an influenza B component of the 2013-2014 Northern Hemisphere quadrivalent influenza vaccine.

*For more information see the section on antigenic characterization in the [MMWR "Update: Influenza Activity — United States and Worldwide, May 19–September 28, 2013"](#).

Antiviral Resistance: Testing of 2009 H1N1, influenza A (H3N2), and influenza B virus isolates for resistance to neuraminidase inhibitors (oseltamivir and zanamivir) is performed at CDC using a functional assay. Additional 2009 H1N1 and influenza A (H3N2) clinical samples are tested for mutations of the virus known to confer oseltamivir resistance. The data summarized below combine the results of both testing methods. These samples are routinely obtained for surveillance purposes rather than for diagnostic testing of patients suspected to be infected with antiviral-resistant virus.

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among 2009 influenza A (H1N1) and A (H3N2) viruses (the adamantanes are not effective against influenza B viruses). As a result, data from adamantane resistance testing are not presented below.

**Neuraminidase Inhibitor Resistance Testing Results
on Samples Collected Since October 1, 2013**


	Oseltamivir		Zanamivir	
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)
Influenza A (H3N2)	56	0 (0.0)	56	0 (0.0)
Influenza B	12	0 (0.0)	12	0 (0.0)
2009 H1N1	768*	10 (1.3)	476	0 (0.0)

*Includes specimens tested in national surveillance and additional specimens tested at public health laboratories in 10 states (AZ, CO, FL, HI, MD, MI, NY, TX, WA, and WI) who share testing results with CDC.

The majority of currently circulating influenza viruses are susceptible to the neuraminidase inhibitor antiviral medications, oseltamivir and zanamivir; however, rare sporadic cases of oseltamivir-resistant 2009 H1N1 and A (H3N2) viruses have been detected worldwide. Antiviral treatment with oseltamivir or zanamivir is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for serious influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at <http://www.cdc.gov/flu/antivirals/index.htm>.

Pneumonia and Influenza (P&I) Mortality Surveillance: During week 51, 6.7% of all deaths reported through the 122 Cities Mortality Reporting System were due to P&I. This percentage was below the epidemic threshold of 6.9% for week 51.


Pneumonia and Influenza Mortality for 122 U.S. Cities Week ending December 21, 2013



Influenza-Associated Pediatric Mortality: One influenza-associated pediatric death that occurred during the 2012-2013 season was reported to CDC during week 51 and was associated with an influenza B virus. This death brings the total number of reported pediatric deaths for that season to 171. A total of four influenza-associated pediatric deaths for the 2013-2014 season have been reported.

Additional data can be found at <http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html>.

Number of Influenza-Associated Pediatric Deaths by Week of Death: 2010-11 season to present



Influenza-Associated Hospitalizations: The Influenza Hospitalization Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-related hospitalizations in children younger than 18 years of age (since the 2003-2004 influenza season) and adults (since the 2005-2006 influenza season).


The FluSurv-NET covers more than 70 counties in the 10 Emerging Infections Program (EIP) states (CA, CO, CT, GA, MD, MN, NM, NY, OR, TN) and additional Influenza Hospitalization Surveillance Project (IHSP) states. The IHSP began during the 2009-2010 season to enhance surveillance during the 2009 H1N1 pandemic. IHSP sites included IA, ID, MI, OK and SD during the 2009-2010

season; ID, MI, OH, OK, RI, and UT during the 2010-2011 season; MI, OH, RI, and UT during the 2011-2012 season; IA, MI, OH, RI, and UT during the 2012-2013 season; and MI, OH, and UT during the 2013-2014 season.

Data gathered are used to estimate age-specific hospitalization rates on a weekly basis, and describe characteristics of persons hospitalized with severe influenza illness. The rates provided are likely to be an underestimate as influenza-related hospitalizations can be missed, either because testing is not performed, or because cases may be attributed to other causes of pneumonia or other common influenza-related complications.


Between October 1, 2013 and December 21, 2013, 1,156 laboratory-confirmed influenza-associated hospitalizations were reported. This is a rate of 4.3 per 100,000 population. Among cases, 1,071 (92.6%) were influenza A, 72 (6.2%) were influenza B, 6 (0.5%) were influenza A and B co-infection, and 7 (0.6%) had no virus type information. Among those with influenza A subtype information, 7 (1.6%) were H3 and 420 (97.7%) were 2009 H1N1. The most commonly reported underlying medical conditions among adults were obesity, metabolic disorders, cardiovascular disease, and chronic lung disease (excluding asthma). The most commonly reported underlying medical conditions in children were asthma, obesity, neurologic disorders, and cardiovascular disease. Approximately 45.5% of hospitalized children had no identified underlying medical conditions. Among 50 hospitalized women of childbearing age (15-44 years), eleven were pregnant.

Additional FluSurv-NET data can be found at: <http://gis.cdc.gov/GRASP/Fluview/FluHospRates.html> and <http://gis.cdc.gov/grasp/fluview/FluHospChars.html>.



Data from the Influenza Hospitalization Surveillance Network (FluSurv-NET), a population-based surveillance for influenza related hospitalizations in children and adults in 13 US states. Incidence rates are calculated using the National Center for Health Statistics' (NCHS) population estimates for the counties included in the surveillance catchment area.


Laboratory-Confirmed Influenza Hospitalizations
Preliminary data as of Dec 21, 2013



Asthma includes a medical diagnosis of asthma or reactive airway disease; Cardiovascular diseases include conditions such as coronary heart disease, cardiac valve disorders, congestive heart failure, and pulmonary hypertension, does not include isolated hypertension; Chronic lung diseases include conditions such as chronic obstructive pulmonary disease, bronchiolitis obliterans, chronic aspiration pneumonia, and interstitial lung disease; Immune suppression includes conditions such as immunoglobulin deficiency, leukemia, lymphoma, HIV/AIDS, and individuals taking immunosuppressive medications; Metabolic disorders include conditions such as diabetes mellitus, thyroid dysfunction, adrenal insufficiency, and liver disease; Neurologic diseases include conditions such as seizure disorders, cerebral palsy, and cognitive dysfunction; Neuromuscular diseases include conditions such as multiple sclerosis and muscular dystrophy; Obesity was assigned if indicated in patient's medical chart or if body mass index (BMI) >30 kg/m²; Pregnancy percentage calculated using number of female cases aged between 15 and 44 years of age as the denominator; Renal diseases include conditions such as acute or chronic renal failure, nephrotic syndrome, glomerulonephritis, and impaired creatinine clearance; No known condition indicates that the case did not have any known underlying medical condition indicated in medical chart at the time of hospitalization. Includes only cases for which data collection has been completed through the medical chart review stage.

Outpatient Illness Surveillance: Nationwide during week 51, 3.0% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is above the national baseline of 2.0%. (ILI is defined as fever (temperature of 100 °F [37.8 °C] or greater) and cough and/or sore throat.)











Percentage of Visits for Influenza-like Illness (ILI) Reported by the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Weekly National Summary, 2013-14 and Selected Previous Seasons



On a regional level, the percentage of outpatient visits for ILI ranged from 0.9% to 7.1% during week 51. Eight regions (Regions 1, 3, 4, 5, 6, 7, 8, and 10) reported a proportion of outpatient visits for ILI at or above their region-specific baseline level.

Region specific data is available at <http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html>.





■ ■ ■ ■ ■ % ILI

■ ■ ■ ■ ■ Basehe*

NOTE: Scabs differ between regions

*Use of the regional basehes for stat edat as not pppropiate.




ILINet Activity Indicator Map: Data collected in ILINet are used to produce a measure of ILI activity* by state. Activity levels are based on the percent of outpatient visits in a state due to ILI and are compared to the average percent of ILI visits that occur during weeks with little or no influenza virus circulation. Activity levels range from minimal, which would correspond to ILI activity from outpatient clinics being below, or only slightly above, the average, to high, which would correspond to ILI activity from outpatient clinics being much higher than average.

During week 51, the following ILI activity levels were experienced:

- Six states experienced high ILI activity (Alabama, Louisiana, Mississippi, Missouri, Oklahoma, and Texas).
- Eight states experienced moderate ILI activity (Colorado, Indiana, Kansas, North Carolina, Pennsylvania, South Carolina, Tennessee, and Utah).
- Six states experienced low ILI activity (Alaska, Arizona, Delaware, Florida, Georgia, and Virginia).
- Twenty-eight states experienced minimal ILI activity (California, Connecticut, Hawaii, Illinois, Idaho, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Ohio, Oregon, Rhode Island, South Dakota, Vermont, Washington, West Virginia, Wisconsin, and Wyoming).
- Data were insufficient to calculate an ILI activity level from the District of Columbia, New York City, and two states (Arkansas and Kentucky).

Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet
2013-14 Influenza Season Week 51 ending Dec 21, 2013



*This map uses the proportion of outpatient visits to health care providers for influenza-like illness to measure the ILI activity level within a state. It does not, however, measure the extent of geographic spread of flu within a state. Therefore, outbreaks occurring in a single city could cause the state to display high activity levels.

Data collected in ILINet may disproportionately represent certain populations within a state, and therefore, may not accurately depict the full picture of influenza activity for the whole state.


Data displayed in this map are based on data collected in ILINet, whereas the State and Territorial flu activity map is based on reports from state and territorial epidemiologists. The data presented in this map is preliminary and may change as more data is received.

Differences in the data presented here by CDC and independently by some state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.

Geographic Spread of Influenza as Assessed by State and Territorial Epidemiologists: The influenza activity reported by state and territorial epidemiologists indicates geographic spread of influenza viruses, but does not measure the severity of influenza activity.

During week 51, the following influenza activity was reported:

- Widespread influenza activity was reported by 10 states (Alabama, Alaska, Kansas, Louisiana, Massachusetts, New York, Pennsylvania, Texas, Virginia, and Wyoming).
- Regional influenza activity was reported by Guam and 23 states (Arkansas, Colorado, Connecticut, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kentucky, Maine, Michigan, Minnesota, Mississippi, Nebraska, Nevada, New Jersey, Oklahoma, Oregon, South Carolina, Tennessee, Utah, and Washington).
- Local influenza activity was reported by 12 states (Arizona, California, Delaware, Maryland, Missouri, Montana, New Hampshire, New Mexico, North Dakota, Ohio, South Dakota, and Wisconsin).
- Sporadic influenza activity was reported by the District of Columbia, Puerto Rico, and four states (Hawaii, Rhode Island, Vermont, and West Virginia).
- The U.S. Virgin Islands and one state (North Carolina) did not report.



* This map indicates geographic spread & does not measure the severity of influenza activity

A description of surveillance methods is available at: <http://www.cdc.gov/flu/weekly/overview.htm>
Report prepared: December 27, 2013.

Additional National and International Influenza Surveillance Information

FluView Interactive: FluView includes enhanced web-based interactive applications that can provide dynamic visuals of the influenza data collected and analyzed by CDC. These FluView Interactive applications allow people to create customized, visual interpretations of influenza data, as well as comparisons across flu seasons, regions, age groups and a variety of other demographics. To access these tools visit www.cdc.gov/flu/weekly/fluviewinteractive.htm.

U.S. State and local influenza surveillance: Click on a jurisdiction below to access the latest local influenza information.

Alabama	Alaska	Arizona	Arkansas	California
Colorado	Connecticut	Delaware	District of Columbia	Florida
Georgia	Hawaii	Idaho	Illinois	Indiana
Iowa	Kansas	Kentucky	Louisiana	Maine
Maryland	Massachusetts	Michigan	Minnesota	Mississippi
Missouri	Montana	Nebraska	Nevada	New Hampshire
New Jersey	New Mexico	New York	North Carolina	North Dakota
Ohio	Oklahoma	Oregon	Pennsylvania	Rhode Island
South Carolina	South Dakota	Tennessee	Texas	Utah
Vermont	Virginia	Washington	West Virginia	Wisconsin
Wyoming	New York City	Virgin Islands		

Google Flu Trends: Google Flu Trends uses aggregated Google search data in a model created in collaboration with CDC to estimate influenza activity in the United States. For more information and activity estimates from the U.S. and worldwide, see <http://www.google.org/flutrends/>.

World Health Organization: Additional influenza surveillance information from participating WHO member nations is available through [FluNet](#) and the [Global Epidemiology Reports](#).

WHO Collaborating Centers for Influenza located in [Australia](#), [China](#), [Japan](#), and the [United Kingdom](#).

Europe: WHO/Europe at <http://www.euroflu.org/index.php> and the European Centre for Disease Prevention and Control at http://ecdc.europa.eu/en/publications/surveillance_reports/influenza/Pages/weekly_influenza_surveillance_overview.aspx.

Public Health Agency of Canada: The most up-to-date influenza information from Canada is available at <http://www.phac-aspc.gc.ca/fluwatch/>.

Health Protection Agency (United Kingdom): The most up-to-date influenza information from the United Kingdom is available at <http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/>.

Any links provided to non-Federal organizations are provided solely as a service to our users. These links do not constitute an endorsement of these organizations or their programs by CDC or the Federal Government, and none should be inferred. CDC is not responsible for the content of the individual organization web pages found at these links.