

**Table 4a. Number of laboratory-confirmed bacterial and parasitic infections in 2010\*, by pathogen and age group, Foodborne Diseases Active Surveillance Network (FoodNet), United States**

Pathogen	Age group (yrs)					Total
	<5	5-9	10-19	20-59	≥60	
<b>Bacteria</b>						
<i>Campylobacter</i>	778	330	634	3,471	1,148	6,361
<i>Listeria</i> <sup>†</sup>	8	1	3	26	87	125
<i>Salmonella</i>	2,217	663	772	3,189	1,409	8,250
<i>Shigella</i>	524	364	137	661	91	1,777
STEC <sup>§</sup> O157	105	77	68	135	57	442
STEC non-O157	160	35	81	133	42	451
<i>Vibrio</i>	0	8	11	107	67	193
<i>Yersinia</i>	60	13	14	40	32	159
<b>Parasites</b>						
<i>Cryptosporidium</i>	162	85	157	676	210	1,290
<i>Cyclospora</i>	0	0	1	22	5	28
<b>Total</b>	<b>4,014</b>	<b>1,576</b>	<b>1,878</b>	<b>8,460</b>	<b>3,148</b>	<b>19,076</b>

\*Data are preliminary. 13 records with missing age not included.

<sup>†</sup>*Listeria* cases defined as isolation of *L. monocytogenes* from a normally sterile site or, in the setting of miscarriage or stillbirth, isolation of *L. monocytogenes* from placental or fetal tissue. In cases of pregnancy-associated listeriosis, case may be mother, infant, or both, depending on source(s) of isolation.

<sup>§</sup>Shiga toxin producing *Escherichia coli*