


ANNUAL SUMMARY 1982

ISSUED SEPTEMBER 1985

CENTERS FOR DISEASE CONTROL  
**FOODBORNE DISEASE**

**SURVEILLANCE**



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES • Public Health Service

PREFACE

This report summarizes information received from state and local health departments, the Food and Drug Administration, the U.S. Department of Agriculture, and private physicians. The information is preliminary and is intended primarily for use by those with responsibility for disease control activities. Anyone desiring to quote this report should contact the Enteric Diseases Branch for confirmation and further interpretation.

Contributions to the report are most welcome. Please address them to

Enteric Diseases Branch  
Division of Bacterial Diseases  
Center for Infectious Diseases  
Centers for Disease Control  
Atlanta, Georgia 30333

SUGGESTED CITATION

Centers for Disease Control: Foodborne Disease Outbreaks  
Annual Summary 1982, Issued September 1985

Centers for Disease Control . . . . . James O. Mason, M.D., Dr. P.H., Director  
Center for Infectious Diseases. . . . . Walter R. Dowdle, Ph.D., Director  
Bacterial Diseases Division . . . . . John C. Feeley, Ph.D., Director  
Enteric Diseases Branch . . . . . Paul A. Blake, M.D., M.P.H., Chief  
Foodborne Disease Outbreaks  
Surveillance Activity . . . . . Kristine L. MacDonald, M.D.\*  
Report preparation. . . . . Dorothy S. Anderson  
Enteric Bacteriology Section. . . . . George K. Morris, Ph.D., Chief  
Statistical Services Branch . . . . . Stanley M. Martin, M.S., Chief  
Nancy Hargrett, Ph.D.  
Barbara Fowler  
Publications and Graphics . . . . . Frances H. Porcher, M.A., Chief  
J. Philip Thompson, M.S., Writer-Editor

\* Through June 1985

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the U.S. Department of Health and Human Services.

## TABLE OF CONTENTS

	<u>Page</u>
I. BACKGROUND	1
A. History	1
B. Objectives	1
II. FOODBORNE DISEASE OUTBREAKS	2
A. Definition of Outbreak	2
B. Source of Data	2
C. Interpretation of Data	2
D. Summary of Data	3
E. Comments	4
F. Guidelines for Confirmation of Foodborne Outbreaks	13
G. Investigation of a Foodborne Outbreak (Sample Form)	17
H. Line Listing of Foodborne Disease Outbreaks, 1982	19
I. Selected Foodborne Outbreak Articles, 1982, Taken from Morbidity and Mortality Weekly Report	25
J. Bibliography	33



## I. BACKGROUND

### A. History

The reporting of foodborne and waterborne diseases in the United States began over half a century ago when state and territorial health officers, concerned about the high morbidity and mortality caused by typhoid fever and infantile diarrhea, recommended that cases of enteric fever be investigated and reported. The purpose was to obtain information about the role of food, milk, and water in outbreaks of intestinal illness as the basis for sound public health action. Beginning in 1923, the U.S. Public Health Service published summaries of outbreaks of gastrointestinal illness attributed to milk. In 1938, it added summaries of outbreaks caused by all foods. These early surveillance efforts led to the enactment of important public health measures which had a profound influence in decreasing the incidence of enteric diseases, particularly those transmitted by milk and water.

From 1951 through 1960, the National Office of Vital Statistics reviewed reports of outbreaks of foodborne illness and published summaries of them annually in Public Health Reports. In 1961 the Centers for Disease Control (CDC), then the Communicable Disease Center, assumed responsibility for publishing reports on foodborne illness. For the period 1961-65, CDC discontinued publication of annual reviews, but reported pertinent statistics and detailed individual investigations in the Morbidity and Mortality Weekly Report (MMWR).

In 1966 the present system of surveillance of foodborne and waterborne diseases began with the incorporation of all reports of enteric disease outbreaks attributed to microbial or chemical contamination of food or water into an annual summary. Since 1966 the quality of investigative reports has improved primarily as a result of more active participation by state and federal agencies in the investigation of foodborne and waterborne disease outbreaks. Due to increasing interest and activity in waterborne disease surveillance, foodborne and waterborne disease outbreaks have been reported in separate annual summaries since 1978. This report summarizes data from foodborne disease outbreaks reported to CDC for 1982.

### B. Objectives

Foodborne disease surveillance has traditionally served 3 objectives:

1. Disease Prevention and Control: Early identification and removal of contaminated products from the commercial market, correction of faulty food preparation practices in food service establishments and in the home, and identification and appropriate treatment of human carriers of foodborne pathogens are the fundamental prevention and control measures resulting from surveillance of foodborne disease.

2. Knowledge of Disease Causation: The responsible pathogen was not identified in over half of the foodborne disease outbreaks reported to CDC in each of the last 5 years. In many of these outbreaks pathogens known to cause foodborne illness may not have been identified because of late or incomplete laboratory investigation. In others, the responsible pathogen may have escaped detection even when a thorough laboratory investigation was carried out because the pathogen may not have been appreciated as a cause of foodborne disease or because the pathogen could not be identified by available laboratory techniques. When more thorough clinical, epidemiologic, and laboratory investigations are employed, perhaps many of these pathogens can be identified, and suitable measures for prevention and control can be instituted.

3. Administrative Guidance: The collection of data from outbreak investigations permits assessment of trends in etiologic agents and food vehicles and focuses on common errors in food handling. By compiling the data in an annual summary, it is hoped that local and state health departments and others involved in the implementation of food protection programs will be kept informed of the factors involved in foodborne disease outbreaks. Comprehensive surveillance would result in a clearer appreciation of priorities in food protection, institution of better training programs, and more effective use of available resources.

## II. FOODBORNE DISEASE OUTBREAKS

### A. Definition of Outbreak

For the purpose of this report, a foodborne disease outbreak is defined as an incident in which (1) 2 or more persons experience a similar illness, after ingestion of a common food, and (2) epidemiologic analysis implicates the food as the source of the illness. A few exceptions exist; for example, 1 case of botulism or chemical poisoning constitutes an outbreak.

Outbreaks of known etiology are those in which laboratory evidence of a specific etiologic agent is obtained, and specified criteria are met (see Section F). Outbreaks of unknown etiology are those in which epidemiologic evidence implicates a food source, but adequate laboratory confirmation is not obtained. These outbreaks are subdivided into 4 subgroups by incubation period of the illnesses: less than 1 hour (probable chemical poisoning), 1 to 7 hours (probable Staphylococcus food poisoning), 8 to 14 hours (probable C. perfringens food poisoning), and greater than 14 hours (other infectious or toxic agents).

### B. Source of Data

Outbreaks are reported to CDC on a standard reporting form (Section G). Reports come most frequently from state and local health departments; reports may also be received from federal agencies such as the Food and Drug Administration (FDA), U.S. Department of Agriculture (USDA), the U.S. Armed Forces, and occasionally from private physicians. Forms are reviewed at CDC to determine if a specific etiologic agent for the outbreak can be confirmed and, in some instances, questions about an etiologic agent may be referred back to the reporting agency. Otherwise, data are otherwise accepted as reported on the forms.

### C. Interpretation of Data

The limitations on the quantity and quality of data presented here must be appreciated to avoid misinterpretation. The number of outbreaks of foodborne disease reported by this surveillance system clearly represents only a small fraction of the total number that occur. The likelihood of an outbreak coming to the attention of health authorities varies considerably depending on consumer and physician awareness, interest, and motivation to report the incident; for example, large outbreaks, interstate outbreaks, restaurant-associated outbreaks, and outbreaks involving serious illness, hospitalizations, or deaths are more likely to come to the attention of health authorities than cases of mild illness following a family cookout.

The quality of the data presented here depends upon the commitment to foodborne surveillance by the state or local health departments. The department's interest in foodborne disease and its investigative and laboratory capabilities are central determinants of the quality of the investigation. Furthermore, the likelihood that the findings of the investigation will be reported varies from one locality to another. This report then should not be the basis of firm conclusions about the absolute incidence of foodborne disease, and it should not be used to draw conclusions about the relative incidence of foodborne diseases of various etiologies. For example, foodborne diseases characterized by short incubation periods, such as those

caused by a chemical agent or staphylococcal enterotoxin, are more likely to be recognized as common-source foodborne disease outbreaks than those diseases with longer incubation periods, such as hepatitis A. Outbreaks involving less common pathogens, such as Bacillus cereus, Escherichia coli, or Giardia lamblia are less likely to be confirmed because these organisms are often not considered in clinical, epidemiologic, and laboratory investigations of foodborne disease outbreaks. Also, pathogens which generally cause mild illness will be under-represented, while those causing serious illness, such as Clostridium botulinum, are more likely to be identified. Similarly, restaurant- or commercial product-associated outbreaks have a higher likelihood of being reported.

D. Summary

In 1982, 656 outbreaks (19,380 cases) of foodborne disease were reported to the Centers for Disease Control. Reports were received from 39 states, as well as from the District of Columbia, the U.S. Virgin Islands, Guam, and Puerto Rico (Figure 1). New York reported the largest number of outbreaks, 323, with 160 from New York City; Washington reported the next largest number of outbreaks (44), followed by California (35).

Fig.1 OUTBREAKS OF FOODBORNE DISEASE REPORTED TO THE CENTERS FOR DISEASE CONTROL, BY STATE, 1982



The etiologic agent was confirmed in 34% of the outbreaks. Bacterial pathogens accounted for 151 outbreaks (5,501 cases). The most frequently isolated bacterial pathogen was Salmonella (55 outbreaks, 2,056 cases), followed by Staphylococcus aureus (28 outbreaks, 669 cases), and Clostridium perfringens (22 outbreaks, 1,189 cases). During 1982, two outbreaks (one in Oregon and one in Michigan) of a previously unrecognized pathogen, E. coli O157:H7, were investigated. Both outbreaks were associated with eating hamburger from the same fast-food restaurant chain. The illness was characterized by bloody diarrhea, abdominal cramps and low-grade or absent fever. Since first described, this pathogen has also been associated with non-bloody diarrhea and a spectrum of clinical illness, including hemolytic-uremic syndrome. Viral agents (Hepatitis A and Norwalk virus) accounted for 21 outbreaks and 5,325 cases. The high number of cases resulting from viral agents is predominantly due to 2 large outbreaks of Norwalk gastroenteritis which occurred in Minnesota. One outbreak involved 3,000 cases and was related to eating bakery items with frosting. The second outbreak involved 2,000 cases and was associated with eating cole slaw. Chemical agents were responsible for 47 outbreaks (220 cases). Only 1 outbreak, involving 4 persons with Trichinella spiralis, was attributed to a parasitic agent. Twenty-six deaths were associated with foodborne diseases in 1982: 11 from Vibrio cholerae O1 (all in Guam and the Trust Territories), 8 from Salmonella, 5 from Clostridium botulinum, and 2 from unknown etiologies.

No pathogen was identified in 436 of the outbreaks (8,330 cases) reported in 1982. Incubation periods were known for illnesses in 412 of the outbreaks. In 13 outbreaks, the incubation period was less than 1 hour, in 123 outbreaks, it was 1 to 7 hours, in 91 outbreaks, it was 8 to 14 hours, and in 185 outbreaks, it was 15 hours or more.

#### E. Comments

There are limitations in the quantity and quality of the data presented in this report. The variability in reporting can be seen by looking at the distribution of outbreaks by state. A few states, such as New York, Washington, and California, reported a disproportionately large number of outbreaks. For example, New York State and New York City reported 323 outbreaks, almost one-half of those reported for the entire United States. While it is possible that states such as New York, Washington, and California have an increased rate of foodborne disease, it is more likely that these differences simply represent differences in surveillance activity. The same variability in reporting can be seen when looking at outbreaks by pathogen. Our data show that C. botulinum is as common a foodborne pathogen as C. jejuni and V. parahaemolyticus, a conclusion that can only be explained by more complete reporting for botulism than for some other pathogens.

The number of outbreaks of foodborne disease of confirmed etiology reported to CDC over the last 5 years has remained relatively constant. The distribution of cases by etiologic agent has also remained fairly constant. Etiologic agents typically have been confirmed in about 40% of outbreaks. When etiologic agents have been confirmed, bacterial pathogens have consistently accounted for approximately two-thirds of outbreaks, with chemical agents accounting for an additional 20%-25%. Many factors contribute to foodborne disease. In 1982, the 5 most common factors, in order of frequency of occurrence, included: 1) improper holding temperature, 2) food from an unsafe source, 3) inadequate cooking, 4) poor personal hygiene, 5) contaminated equipment. In most of the outbreaks caused by a bacterial pathogen, the food had been stored at improper holding temperatures. In outbreaks of botulism or trichinosis, the food had usually been inadequately cooked. In outbreaks of ciguatera and mushroom poisoning, the food itself was unsafe, and illness was not in any sense related to improper handling or preparation.

The large number of outbreaks in which no pathogen was identified serves as a challenge to improve investigative skills so that known pathogens can be identified more frequently and new and as-yet unidentified pathogens may be recognized.

No.	Name	Description				Remarks
		Quantity	Value	Weight	Measure	
1	...	...	...	...	...	
2	...	...	...	...	...	
3	...	...	...	...	...	
4	...	...	...	...	...	
5	...	...	...	...	...	
6	...	...	...	...	...	
7	...	...	...	...	...	
8	...	...	...	...	...	
9	...	...	...	...	...	
10	...	...	...	...	...	
11	...	...	...	...	...	
12	...	...	...	...	...	
13	...	...	...	...	...	
14	...	...	...	...	...	
15	...	...	...	...	...	
16	...	...	...	...	...	
17	...	...	...	...	...	
18	...	...	...	...	...	
19	...	...	...	...	...	
20	...	...	...	...	...	
21	...	...	...	...	...	
22	...	...	...	...	...	
23	...	...	...	...	...	
24	...	...	...	...	...	
25	...	...	...	...	...	
26	...	...	...	...	...	
27	...	...	...	...	...	
28	...	...	...	...	...	
29	...	...	...	...	...	
30	...	...	...	...	...	
31	...	...	...	...	...	
32	...	...	...	...	...	
33	...	...	...	...	...	
34	...	...	...	...	...	
35	...	...	...	...	...	
36	...	...	...	...	...	
37	...	...	...	...	...	
38	...	...	...	...	...	
39	...	...	...	...	...	
40	...	...	...	...	...	
41	...	...	...	...	...	
42	...	...	...	...	...	
43	...	...	...	...	...	
44	...	...	...	...	...	
45	...	...	...	...	...	
46	...	...	...	...	...	
47	...	...	...	...	...	
48	...	...	...	...	...	
49	...	...	...	...	...	
50	...	...	...	...	...	

Table 1 Confirmed Foodborne Disease Outbreaks, Cases, and Deaths, by Etiologic Agents, United States, 1982

<u>Etiologic Agent</u>	<u>Outbreaks</u>		<u>Cases</u>		<u>Deaths</u>	
	<u>No.</u>	<u>(%)</u>	<u>No.</u>	<u>(%)</u>	<u>No.</u>	<u>(%)</u>
<u>BACTERIAL</u>						
<u>Bacillus cereus</u>	8	(3.6)	200	(1.8)	0	(0.0)
<u>Brucella</u>	1	(0.5)	3	(<0.1)	0	(0.0)
<u>Campylobacter jejuni</u>	2	(0.9)	31	(0.3)	0	(0.0)
<u>Clostridium botulinum</u>	21	(9.5)	30	(0.3)	5	(20.9)
<u>Clostridium perfringens</u>	22	(10.0)	1,189	(10.8)	0	(0.0)
<u>Escherichia coli</u>	2	(0.9)	47	(0.4)	0	(0.0)
<u>Salmonella</u>	55	(25.0)	2,056	(18.6)	8	(33.3)
<u>Shigella</u>	4	(1.8)	116	(1.1)	0	(0.0)
<u>Staphylococcus aureus</u>	28	(12.7)	669	(6.0)	0	(0.0)
<u>Streptococcus Group A</u>	1	(0.5)	34	(0.3)	0	(0.0)
<u>Vibrio cholerae 01</u>	1	(0.5)	892	(8.0)	11	(45.8)
<u>Vibrio cholerae non-01</u>	1	(0.5)	7	(0.1)	0	(0.0)
<u>Vibrio parahaemolyticus</u>	3	(1.4)	39	(0.4)	0	(0.0)
<u>Yersinia enterocolitica</u>	2	(0.9)	188	(1.7)	0	(0.0)
Total	151	(68.7)	5,501	(49.9)	24	(100.0)
<u>CHEMICAL</u>						
Ciguatoxin	8	(3.6)	37	(0.3)	0	(0.0)
Heavy metals	5	(2.3)	26	(0.2)	0	(0.0)
Monosodium glutamate	3	(1.4)	10	(0.1)	0	(0.0)
Mushroom poisoning	4	(1.8)	9	(0.1)	0	(0.0)
Scombrototoxin	18	(8.2)	58	(0.5)	0	(0.0)
Paralytic shellfish	1	(0.5)	5	(<0.1)	0	(0.0)
Other	8	(3.6)	75	(0.7)	0	(0.0)
Total	47	(21.4)	220	(1.9)	0	(0.0)
<u>PARASITIC</u>						
<u>Trichinella spiralis</u>	1	(0.5)	4	(<0.1)	0	(0.0)
Total	1	(0.5)	4	(<0.1)	0	(0.0)
<u>VIRAL</u>						
Hepatitis A	19	(8.5)	325	(2.9)	0	(0.0)
Norwalk Virus	2	(0.9)	5,000	(45.2)	0	(0.0)
Total	21	(9.4)	5,325	(48.1)	0	(0.0)
CONFIRMED TOTAL	220	(100.0)	11,050	(100.0)	24	(100.0)

Table 2 Confirmed Foodborne Disease Outbreaks, by Etiologic Agent, United States, 1978-1982

Etiologic Agent	1978		1979		1980		1981		1982	
	No.	(%)								
<b><u>BACTERIAL</u></b>										
<u>B. cereus</u>	6	(3.9)	-	-	9	(4.1)	8	(3.2)	8	(3.6)
<u>Brucella</u>	-	-	2	(1.3)	-	-	-	-	1	(0.5)
<u>C. jejuni</u>	-	-	-	-	5	(2.3)	10	(4.0)	2	(0.9)
<u>C. botulinum</u>	12	(7.8)	7	(4.0)	14	(6.3)	11	(4.4)	21	(9.5)
<u>C. perfringens</u>	9	(5.9)	20	(11.6)	25	(11.3)	28	(11.2)	22	(10.0)
<u>E. cloacae</u>	-	-	1	(0.6)	-	-	-	-	-	-
<u>E. coli</u>	1	(0.7)	-	-	1	(0.5)	-	-	2	(0.9)
<u>Salmonella</u>	45	(29.3)	44	(25.5)	39	(17.7)	66	(26.4)	55	(25.0)
<u>Shigella</u>	4	(2.6)	7	(4.0)	11	(5.0)	9	(3.6)	4	(1.8)
<u>S. aureus</u>	23	(14.9)	34	(19.7)	27	(12.2)	44	(17.6)	28	(12.7)
<u>Streptococcus Group A</u>	-	-	-	-	-	-	2	(0.8)	1	(0.5)
<u>Streptococcus Group D</u>	1	(0.7)	-	-	-	-	1	(0.4)	-	-
<u>Streptococcus Group G</u>	-	-	1	(0.6)	-	-	-	-	-	-
<u>V. cholerae 01</u>	1	(0.7)	-	-	-	-	-	-	1	(0.5)
<u>V. cholerae non-01</u>	-	-	1	(0.6)	-	-	1	(0.4)	1	(0.5)
<u>V. parahaemolyticus</u>	2	(1.3)	2	(1.3)	4	(1.8)	2	(0.8)	3	(1.4)
<u>Y. enterocolitica</u>	-	-	-	-	-	-	2	(0.8)	2	(0.9)
Other	1	(0.7)	-	-	1	(0.5)	1	(0.4)	-	-
Total	105	(68.5)	119	(69.2)	136	(61.7)	185	(74.0)	151	(68.7)
<b><u>CHEMICAL</u></b>										
Ciguatoin	19	(12.3)	18	(10.4)	15	(6.8)	15	(6.0)	8	(3.6)
Heavy metals	1	(0.7)	1	(0.6)	1	(0.5)	2	(0.8)	5	(2.3)
Monosodium glutamate	-	-	-	-	-	-	2	(0.8)	3	(1.4)
Mushroom poisoning	1	(0.6)	1	(0.6)	-	-	11	(4.4)	4	(1.8)
Paralytic shellfish	4	(2.6)	-	-	5	(2.2)	-	-	1	(0.5)
Scombrototoxin	7	(4.5)	12	(7.0)	29	(13.0)	7	(2.8)	18	(8.2)
Other	5	(3.2)	4	(2.3)	16	(7.2)	14	(5.6)	8	(3.6)
Total	37	(23.9)	36	(20.9)	66	(29.7)	51	(20.4)	47	(21.4)
<b><u>PARASITIC</u></b>										
<u>Trichinella spiralis</u>	7	(4.5)	11	(6.4)	5	(2.3)	7	(2.8)	1	(0.5)
Other	-	-	-	-	2	(0.9)	1	(0.4)	-	-
Total	7	(4.5)	11	(6.4)	7	(3.2)	8	(3.2)	1	(0.5)
<b><u>VIRAL</u></b>										
Hepatitis A	5	(3.2)	5	(2.9)	10	(4.5)	6	(2.4)	19	(8.6)
Norwalk Virus	-	-	1	(0.6)	2	(0.9)	-	-	2	(0.9)
Total	5	(3.2)	6	(3.5)	12	(5.4)	6	(2.4)	21	(9.4)
CONFIRMED TOTAL	154	(100.0)	172	(100.0)	221	(100.0)	250	(100.0)	220	(100.0)

Table 3 Foodborne Outbreaks by Specific Etiologic Agent and Vehicle of Transmission, United States, 1982

<u>Etiologic Agent</u>	<u>Beef</u>	<u>Ham</u>	<u>Pork</u>	<u>Sau- sage</u>	<u>Chick- en</u>	<u>Tur- key</u>	<u>Other Meat</u>	<u>Shell Fish</u>	<u>Tuna</u>	<u>Mahi- Mahi</u>	<u>Other Fish</u>	<u>Milk</u>	<u>Eggs</u>	<u>Ice Cre</u>
<u>BACTERIAL</u>														
<u>B. cereus</u>	1	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Brucella</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>C. jejuni</u>	-	-	-	-	-	-	-	-	-	-	-	2	-	-
<u>C. botulinum</u>	1	-	-	-	-	-	-	-	-	-	1	-	-	-
<u>C. perfringens</u>	6	-	-	-	-	4	2	-	-	-	-	-	-	-
<u>E. coli</u>	2	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Salmonella</u>	1	-	2	1	2	4	1	-	-	-	1	1	2	-
<u>Shigella</u>	-	-	-	-	-	-	-	-	1	-	-	-	-	-
<u>S. aureus</u>	5	3	2	1	4	3	1	-	-	-	-	-	-	-
<u>Streptococcus Group A</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>V. cholerae 01</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>V. cholerae non-01</u>	-	-	-	-	-	-	-	1	-	-	-	-	-	-
<u>V. parahaemolyticus</u>	-	-	-	-	-	-	-	3	-	-	-	-	-	-
<u>Y. enterocolitica</u>	-	-	-	-	-	-	-	-	-	-	-	1	-	-
Other	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total	16	3	4	2	6	11	4	4	1	-	2	4	2	-
<u>CHEMICAL</u>														
Ciguatoxin	-	-	-	-	-	-	-	-	-	-	8	-	-	-
Heavy metals	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Monosodium glutamate	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mushroom poisoning	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Paralytic shellfish	-	-	-	-	-	-	-	1	-	-	-	-	-	-
Scombrototoxin	-	-	-	-	-	-	-	-	3	5	10	-	-	-
Other	1	-	-	1	-	-	-	-	-	-	-	-	-	-
Total	1	-	-	1	-	-	-	1	3	5	18	-	-	-
<u>PARASITIC</u>														
<u>Trichinella spiralis</u>	-	-	1	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	1	-	-	-	-	-	-	-	-	-	-	-
<u>VIRAL</u>														
Hepatitis A	-	-	-	-	-	-	-	5	1	-	-	-	-	-
Norwalk Virus	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	-	-	-	5	1	-	-	-	-	-
CONFIRMED TOTAL	17	3	5	3	6	11	4	10	5	5	20	4	2	-
UNKNOWN	5	2	1	1	7	-	-	56	2	-	1	-	1	-
TOTAL 1982	22	5	6	4	13	11	4	66	7	5	21	4	3	-

Table 3 Foodborne Outbreaks by Specific Etiologic Agent and Vehicle of Transmission, U.S., 1982 (Contd)

<u>Baked Foods</u>	<u>Fruits &amp; Vegetable</u>	<u>Potato Salad</u>	<u>Poultry, Fish, Egg Salad</u>	<u>Other Salad</u>	<u>Fried Rice</u>	<u>Chinese Food</u>	<u>Mexican Food</u>	<u>Carbonated Bev</u>	<u>Non-Dairy Bev</u>	<u>Multiple Foods</u>	<u>Mushrooms</u>	<u>Other Foods</u>	<u>Unknown</u>	<u>Total</u>
1	-	-	-	1	3	-	-	-	-	-	-	1	-	8
-	-	-	-	-	-	-	-	-	-	-	-	-	1	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	2
-	16	-	-	-	-	-	-	-	-	-	1	1	1	21
-	-	-	-	-	-	-	2	-	-	1	-	-	7	22
-	-	-	-	-	-	-	-	-	-	-	-	-	-	2
-	-	-	1	2	-	-	1	-	-	-	-	3	29	55
-	-	-	-	1	-	-	-	-	-	-	-	-	2	4
2	-	2	1	-	-	-	1	-	-	1	-	-	2	28
-	-	-	-	1	-	-	-	-	-	-	-	-	-	1
-	-	-	-	-	-	-	-	-	-	1	-	-	-	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	3
-	1	-	-	-	-	-	-	-	-	-	-	-	-	2
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>3</u>	<u>17</u>	<u>2</u>	<u>2</u>	<u>5</u>	<u>3</u>	<u>-</u>	<u>4</u>	<u>-</u>	<u>-</u>	<u>3</u>	<u>1</u>	<u>5</u>	<u>42</u>	<u>151</u>
-	-	-	-	-	-	-	-	-	-	-	-	-	-	8
-	-	-	-	-	-	-	-	4	1	-	-	-	-	5
-	-	-	-	-	-	2	-	-	-	1	-	-	-	3
-	-	-	-	-	-	-	-	-	-	-	4	-	-	4
-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	18
<u>-</u>	<u>1</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>2</u>	<u>-</u>	<u>1</u>	<u>2</u>	<u>1</u>	<u>4</u>	<u>3</u>	<u>-</u>	<u>8</u>
<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>47</u>
<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>
<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>
1	-	-	-	-	-	-	-	-	-	1	-	-	10	19
<u>1</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>	<u>-</u>	<u>-</u>	<u>10</u>	<u>21</u>
5	18	2	2	6	3	2	4	5	2	5	5	8	52	220
5	1	3	1	6	1	1	2	-	-	5	1	9	325	436
10	19	5	3	12	4	3	6	5	2	10	6	17	377	656

Table 4 Foodborne Disease Outbreaks, by Specific Etiologic Agent and Place Where Food Was Eaten, United States, 1982

Etiologic Agent	Delicatessen, Cafeteria, or Restaurant							Un-known	Total
	Home	School	Picnic	Church	Camp	Other			
<b><u>BACTERIAL</u></b>									
<u>B. cereus</u>	-	3	2	-	-	1	2	-	8
<u>Brucella</u>	1	-	-	-	-	-	-	-	1
<u>C. jejuni</u>	1	-	-	-	-	-	1	-	2
<u>C. botulinum</u>	21	-	-	-	-	-	-	-	21
<u>C. perfringens</u>	-	9	1	-	-	-	12	-	22
<u>E. coli</u>	-	-	-	-	-	-	2	-	2
<u>Salmonella</u>	12	17	6	3	4	1	12	-	55
<u>Shigella</u>	1	2	-	-	1	-	-	-	4
<u>S. aureus</u>	4	7	2	2	2	1	10	-	28
<u>Streptococcus Group A</u>	-	-	-	-	-	-	1	-	1
<u>V. cholerae 01</u>	1	-	-	-	-	-	-	-	1
<u>V. cholerae non-01</u>	1	-	-	-	-	-	-	-	1
<u>V. parahaemolyticus</u>	1	-	-	1	-	-	1	-	3
<u>Y. enterocolitica</u>	1	-	-	-	-	-	1	-	2
Total	44	38	11	6	7	3	42	-	151
<b><u>CHEMICAL</u></b>									
Ciguatoxin	3	5	-	-	-	-	-	-	8
Heavy metals	1	3	-	-	-	-	1	-	5
Monosodium glutamate	1	2	-	-	-	-	-	-	3
Mushroom poisoning	3	-	-	-	-	-	1	-	4
Paralytic shellfish	1	-	-	-	-	-	-	-	1
Scombrototoxin	7	9	-	-	-	-	2	-	18
Other chemical	4	4	-	-	-	-	-	-	8
Total	20	23	-	-	-	-	4	-	47
<b><u>PARASITIC</u></b>									
<u>Trichinella spiralis</u>	1	-	-	-	-	-	-	-	1
Total	1	-	-	-	-	-	-	-	1
<b><u>VIRAL</u></b>									
Hepatitis A	5	6	-	1	-	1	6	-	19
Norwalk Virus	-	1	-	-	-	-	1	-	2
Total	5	7	-	1	-	1	7	-	21
CONFIRMED TOTAL	70	68	11	7	7	4	53	-	220
UNKNOWN	127	221	17	16	2	1	49	3	436
TOTAL 1982	197	289	28	23	9	5	102	3	656

Table 5 Foodborne Disease Outbreaks by Specific Etiologic Agent and Month of Occurrence, United States, 1982

<u>Etiologic Agent</u>	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>	<u>Unknown</u>	<u>Total</u>
<u>BACTERIAL</u>														
<u>Brucella</u>	-	-	1	-	-	-	-	-	-	-	-	-	-	1
<u>B. cereus</u>	2	-	-	-	-	-	2	2	2	-	-	-	-	8
<u>C. jejuni</u>	-	-	-	-	-	1	-	-	1	-	-	-	-	2
<u>C. botulinum</u>	-	3	-	-	2	3	3	2	1	1	3	3	-	21
<u>C. perfringens</u>	-	2	1	1	5	2	1	2	2	1	2	3	-	22
<u>E. coli</u>	-	1	-	-	1	-	-	-	-	-	-	-	-	2
<u>Salmonella</u>	2	1	1	5	4	5	10	11	3	5	3	5	-	55
<u>Shigella</u>	-	-	-	1	-	-	-	-	1	1	1	-	-	4
<u>S. aureus</u>	2	1	1	2	2	3	5	5	1	3	1	2	-	28
<u>Streptococcus Group A</u>	-	-	-	-	-	-	1	-	-	-	-	-	-	1
<u>V. cholerae 01</u>	-	-	-	-	-	-	-	1	-	-	-	-	-	1
<u>V. cholerae non-01</u>	-	-	-	-	-	-	-	-	-	-	-	1	-	1
<u>V. parahaemolyticus</u>	-	-	-	-	-	-	2	1	-	-	-	-	-	3
<u>Y. enterocolitica</u>	-	1	-	-	-	1	-	-	-	-	-	-	-	2
<u>Total</u>	<u>6</u>	<u>9</u>	<u>4</u>	<u>9</u>	<u>14</u>	<u>15</u>	<u>24</u>	<u>24</u>	<u>11</u>	<u>11</u>	<u>10</u>	<u>14</u>	<u>-</u>	<u>151</u>
<u>CHEMICAL</u>														
<u>Ciguatoxin</u>	1	-	-	-	-	-	3	3	-	-	1	-	-	8
<u>Heavy metals</u>	1	1	-	-	1	-	-	-	-	-	2	-	-	5
<u>Monosodium glutamate</u>	1	-	1	-	-	-	-	1	-	-	-	-	-	3
<u>Mushroom poisoning</u>	-	-	1	-	-	-	2	-	1	-	-	-	-	4
<u>Paralytic shellfish</u>	-	-	-	-	-	-	-	1	-	-	-	-	-	1
<u>Scombrototoxin</u>	1	-	-	-	-	1	1	4	6	4	-	1	-	18
<u>Other</u>	-	-	2	2	1	-	-	-	1	1	1	-	-	8
<u>Total</u>	<u>4</u>	<u>1</u>	<u>4</u>	<u>2</u>	<u>2</u>	<u>1</u>	<u>6</u>	<u>9</u>	<u>8</u>	<u>5</u>	<u>4</u>	<u>1</u>	<u>-</u>	<u>47</u>
<u>PARASITIC</u>														
<u>Trichinella spiralis</u>	-	-	-	-	1	-	-	-	-	-	-	-	-	1
<u>Total</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>	<u>-</u>	<u>1</u>							
<u>VIRAL</u>														
<u>Hepatitis A</u>	3	-	-	-	2	3	3	4	1	1	-	2	-	19
<u>Norwalk Virus</u>	-	-	-	-	-	-	-	1	-	-	1	-	-	2
<u>Total</u>	<u>3</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>2</u>	<u>3</u>	<u>3</u>	<u>5</u>	<u>1</u>	<u>1</u>	<u>1</u>	<u>2</u>	<u>-</u>	<u>21</u>
<u>CONFIRMED TOTAL</u>	<u>13</u>	<u>10</u>	<u>8</u>	<u>11</u>	<u>19</u>	<u>19</u>	<u>33</u>	<u>38</u>	<u>20</u>	<u>17</u>	<u>15</u>	<u>17</u>	<u>-</u>	<u>220</u>
<u>UNKNOWN</u>	<u>27</u>	<u>31</u>	<u>27</u>	<u>29</u>	<u>43</u>	<u>31</u>	<u>21</u>	<u>32</u>	<u>30</u>	<u>27</u>	<u>26</u>	<u>112</u>	<u>-</u>	<u>436</u>
<u>TOTAL 1982</u>	<u>40</u>	<u>41</u>	<u>35</u>	<u>40</u>	<u>62</u>	<u>50</u>	<u>54</u>	<u>70</u>	<u>50</u>	<u>44</u>	<u>41</u>	<u>129</u>	<u>-</u>	<u>656</u>

Table 6 Foodborne Disease Outbreaks by Etiologic Agent and Contributing Factors, United States, 1982

<u>Etiologic Agent</u>	<u>Number of Reported Outbreaks</u>	<u>Number of Outbreaks In Which Factors Reported</u>	<u>Improper Holding Temperatures</u>	<u>Inadequate Cooking</u>	<u>Contaminated Equipment</u>	<u>Food From Unsafe Source</u>	<u>Poor Personal Hygiene</u>	<u>Other</u>
<u>BACTERIAL</u>								
<u>B. cereus</u>	8	5	5	-	-	-	1	-
<u>Brucella</u>	1	-	-	-	-	-	-	-
<u>C. jejuni</u>	2	2	-	-	-	1	-	1
<u>C. botulinum</u>	21	4	1	-	-	-	-	3
<u>C. perfringens</u>	22	20	19	8	3	-	-	2
<u>E. coli</u>	2	1	1	-	-	-	-	-
<u>Salmonella</u>	55	34	16	18	6	6	7	4
<u>Shigella</u>	4	4	1	-	-	-	4	-
<u>S. aureus</u>	28	21	20	3	4	1	9	1
<u>Streptococcus Group A</u>	1	1	1	-	-	-	1	-
<u>Streptococcus Group D</u>	1	-	-	-	-	-	-	-
<u>V. cholerae 01</u>	1	-	-	-	-	-	-	-
<u>V. cholerae non-01</u>	1	1	-	1	-	-	-	-
<u>V. parahaemolyticus</u>	3	2	-	-	-	2	-	-
<u>Y. enterocolitica</u>	2	-	-	-	-	-	-	-
Total	151	95	64	30	13	10	22	11
<u>CHEMICAL</u>								
Ciguatoxin	8	1	-	-	-	1	-	-
Heavy metals	5	4	-	-	2	-	1	2
Monosodium glutamate	3	2	-	-	-	-	-	2
Mushroom poisoning	4	2	-	-	-	2	-	-
Paralytic shellfish	1	1	-	-	-	1	-	-
Scombrototoxin	18	9	5	-	-	5	-	-
Other	8	6	-	-	3	-	-	3
Total	47	25	5	-	5	9	1	7
<u>PARASITIC</u>								
<u>Trichinella spiralis</u>	1	1	-	1	-	1	-	-
Total	1	1	-	1	-	1	-	-
<u>VIRAL</u>								
Hepatitis A	19	14	2	2	-	6	9	3
Norwalk Virus	2	2	-	-	-	-	2	-
Total	21	16	2	2	-	6	11	3
CONFIRMED TOTAL	220	137	71	33	18	26	34	21
UNKNOWN	436	225	87	73	35	94	59	16
TOTAL 1982	656	362	158	106	53	120	93	37

F. Guidelines for Confirmation of Foodborne Disease Outbreak

<u>BACTERIAL</u>	<u>Clinical Syndrome</u>	<u>Laboratory, clinical, and/or epidemiologic criteria for confirmation</u>
1. <u>Bacillus cereus</u>	<p>Vomiting toxin:                      a) incubation period 1-6 hrs.                      b) vomiting, some cases with diarrhea</p> <p>Diarrheal toxin:                      a) incubation period 6-24 hrs.                      b) diarrhea, abdominal cramps, some cases with vomiting</p>	<p>a) isolation of <math>\geq 10^5</math> organisms per gram in epidemiologically incriminated food</p> <p style="text-align: center;">OR</p> <p>b) isolation of organism from stools of ill persons and not in stools of controls</p>
2. <u>Brucella</u>	<p>a) incubation period several days to several months</p> <p>b) clinical syndrome compatible with brucellosis</p>	<p>a) 4-fold increase in titer</p> <p style="text-align: center;">OR</p> <p>b) positive blood culture</p>
3. <u>Campylobacter jejuni</u>	<p>a) incubation period 2-10 days, usually 4-7</p> <p>b) gastrointestinal syndrome-- abdominal pain, often severe; bloody diarrhea common</p>	<p>Isolation of organisms from stool/ blood of ill individuals</p>
4. <u>Clostridium botulinum</u>	<p>a) incubation 2 hours-8 days, usually 12-48 hours</p> <p>b) clinical syndrome compatible with botulism (see CDC Botulism Manual)</p>	<p>a) detection of botulinum toxin in human sera, feces, or food</p> <p style="text-align: center;">OR</p> <p>b) isolation of <u>C. botulinum</u> organism from stools</p> <p style="text-align: center;">OR</p> <p>c) clinical syndrome in persons known to have consumed same food as other individuals with laboratory-proven cases</p>
5. <u>Clostridium perfringens</u>	<p>a) incubation period 9-15 hrs.</p> <p>b) lower intestinal syndrome-- majority of cases with diarrhea but little vomiting or fever</p>	<p>a) organisms of same serotype in epidemiologically incriminated food and stool of ill individuals.</p> <p style="text-align: center;">OR</p> <p>b) isolation of organisms with same serotype in stool of most ill individuals and not in stool of controls</p> <p style="text-align: center;">OR</p> <p>c) <math>\geq 10^5</math> organisms per gram in epidemiologically incriminated food provided specimen properly handled</p>
6. <u>Escherichia coli</u>	<p>a) incubation period 6-36 hrs.</p> <p>b) gastrointestinal syndrome-- majority of cases with diarrhea</p>	<p>a) demonstration of organisms of same serotype in epidemiologically incriminated food and stool of ill individuals and not in stool of controls</p> <p style="text-align: center;">OR</p> <p>b) isolation from stool of most ill individuals, organisms of the same serotype which have been shown to be enterotoxigenic or invasive by laboratory techniques</p>

	<u>Clinical Syndrome</u>	<u>Laboratory, clinical, and/or epidemiologic criteria for confirmation</u>
7. <u>Salmonella</u>	a) incubation period 6-48 hrs.  b) gastrointestinal syndrome-- majority of cases with diarrhea	a) isolation of <u>Salmonella</u> organism from epidemiologically implicated food  OR b) isolation of <u>Salmonella</u> organism from stools of ill individuals
8. <u>Shigella</u>	a) incubation period 12-50 hours  b) gastrointestinal syndrome-- majority of cases with diarrhea	a) isolation of <u>Shigella</u> organism from epidemiologically implicated food  OR b) isolation of <u>Shigella</u> organism from stools of ill individuals
9. <u>Staphylococcus aureus</u>	a) incubation period 30 min.- 8 hours (usually 2-4 hrs.)  b) gastrointestinal syndrome-- majority of cases with vomiting	a) detection of enterotoxin in epidemiologically implicated food  OR b) organisms with same phage type in stools or vomitus of ill individ- uals; isolation from epidemiologi- cally implicated food and/or skin or nose of food handler is support- ive evidence  OR c) isolation of $\geq 10^5$ organisms per gram in epidemiologically implicated food
10. <u>Streptococcus Group A</u>	a) incubation period 1-4 days  b) febrile URI syndrome	a) isolation of organisms with same M and T type from implicated food  OR b) isolation of organisms with same M and T type from throats of ill individuals
11. <u>Vibrio cholerae</u> 01	a) incubation period 1-5 days  b) gastrointestinal syndrome-- majority of cases with diarrhea and without fever	a) isolation of toxigenic <u>V. cholerae</u> 01 from epidemiologically incriminated food  OR b) isolation of organisms from stools or vomitus of ill individuals  OR c) significant rise in vibriocidal, bacterial agglutinating or anti- toxin antibodies in acute and early convalescent sera, or significant fall in vibriocidal antibodies in early and late convalescent sera in persons not recently immunized
<u>Vibrio cholerae</u> Non-01	a) incubation period up to 3 days  b) gastrointestinal syndrome-- majority of cases with diarrhea	a) isolation of non-01 <u>V. cholerae</u> of same serotype from stools of ill persons; isolation from epidemio- logically implicated food is sup- portive evidence
12. <u>Vibrio parahaemolyticus</u>	a) incubation period 4-30 hrs.  b) gastrointestinal syndrome-- majority of cases with diarrhea	a) isolation of $\geq 10^5$ organisms from epidemiologically implicated food (usually seafood)  OR b) isolation of Kanagawa-positive organisms from stool of ill individuals

	<u>Clinical Syndrome</u>	<u>Laboratory, clinical, and/or epidemiologic criteria for confirmation</u>
13. Others	clinical data appraised in individual circumstances	laboratory data appraised in individual circumstances
<b><u>CHEMICAL</u></b>		
1. Heavy metals	a) incubation period 5 min. to 8 hrs. (usually less than 1 hr)	demonstration of high concentration of metallic ion in epidemiologically incriminated food or beverage
Antimony Cadmium Copper Iron Tin	b) clinical syndrome compatible with heavy metal poisoning--usually gastrointestinal syndrome and often metallic taste	
<b>2. Ichthyosarcotoxin</b>		
Ciguatoxin	a) incubation period 1-48 hrs. (usually 2-8 hrs.)	a) demonstration of ciguatoxin in epidemiologically incriminated fish
	b) usually gastrointestinal symptoms followed by neurologic manifestations, including paresthesia of lips, tongue, throat or extremities, and reversal of hot and cold sensation	<u>OR</u> b) clinical syndrome in person(s) who have eaten a type of fish previously associated with ciguatera fish poisoning (e.g., snapper, grouper)
Puffer fish (tetrodotoxin)	a) incubation period 10 min. to 3 hrs. (usually 10-45 min.)	a) demonstration of tetrodotoxin in fish
	b) paresthesia of lips, tongue, face or extremities often followed by numbness, loss of proprioception or a "floating" sensation	<u>OR</u> b) puffer fish epidemiologically incriminated
Scombrototoxin	a) incubation period 1 min. to 3 hours (usually less than 1 hour)	a) demonstration of elevated histamine levels in epidemiologically incriminated fish
	b) flushing, headache, dizziness, burning of mouth and throat, upper and lower gastrointestinal symptoms, urticaria and generalized pruritus	<u>OR</u> b) clinical syndrome in person(s) known to have eaten a fish of order Scombroidei or type of fish previously associated with scombroid poisoning (e.g., mahi-mahi)
3. Monosodium glutamate	a) incubation period 3 min. to 2 hours (usually less than 1 hour)	history of large amounts (usually >1.5 grams) of MSG having been added to epidemiologically incriminated food
	b) burning sensations in chest, neck, abdomen or extremities, sensations of lightness and pressure over face, or a heavy feeling in the chest	
4. Mushroom poison Group containing ibotenic acid and muscimol	a) incubation period 1-12 hrs. (usually less than 4 hrs.)	a) demonstration of toxic chemical in epidemiologically incriminated mushrooms
	b) clinical syndrome compatible w/mushroom poisoning by this group--often including confusion, delirium, visual disturbances	<u>OR</u> b) epidemiologically incriminated mushrooms identified as a toxic type

	<u>Clinical Syndrome</u>	<u>Laboratory, clinical, and/or epidemiologic criteria for confirmation</u>
Group containing amanitotoxins and phallotoxins, or gyromitrin	a) incubation period 5-18 hrs. b) characteristic clinical syndrome compatible with mushroom poisoning by this group--upper and lower gastrointestinal symptoms followed by hepatic and/or renal failure	a) demonstration of toxic chemical in epidemiologically incriminated mushrooms <u>OR</u> b) epidemiologically incriminated mushrooms identified as a toxic type
Groups containing muscarine, psilocybin and psilocin, gastrointestinal irritants, disulfiram-like compounds	a) characteristic incubation period b) clinical syndrome compatible with mushroom poisoning by these groups	a) demonstration of toxic chemical in epidemiologically incriminated mushrooms <u>OR</u> b) epidemiologically incriminated mushroom identified as toxic type
5. Paralytic or neurotoxic shellfish poison	a) incubation period 30 min. to 3 hours b) paresthesias of lips, mouth or face, and extremities; weakness, including respiratory difficulty in most severe cases; upper and lower gastrointestinal symptoms in some cases	a) detection of toxin in epidemiologically incriminated mollusks <u>OR</u> b) detection of large numbers of shellfish poisoning-associated species of dinoflagellates in water from which epidemiologically incriminated mollusks gathered
6. Other chemical	clinical data appraised in individual circumstances	laboratory data appraised in individual circumstances
<u>PARASITIC AND VIRAL</u>		
1. <u>Trichinella spiralis</u>	a) incubation period 3-30 days b) clinical syndrome compatible with trichinosis--often including fever, high eosinophil count, orbital edema, myalgia	a) muscle biopsy from ill individual <u>OR</u> b) serological tests <u>OR</u> c) demonstration of larvae in incriminated food
2. Hepatitis A	a) incubation period 10-45 days b) clinical syndrome compatible with hepatitis--usually including jaundice, GI symptoms, dark urine	liver function tests compatible with hepatitis in affected persons who consumed the epidemiologically incriminated food
3. Others	clinical evidence appraised in individual circumstances	laboratory evidence appraised in individual circumstances





H. LINE LISTING OF FOODBORNE DISEASE OUTBREAKS, 1982

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled and Eaten
				Patient	Vehicle	Food-Handler		
<b><u>BACTERIAL</u></b>								
<b><u>Bacillus cereus</u></b>								
	California	44	7/19		+		H/M ice cream	Camp
	Illinois	9	8/23		+		Fried rice	Restaurant
	Kentucky	7	7/21		+		Reuben sandwich	Hospital Cafeteria
	New Mexico	22	1/18		+		Beef taco	School
	New Mexico	5	8/04	+	+		Fried rice	Restaurant
	New York	100	1/09		+		Steamed rice	Fire Hall
	New York City	6	9/17		+		Baked food	School
	Wyoming	7	9/14		+		Fried rice	Restaurant
<b><u>Brucella</u></b>								
	Connecticut	3	3/	+			Unknown	Private home
<b><u>Campylobacter jejuni</u></b>								
	Vermont	15	9/22	+			Raw milk	Variety places
	Wisconsin	16	6/27	+			Raw milk	Private home
<b><u>Clostridium botulinum</u></b>								
	Alaska	2	8/		+		Fermented fish eggs	Private home
	California	1	2/		+		H/C beans	Private home
	California	2	2/	+	+		C/C mushrooms	Private home
	California	2	6/	+	+		H/C peppers	Private home
	California	1	7/	+	+		C/C tomatoes	Private home
	California	1	8/03	+	+		Beef pot pie	Private home
	California	4	11/	+	+		H/C swiss chard	Private home
	California	2	12/	+	+		H/C bean salad	Private home
	Colorado	1	7/	+	+		H/C eggplant	Private home
	Massachusetts	1	11/		+		H/C relish	Private home
	North Dakota	1	9/05	+	+		H/C beans	Private home
	Oregon	3	5/		+		H/C asparagus	Private home
	Utah	1	6/	+			Unknown	Private home
	Utah	1	12/	+			H/C peppers	Private home
	Washington	1	5/09		+		H/C salmon	Private home
	Washington	1	6/07		+		H/C asparagus	Private home
	Washington	1	7/11	+	+		H/C green beans	Private home
	Washington	1	12/16		+		H/C asparagus	Private home
	West Virginia	1	2/		+		H/C green beans	Private home
	Wisconsin	1	10/	+			H/C zucchini	Private home
	Wyoming	1	11/29		+		H/C green beans	Private home
<b><u>Clostridium perfringens</u></b>								
	Arizona	300	5/05		+		Tamales, enchiladas	Prison
	Arizona	108	7/25		+		Barbacoa and beans	Meeting hall
	California	166	11/02	+			Spaghetti, meat sauce	Prison
	Connecticut	83	3/08		+		Sloppy Joe	School
	Illinois	39	12/19	+	+		Turkey & gravy	Party rented
	Michigan	35	9/19	+	+	+	Swedish meatballs	Wedding party
	Minnesota	70	9/26	+	+		Beef Bourguignon	Restaurant

H/M = home-made

H/C = home-canned

C/C = commercial-canned

Etiology	State	Number of Cases	Date of Onset	Lab Data			Location Where Food Mishandled and Eaten
				Patient	Vehicle	Food-Handler	
<u>Clostridium perfringens</u> (Cont'd)							
	Minnesota	32	12/23	+	+		Moose Lodge hall
	New Jersey	17	6/28		+		Meatloaf Hospital
	New Mexico	14	5/13		+		Enchiladas Cafeteria
	New York	10	6/29	+			Unknown Restaurant
	Pennsylvania	12	5/09	+			Turkey Restaurant
	Pennsylvania	25	5/22	+			Unknown Banquet
	Pennsylvania	25	8/14	+	+		Roast beef Banquet
	Tennessee	34	12/23	+	+		Turkey Christmas party
	Washington	5	2/23	+			Beef enchiladas Restaurant
	Washington	73	2/26	+			Unknown Prison
	Washington	2	4/26	+			Unknown Restaurant
	Washington	5	5/09	+			Unknown Restaurant
	Washington	3	8/04	+			Unknown Restaurant
	Washington	6	10/16	+			Unknown Restaurant
	Washington	125	11/06	+			Turkey salad Reformatory spread
<u>Escherichia coli</u>							
	Michigan	21	5/28	+	+		Ground beef Restaurant
	Oregon	26	2/05	+			Ground beef Restaurant
<u>Salmonella</u>							
<u>S. typhimurium, infantis</u>	Arizona	19	12/05	+	+		Raw milk Private home
<u>S. montevidео</u>	California	63	7/18	+		+	Unknown Restaurant
Group B	California	120	10/23	+			Chicken salad Picnic
<u>S. enteritidis*</u>	Connecticut	15	4/12	+		+	Unknown Hospital
<u>S. typhimurium</u>	Connecticut	3	6/19	+			Unknown Restaurant
<u>S. heidelberg</u>	Connecticut	15	8/08	+		+	Macaroni/cheese Hosp cafeteria
<u>S. enteritidis</u>	Connecticut	7	8/09	+		+	Unknown Restaurant
<u>S. typhimurium, copenhagen</u>	D. C.	55	2/18	+	+	+	Prime rib School
<u>S. thompson</u>	Florida	36	4/06	+		+	Unknown Restaurant
<u>S. montevidео</u>	Georgia	13	7/11	+	+		H/M chocolate Church
<u>S. heidelberg</u>	Hawaii	4	8/07	+			Unknown Restaurant
<u>S. typhimurium</u>	Illinois	7	6/19	+		+	Refried beans Restaurant
<u>S. anatum, infantis</u>	Illinois	9	8/08	+	+		H/M ice cream Private home
<u>S. montevidео</u>	Illinois	5	9/09	+		+	Salad bar Restaurant
<u>S. typhimurium, hadar</u>	Kansas	30	5/30	+			Unknown Picnic
<u>S. unspecified</u>	Kentucky	15	10/23	+			Unknown Restaurant
<u>S. enteritidis</u>	Maine	160	11/25	+	+	+	Turkey and giblets Restaurant
<u>S. meleagridis</u>	Massachusetts	104	3/	+			Unknown Statewide
<u>S. typhimurium</u>	Massachusetts	10	5/	+		+	Unknown Nursing home
<u>S. enteritidis</u>	Massachusetts	72	7/	+			Unknown School
<u>S. enteritidis</u>	Massachusetts	428	7/	+		+	Unknown Statewide
<u>S. thompson</u>	Massachusetts	3	8/	+			Unknown Buffet lunch
<u>S. enteritidis</u>	Massachusetts	48	8/	+			Unknown School
<u>S. heidelberg</u>	Massachusetts	30	8/31	+		+	Unknown Training camp
<u>S. enteritidis</u>	Massachusetts	33	9/15	+		+	Unknown Hospital
<u>S. hadar, agona</u>	Massachusetts	24	11/25	+		+	Turkey Restaurant
<u>S. enteritidis</u>	Massachusetts	34	12/09	+	+	+	Cornish hens School
Group D	Massachusetts	10	12/17	+			Unknown Hospital
<u>S. unspecified</u>	Minnesota	47	8/22	+			Pistachio nut salad Church
<u>S. enteritidis</u>	New Hampshire	5	7/	+			Unknown Restaurant
<u>S. enteritidis</u>	New Hampshire	26	7/24	+		+	Scrambled eggs Nursing home
<u>S. unspecified</u>	New Jersey	5	4/30	+			Unknown Restaurant
<u>S. enteritidis</u>	New Jersey	47	6/05	+		+	Seafood combo Restaurant
Groups B, C	New York	16	5/08	+			Unknown School
Group B	New York	8	7/10	+		+	Tuna macaroni salad Restaurant
Group B	New York	31	7/11	+			Chicken Private home

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled and Eaten
				Patient	Vehicle	Food-Handler		
<u>Salmonella (Cont'd)</u>								
<u>S. enteritidis</u>	New York	114	7/19	+	+		Chicken/turkey salad	Camp
Group B	New York	2	7/24	+			Unknown	Private home
Group B	New York	21	8/01	+			Unknown	Private camp
<u>S. enteritidis</u> CZ	New York	3	12/26	+			Unknown	Private home
<u>S. uganda</u>	New York City	2	1/23		+		Pork	Restaurant
<u>S. chester</u>	North Dakota	18	1/31	+	+		Unknown	Picnic
<u>S. typhimurium</u> , <u>heidelberg</u>	North Dakota	24	4/22	+			Unknown	Nursing home
<u>S. thompson</u>	Oklahoma	16	10/02	+	+		Strawberry ice cream	Private home
<u>S. poona</u>	Oregon	38	6/01	+		+	Unknown	Church
<u>S. typhimurium</u>	Pennsylvania	156	4/24	+		+	Turkey	School
<u>S. heidelberg</u>	Pennsylvania	15	8/30	+			Lasagna	Restaurant
<u>S. Arizona hinshawii</u>	Pennsylvania	21	11/03	+	+	+	Turkey	Church
<u>S. typhimurium</u>	Texas	16	8/10	+	+		H/M ice cream	Private home
<u>S. typhimurium</u>	Washington	13	5/30	+			Unknown	Private home
<u>S. typhimurium</u>	Washington	2	9/21	+			Unknown	Private home
<u>S. enteritidis</u>	Washington	11	10/25	+	+		Pork	Private home
<u>S. typhimurium</u>	Washington	17	10/30	+		+	Unknown	Restaurant
<u>S. typhimurium</u>	Washington	2	12/28	+	+		H/M sausage	Private home
<u>S. typhimurium</u>	Wyoming	8	6/27	+	+		Eggs in h/m ice cream	Private home
<u>Shigella</u>								
<u>S. flexneri</u> 31	California	7	4/21	+		+	Unknown	Church
<u>S. sonnei</u>	Colorado	75	10/14	+		+	Salad bar	Restaurant
<u>S. sonnei</u>	Hawaii	11	11/28	+			Unknown	Private home
<u>S. sonnei</u>	Pennsylvania	23	9/27	+		+	Tuna salad	Restaurant
<u>Staphylococcus aureus</u>								
	California	26	1/01	+	+		Beef tacos	Church
	Connecticut	70	10/29		+		Turkey roast	School
	Delaware	42	9/24		+		Chicken	Retirement home
	Hawaii	24	10/10		+		Coconut-cream cake	Hospital
	Hawaii	2	10/22		+		Chicken	Private home
	Illinois	108	4/22		+	+	Turkey	School
	Illinois	37	8/15	+		+	Potato salad	Nursing home
	Iowa	40	8/01	+	+		Bologna sandwich	Bus
	Kansas	6	5/27		+		Ham	Restaurant
	Massachusetts	8	4/	+	+		Turkey	State Hospital
	Massachusetts	19	8/	+			Unknown	Camp
	Minnesota	5	5/04		+		Chicken pot pie	Restaurant
	New Mexico	4	7/22		+		Beef jerky	Private home
	New York	8	6/20	+	+		Cream puffs	Private home
	New York	2	7/23		+		Pastrami	Private home
	Pennsylvania	52	6/06	+	+	+	Ham	Street fair
	Pennsylvania	14	7/03	+	+	+	Ground beef taco mix	Restaurant
	Pennsylvania	14	7/21	+	+		Ham/cheese sandwich	Chartered bus
	Pennsylvania	33	7/22	+	+		Stuffed chicken breast	Picnic
	Pennsylvania	51	8/04	+	+		Pork	Boat trip
	Pennsylvania	29	8/07		+	+	Potato salad	Private club
	Pennsylvania	13	11/21	+	+		Roast beef	Firehouse
	Pennsylvania	15	12/01	+	+	+	Chicken salad	Church
	Tennessee	4	2/05	+	+		Hot dogs	Restaurant
	Tennessee	6	3/04		+		BRO pork	Private home
	Texas	25	6/21		+		Unknown	Picnic
	Vermont	6	12/13		+	+	Refried beans	Restaurant
	Washington	6	1/20	+		+	Roast beef sandwich	Restaurant

<u>Etiology</u>	<u>State</u>	<u>Number of Cases</u>	<u>Date of Onset</u>	<u>Lab Data</u>			<u>Vehicle</u>	<u>Location Where Food Mishandled and Eaten</u>
				<u>Patient</u>	<u>Vehicle</u>	<u>Food-Handler</u>		
<u>Streptococcus</u>								
Group A B-Hemolytic	New Hampshire	34	7/18	+			Onion and clam dip	Party
<u>Vibrios</u>								
<u>V. cholerae</u> O1	Guam/Trust Ter.	892	8/01	+			Uncooked shell-fish	Private home
<u>V. cholerae</u> non-O1	Guam/Trust Ter.	7	12/08	+			Shrimp	Private home
<u>V. parahaemolyticus</u>	Massachusetts	26	8/	+			Raw clams	Clam bake
<u>V. parahaemolyticus</u>	New York	10	7/06	+			Raw clams	Picnic
<u>V. parahaemolyticus</u>	New York	3	7/16		+		Steamed clams	Private home
<u>Yersinia enterocolitica</u>								
	Pennsylvania	16	2/10	+			Bean sprouts	Brownie meeting
	Multiple	172	6/11	+			Pasteurized milk	Private home
<u>CHEMICAL</u>								
<u>Ciguatera</u>								
	Hawaii	2	7/30				Forktailed snapper	Private home
	Hawaii	8	8/16				Barracuda	Private home
	Hawaii	6	11/12				Royal seabass	Private home
	Virgin Islands	5	7/30				Fish chowder	Restaurant
	Virgin Islands	4	8/01				Other fish	Restaurant
	Virgin Islands	7	8/08				Barracuda	Restaurant
	Puerto Rico	15	3/27				Other fish	Unknown
	Puerto Rico	15	3/28				Other fish	Private home
<u>Heavy metals</u>								
Copper	Illinois	3	1/11		+		Soft drink	Private home
Copper	New York	1	11/09				Water fountain	Restaurant
Copper	Washington	3	5/16				Carbonated drink	Restaurant
Zinc	New York City	1	2/07		+		Ginger ale	Restaurant
Zinc and iron	New Mexico	18	11/19		+		Punch	Class party
<u>Monosodium glutamate</u>								
	Minnesota	2	1/19				Chinese food	Restaurant
	Washington	2	8/21				Chinese food	Restaurant
	New York City	6	3/28		+		Turkey stuffing/gravy	
	Private home							
<u>Mushroom poisoning</u>								
<u>Amanita bruenscens</u>	California	5	3/07	+	+		Mushrooms	Private home
<u>Amanita muscaria</u>	New York	1	9/		+		Wild mushrooms	Private home
<u>Leucosarcus naucin</u>	New York	2	7/		+		Wild mushrooms	Private home
<u>T. platyphylla</u>	New York	1	7/		+		Wild mushrooms	Outside
<u>Paralytic Shellfish</u>								
	Alaska	5	8/10		+		Mussels	Private home

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled and Eaten
				Patient	Vehicle	Food-Handler		
<u>Scambrotoxin</u>								
	Alaska	2	8/03				Raw salmon	Private home
	Alaska	9	8/09				Raw fish	Ship
	California	2	9/26		+		Mahi-Mahi	Private home
	Connecticut	2	9/06		+		Bluefish	Restaurant
	Connecticut	1	9/08		+		Bluefish	Restaurant
	Connecticut	8	9/15		+		Bluefish	Work
	Connecticut	4	10/		+		Bluefish	Restaurant
	Connecticut	5	10/13		+		Bluefish	Cafeteria
	Hawaii	2	6/14				Blue ulua	Private home
	Hawaii	4	12/09				Mahi-Mahi	Restaurant
	New York	2	1/20		+		Mahi-Mahi	Restaurant
	New York	3	7/15				Bluefish	Restaurant
	New York	2	8/04		+		Tuna	Private home
	New York	2	8/06		+		Bluefish	Restaurant
	New York	1	9/13				Tuna sandwich	Private home
	New York	3	10/03				Tuna	Private home
	Washington	4	9/03		+		Mahi-Mahi	Private home
	Washington	2	10/29				Mahi-Mahi	Restaurant

Other chemical

Calcium chloride	Hawaii	2	11/17		+		Frozen snack	Private home
Hot peppers	Minnesota	2	10/21				Fruit or vegetables	Restaurant
Lye	New Jersey	5	5/12		+		Tea	Restaurant
Nitrites	New York City	1	4/01		+		Franks and sauerkraut	Restaurant
Sodium hydroxide	Ohio	4	3/27		+		Soft drink	Private home
Cardol cashew shell	Pennsylvania	54	4/06	+	+		Cashew cardol oil	Private home
Nitrite	Utah	6	9/26		+		Beef jerky	Private home
Trisodium phosphate	Washington	1	3/23		+		Cleaning compound	Restaurant

PARASITIC

Trichinella spiralis

	Hawaii	4	5/28	+			Smoked wild pork	Private home
--	--------	---	------	---	--	--	------------------	--------------

VIRAL

Hepatitis A

California		6	10/28	+		+	Unknown	Hosp. cafeteria
Connecticut		30	8/08	+			Unknown	Camp
Florida		9	1/27	+		+	Unknown	Restaurant
Florida		11	12/25	+			Oysters	Store
Hawaii		6	12/01	+			Unknown	Private home
Iowa		16	7/18	+		+	Unknown	Army Reserve
Massachusetts		6	1/01	+		+	Unknown	Corporate party
Massachusetts		7	1/25	+		+	Unknown	Cafeteria
Michigan		10	7/12	+			Peanut butter cake	Natl Guard Camp
Michigan		3	8/13	+			Ice cream	Beach
Minnesota		24	5/10	+		+	Salad and ice	Restaurant
Minnesota		24	6/10	+			Unknown	Restaurant
New York		18	5/30				Raw & steamed clams	Private home
New York		5	6/01	+			Raw clams	Private home
New York		46	6/06	+			Clams	Picnic
New York		7	7/05	+			Raw clams	Private home
New York		2	8/01	+			Unknown	Private home
New York		10	9/01	+			Unknown	Restaurant
New York City		85	8/04	+		+	Tuna fish salad	Restaurant

<u>Etiology</u>	<u>State</u>	<u>Number of Cases</u>	<u>Date of Onset</u>	<u>Lab Data</u>			<u>Location Where Food Mishandled and Eaten</u>
				<u>Patient</u>	<u>Vehicle</u>	<u>Food-Handler</u>	
<u>Norwalk Virus</u>							
	Minnesota	3000	8/21	+		+	Cake/frosting
	Minnesota	2000	11/11	+			Cole slaw Variety places Restaurant

UNKNOWN

A line listing of outbreaks of unknown etiology may be obtained by writing to the Enteric Diseases Branch, Division of Bacterial Diseases, Center for Infectious Diseases, Centers for Disease Control, Atlanta, Georgia 30333.

I. Selected Foodborne Outbreak Articles Taken From Morbidity and Mortality Weekly Report, 1982

Botulism and Commercial Pot Pie--California  
MMWR 1983;32:39-45

On August 3, 1982, a 56-year-old woman residing in Los Angeles County, California, developed diplopia, weakness, difficulty breathing, and chest pain. She had respiratory arrest on admission to the hospital but was intubated, resuscitated, and placed in intensive care. Examination showed complete bilateral ptosis, ophthalmoplegia, facial muscle weakness, and areflexia. Cerebrospinal fluid was normal except for increased glucose; Tensilon test was negative. She had a history of seizure disorder, diabetes mellitus, and organic brain syndrome. An infectious disease consultant thought her subsequent fever was due to pneumonia secondary to aspiration, and he suspected botulism as the underlying cause of her illness.

The patient lives with her husband and grown son who both prepare meals for her and attempt a strict diet in consideration of her diabetes. When asked about the patient's food history before onset of illness, the husband and son named no likely suspects for botulism. No home-preserved foods had been served, and, with one exception, she had not eaten other foods that were not freshly prepared for her or were not also consumed by her husband and son. The exception was commercial beef pot pie, which was accidentally mishandled, then consumed by the patient 1 day before illness began.

The son had prepared the pot pie for an earlier evening meal. The frozen pie was baked in an oven for 40-45 minutes. As he was about to serve it to his mother, his father came home with some freshly cooked hamburgers just purchased at a take-out restaurant. The pot pie was put aside on an unrefrigerated shelf. Two and one-half days later, the son came home and found his mother had just consumed this pot pie without reheating it.

An uneaten portion of the pot pie, still in its metal plate, was retrieved by the family members. Type A botulism toxin was found in this pie by a mouse-inoculation test performed at a U.S. Department of Agriculture laboratory in Beltsville, Maryland, and type A toxin was also demonstrated in the patient's serum by the state's Microbial Disease Laboratory.

Editorial Note: This is the third case of botulism associated with commercial pot pies reported from California (1,2); one other episode (involving two clinically diagnosed patients) was reported from Minnesota in 1960 (3). Mishandling of the pot pies occurred in three of these episodes, and mishandling was also suspected in the fourth. The known mishandlings consisted of leaving the baked pot pie in the oven with the pilot light on, thereby maintaining "incubator" temperatures overnight. The pies were then eaten with no (or insufficient) reheating to destroy toxin. Or, as in the present case, the baked pie sat out at room temperature for over 2 days during hot weather--conditions that also could simulate an incubator.

In these situations, it is suspected that the original baking killed competing organisms in the pies and eliminated much of the oxygen. The heat-resistant, anaerobic Clostridium botulinum, which was evidently present and can be found in many fresh, frozen, and other food products, was then presumably able to germinate and produce toxin under the crust during storage at warm, incubator-like temperatures. Products such as pot pies should be kept frozen before heating and ideally should be served hot after the first cooking. If any such products are to be saved, they should be quickly refrigerated, then reheated to hot temperatures. This would minimize any risk of botulinal poisoning.

## References

1. State of California, Department of Health Services. Botulism--home-canned figs and chicken pot pie. California Morbidity 1975, No. 46.
2. State of California, Department of Health Services. Type A botulism associated with commercial pot pie. California Morbidity 1976, No. 51.
3. CDC. Botulism. MMWR 1960;9(27):2

### Ciguatera Fish Poisoning--Bahamas, Miami MMWR 1982;28:391-2

On March 6, 1982, the U.S. Coast Guard in Miami, Florida, received a request for medical assistance from an Italian freighter located in waters off Freeport, Bahamas. Numerous crew members were ill with nausea, vomiting, and muscle weakness and required medical evacuation for hospitalization and treatment.

A total of 14 ill crew members were airlifted to three Florida hospitals. Three were seen in emergency rooms and later released. Eleven were hospitalized; seven required admission to intensive care units. All patients were Italian males, age 24-40 years; symptoms included diarrhea--12 patients (86%), vomiting--11 (79%), paresthesias--11 (79%), hypotension--10 (71%), peripheral muscular weakness--9 (65%), nausea--8 (57%), abdominal cramping--6 (43%), pruritis--4 (29%), and peripheral numbness--2 (14%). These findings were consistent with ciguatera fish poisoning, and an epidemiologic investigation was initiated.

The ship employed 26 crew members and is permanently based near Freeport, where it ferries petroleum products ashore from large tankers. On March 4, a crew member caught a 25-pound barracuda while fishing from the ship. On March 6, 14 crew members cooked and ate the barracuda; all became ill within 6 hours. None of the 12 crew members who did not eat the barracuda became ill. Six of the ill crew members reported becoming sick 45 minutes to 6 hours after the implicated meal (median: 2.5 hours). All 14 crew members eventually recovered without sequelae and returned to work. Median length of hospital stay was 6 days.

**Editorial Note:** Ciguatera is a human intoxication syndrome associated with the consumption of marine tropical reef fishes. Although recent surveys indicate that poisonings are relatively uncommon in Florida (1,2), one investigator recorded 280 intoxications from January 1978 to June 1980 (2).

The ichthyosarcotoxins are thought to be accumulated through the food chain, the toxins being produced by microalgae known as dinoflagellates (3,4). The toxins are lipid-soluble and appear to accumulate in the flesh, fatty tissue, and viscera of large predatory species of fish, such as barracuda, grouper, and snapper (5,6). The isolation, purification, and characterization of the suspected toxins have been hampered by limited availability of authentic ciguatoxic fish, lack of a specific sensitive assay, and the low concentration and heterogeneity of toxins present in specimens.

The assessment of toxicity most often used is the mouse bio-assay. Based on signs elicited following intraperitoneal (IP) injection, it includes, but is not limited to inactivity, diarrhea, labored breathing, cyanosis, piloerection, tremors, paralysis, and staggering gait. Death occurs when the injection is given in higher doses, with a lethal dose, 50% kill (LD<sub>50</sub>), of 0.45 ug/kg for purified toxin (5). Thus, ciguatoxin is one of the most potent marine toxins known. The barracuda's head was toxic by mouse bio-assay with an LD<sub>50</sub> (IP) of 2-5 gram equivalents of original fish meat. Thin-layer chromatographic separation of extracts revealed the presence of at least two major toxins. Further purification is under way to define more clearly the toxin(s) implicated in this outbreak.

## References

1. Lawrence DN, Enriquez MB, Lumish RM, Maceo A. Ciguatera fish poisoning in Miami. JAMA 1980;244:254-8.
2. Poli MA. A review of ciguatera, with special reference to the Caribbean, and an investigation into its significance and incidence in Florida. Master's thesis, University of Miami, 1982.
3. CDC. Ciguatera fish poisoning--St. Croix, Virgin Islands of the United States. MMWR 1981;30:138-9.
4. Bagnis R, Chanteau S, Chunque E, Hurtel JM, Yasumoto T, Inoue A. Origins of ciguatera fish poisoning: a new dinoflagellate, Gambierdiscus toxicus Adachi and Fukuyo, definitely involved as a causal agent. Toxicol 1980;18:199-208.
5. Withers NW. Ciguatera fish poisoning. Ann Rev Med 1982;3:97-111.
6. Banner AH. Ciguatera: a disease from coral reef fish. In: Jones OA, Endean R, eds. Biology and geology of coral reefs. New York: Academic Press, 1975;111:177-213.

### Enteric Illness Associated with Raw Clam Consumption--New York MMWR 1982;31:449-51

Since June 1, 1982, the New York State Health Department has received reports of at least 14 separate outbreaks of gastroenteritis associated with consumption of raw clams. Approximately 150 persons have been affected. Typical symptoms have included diarrhea and abdominal cramps beginning 12-72 hours after eating clams, with nausea, vomiting, and fever occurring less often. In three of these outbreaks, seven individuals subsequently developed hepatitis A 21-37 days after eating clams. Three other persons developed hepatitis A without initial gastrointestinal symptoms. Eight of the 10 cases were verified by the presence of IgM antibody to hepatitis A virus (HAV); results on the others are pending.

A summary of four of these outbreaks follows:

Outbreak A: On May 29, 24 individuals attended a private party in Albany County at which raw clams were served. Within 6-24 hours, 18 (90%) of 20 persons who had eaten clams developed diarrhea and abdominal cramps, which lasted 1-3 days. None of four persons who remained well had consumed clams. Stool specimens obtained shortly after onset of illness from seven persons with gastroenteritis were negative for Salmonella, Shigella, and Campylobacter. Two persons who ate clams from the same lot as those consumed at the party were evaluated at the New York State Health Department's laboratory. Although cultures of extracts from these clams did not grow enteric bacterial pathogens, both 27-nm and 40-nm virus-like particles were observed by electron microscopy.

Outbreak B: On May 30, fourteen people attended a private party in Rensselaer County at which clams were served. Five (83%) of six persons who ate raw clams developed diarrhea, nausea, vomiting, and abdominal cramps 36-72 hours later; symptoms persisted for 1-2 days. None of the eight persons who did not eat raw clams became ill. One of the five individuals with gastroenteritis, who worked as a food handler, developed hepatitis A (confirmed by the presence of HAV-specific IgM antibody) 34 days after eating clams, prompting county health officers to administer immunoglobulin (IG) as a preventive measure to 850 people exposed to foods he had prepared.

Outbreak C: On June 5, members of multiple bowling leagues attended a picnic in Albany County. Many of the approximately 200 attendees developed diarrhea, nausea, vomiting, and abdominal cramps 12-72 hours after the event. Forty-five of 126 persons interviewed reported gastroenteritis; 42 (89%) of these had eaten raw clams. Only raw clams were significantly associated with illness ( $p < 0.001$ ). Four persons who consumed clams and were affected by gastroenteritis developed hepatitis A 29-37

days later. This outbreak was not recognized in time to obtain specimens from persons with acute gastrointestinal illness.

Outbreak D: On July 11, 11 persons attended a party in Schenectady County at which raw clams were served. All seven individuals who ate clams developed diarrhea and abdominal cramps 15-60 hours later; none had fever or vomiting. Diarrhea persisted for up to 1 week in several persons. None of four persons who did not eat clams became ill. Thus, clams were epidemiologically implicated as the vehicle of transmission. Stool samples from five ill individuals were negative for enteric bacterial pathogens (Salmonella, Shigella, Vibrio, Campylobacter, and Yersinia). Examination of stools for virus is pending.

Inadequate or absent tagging of the clams implicated in these outbreaks has made it difficult to accurately determine the clams' source. However, current information indicates clams responsible for the outbreaks originated in coastal waters from at least three states: Massachusetts, New York, and Rhode Island. The timing of these outbreaks may be related to contamination of harvesting beds by the heavy rains and subsequent runoff that occurred in the Northeast during May and early June. Preliminary data from New York and Rhode Island indicate an increase in coliform counts in clam-harvesting waters monitored during this time.

Since December 1981, the New York State Department of Health has been informed of 33 outbreaks of clam-related illness involving more than 250 cases of gastroenteritis and 20 cases of hepatitis A. One county where clams are harvested has noted a two-fold increase in reported cases of hepatitis A. One county where clams are harvested has noted a two-fold increase in reported cases of hepatitis A for the first 6 months of this year compared with the same period last year (60 in 1982 vs 31 in 1981); 45% of the 1982 patients had histories of clam consumption consistent with the incubation period of hepatitis A. An intensive evaluation of 1,559 food establishments, conducted between July 22 and July 29, revealed that 125 (14%) of 908 that stock shellfish sold clams that were untagged or improperly certified (to identify their waters of origin).

Because these outbreaks suggested a recent problem of clam contamination, New York State Health Department officials currently advise individuals to refrain from eating raw clams. In addition, they advise giving IG to persons involved in clam-associated outbreaks of gastroenteritis, provided it can be administered within 2 weeks of clam consumption.

Editorial Note: Ingestion of shellfish has been known for over 50 years to cause outbreaks of bacterial and viral enteric diseases (1). Typhoid fever (2), hepatitis A (3,4), cholera (5), and Vibrio parahaemolytic (6) have long been associated with ingestion of raw clams and oysters. More recently, raw shellfish contaminated with non-01 V. cholerae (7) and Norwalk virus (8,9) have also been reported as causes of gastroenteritis outbreaks. Although gastroenteritis (due to bacterial pathogens) and hepatitis A have recently been reported among persons drinking contaminated water (10), this is the first report in several years of outbreaks of these illnesses occurring jointly after shellfish consumption (11,12). Viral gastroenteritis in association with hepatitis A is not known to have been reported following shellfish consumption. The clinical findings observed in several of the New York outbreaks are compatible with a viral etiology, such as the Norwalk virus: a short incubation period, abrupt onset of upper and/or lower gastrointestinal illness, and brief duration (1-2 days). The absence of bacterial pathogens and the visualization of virus-like particles in clams from one outbreak further support a viral etiology in several of these outbreaks.

The recent New York State outbreaks may be related to periods of heavy rain and flooding. Run-off at these times, especially when sewage systems overflow, characteristically increases coliform counts in monitored coastal waters. However, the numerous outbreaks in New York before the May-June flooding suggest an endemic degree of clam contamination, some of which may be attributable to harvesting from

uncertified, sewage-contaminated waters. This practice is likely to continue, because taking clams from highly populated, polluted beds is economically profitable and difficult to prevent. These outbreaks emphasize that clams may contain multiple enteric pathogens, including viruses, and consumption of clams--especially raw or partially cooked--continues to pose substantial risk of transmitting disease. Although the most effective way of avoiding the problem is to prevent the distribution of illegally gathered, untagged clams, such measures are not always possible. Therefore, because steaming or other forms of cooking do not always kill the enteric viruses in clams (13,14), the most effective means of preventing clam-associated illness is to adequately deplete them.

#### References

1. Earampamoorthy S, Koff RS. Health hazards of bivalve-mollusk ingestion. *Ann Intern Med* 1975;83:107-10.
2. Lumsden LL, Hasseltine HE, Leake JP, Veldee MV. A typhoid fever epidemic caused by oyster-borne infection (1924-1925). *Public Health Rep* 1925;Suppl 50:1-102
3. Roos B. Hepatitis epidemic transmitted by oysters. *Svenska lak-tidning* 1956;53:989-1003.
4. Mason JO, McLean WR. Infectious hepatitis traced to the consumption of raw oysters. An epidemiologic study. *Am J Hyg* 1962;75:90-111.
5. McIntyre RC, Tira T, Flood T, Blake PA. Modes of transmission of cholera in a newly infected population on an atoll: implication for control measures. *Lancet* 1979;1:311-4.
6. Barker WH. Vibrio parahaemolyticus outbreaks in the United States. *Lancet* 1974;1:551-4.
7. Wilson R, Lieb S, Roberts A, et al. Non-O group 1 Vibrio cholerae gastroenteritis associated with eating raw oysters. *Am J Epidemiol* 1981;114:293-8.
8. Murphy AM, Grohmann GS, Christopher PJ, Lopez WA, Davey GR, Millsom RH. An Australia-wide outbreak of gastroenteritis from oysters caused by Norwalk virus. *Med J Aust* 1979;2:329-33.
9. Gunn RA, Janowski HT, Lieb S, Prather EC, Greenberg HB. Norwalk virus gastroenteritis following raw oyster consumption. *Am J Epidemiol* 1982;115:348-51.
10. Sanchez Y, LaBelle RL, Hejkol T. Identification of hepatitis A antigen in sewage and well water prior to an outbreak of waterborne infectious hepatitis. Environmental aspects of viral hepatitis transmission. Proceedings of the Third International Symposium. In: Szmuness W, Alter HJ, Maynard JE, eds. on *Viral Hepatitis Philadelphia: Franklin Institute Press* 1981:629-30.
11. Dismukes WE, Bisno AL, Katz S, Johnson RF. An outbreak of gastroenteritis and infectious hepatitis attributed to raw clams. *Am J Epidemiol* 1969;89:555-61.
12. Begg RC. Food poisoning--four unusual episodes. *NZ Med J* 1975;82:52-4.
13. Koff RS, Sear HS. Internal temperature of steamed clams. *N Engl J Med* 1967;276:737-9.
14. Feachem R, Garelick H, Slade J. Enteroviruses in the environment. *Trop Dis Bull* 1981;78:185-230.

#### Outbreak of Yersinia enterocolitica--Washington State MMWR 1982;31:562-4

In December 1981 and January 1982, an outbreak of predominantly gastroenteritis caused by Yersinia enterocolitica occurred among 87 persons in Washington state. The illness was associated with the ingestion of a locally produced brand of tofu, an oriental soybean curd, packed in untreated spring water. It was sold primarily in western Washington with limited distribution in Alaska, Idaho, and Oregon. Y.

enterocolitica was isolated from the tofu, the processing plant's water supply, and several sites within the plant.

In mid-January 1982, the Seattle-King County Health Department received reports from two hospital laboratories of 12 positive stool cultures of Y. enterocolitica associated with gastrointestinal illness; during the previous year, 10 Y. enterocolitica isolates were reported in the entire county. Increased surveillance by Seattle-King County and the Washington State Department of Social and Health Services over several months identified additional cases, for a total of 87. A case was defined as anyone who was culture-positive and/or who had had contact with a case and had fever in conjunction with diarrhea or abdominal cramps.

Of the 87 cases, 56 were culture-positive; 38 patients had enteritis, six had only extra-intestinal infections, four had both extra-intestinal infections and enteritis, and eight were asymptomatic carriers. The 10 cases of extra-intestinal infection included patients with wound ulcers (two), inguinal lymphadenopathy (two, one with a perineal ulcer), pneumonia (two), labial infection (one), arthritis (one), septicemia (one), and pharyngitis (one). Nine (16.1%) of the 56 culture-positive patients were < 1 year of age, 12 (21.4%) were 1 to 4 years old, 7 (12.5%) were 5 to 18 years old, and 28 (50.0%) were > 18 years old. Among 38 culture-positive cases with enteritis, who tended to have more severe illness and on whom more complete information was available, the following were reported: fever (91%), abdominal pain (81%), diarrhea (76%), nausea (54%), vomiting (39%), bloody stools (27%), joint pain (42%), and skin rash (43%). Symptoms lasted from 1 day to 4 weeks (mean 10 days). Two patients, however, were ill for over 2 months. Seventeen patients were hospitalized for from 2 to 11 days (average 9.7 days); two of those hospitalized had appendectomies and one, a partial colectomy. One patient was also culture-positive for Salmonella typhimurium and one for rotavirus, as well as for Y. enterocolitica.

A neighborhood case-control study of 11 ill persons and 11 controls revealed an association between Y. enterocolitica infection and tofu consumption ( $p < 0.01$ ). Questions regarding animal contacts, water sources, raw milk consumption, travel, and day-care settings, as well as extensive food histories, did not identify any other common sources. Further investigation revealed that 70 (80.5%) of the 87 persons interviewed had consumed the same brand of tofu within the 2 weeks before onset of symptoms. For five culture-positive persons who had consumed only one meal of tofu, the incubation period averaged 6.6 days (range 4-11 days).

The tofu plant in King County is located on a rural island in Puget Sound. The plant water supply, which originates from a spring approximately 0.5 mile from the plant, is shared by four residences and an apple-cider plant. No illness was reported among consumers of the cider. Inspection of the tofu plant on January 20, 1982, disclosed unsanitary conditions, including poor personal hygiene, use of an outdoor privy, and unsanitary equipment. Samples of tofu and the plant water supply were positive for Y. enterocolitica, as were stool specimens collected from two of 12 employees; both employees were asymptomatic.

A voluntary recall of the product was instituted from January 21 to January 25. Further sampling and laboratory analysis of the tofu demonstrated high fecal-coliform counts. Production was resumed after a water-purification system was installed. Laboratory results of plasmid analysis, determination of enterotoxin production, and serotyping are pending.

#### Multi-State Outbreak of Yersiniosis MMWR 1982;31:505-6

Between June 11 and July 29, 1982, a large interstate outbreak of enteritis caused by Yersinia enterocolitica occurred. State health departments became aware of a potential problem when hospitals reported increased numbers of Y. enterocolitica

isolates. Epidemiologic investigation implicated milk pasteurized at a plant in Memphis, Tennessee, as the vehicle of infection.

One hundred seventy-two culture-positive Y. enterocolitica infections were identified; 67 in the Little Rock, Arkansas, area; 80 in Memphis, Tennessee, and its northern Mississippi suburbs; and 25 in the Greenwood, Mississippi, area. One hundred forty-eight (86%) patients had enteric infections with diarrhea and/or abdominal pain, usually accompanied by fever; 24 patients had extra-intestinal infections of throat, blood, urinary tract, central nervous system, and wounds. Forty-one percent of cases occurred among children less than 5 years of age. Most patients required hospitalization, and 17 underwent appendectomies. The epidemic strain is agglutinated most strongly by antisera to Y. enterocolitica O groups 13 and 18.

Separate case-control studies in each city showed that drinking milk pasteurized by a milk plant in Memphis was associated with illness (in Little Rock,  $p=0.03$ ; in Memphis,  $p=0.01$ ; in Greenwood,  $p=0.004$ ). Overall, 71% of cases and 39% of controls recalled drinking milk from the plant in the 2 weeks before onset of symptoms.

In an effort to estimate the size of the outbreak, a survey was made by telephone of 100 randomly chosen households in Greenwood. Heads of household were queried concerning illness and milk drinking history within the last two months. Eleven cases of yersiniosis-like illness, defined as either 1) fever  $\geq 101$  F (38.3 C) and diarrhea or 2) fever  $\geq 101$  F and abdominal pain at any time during the previous 6 weeks, were identified among the 260 members of these households. All patients resided in households that used milk from the implicated plant, and 10 of the 11 (91%) recalled drinking its milk within the previous 2 months. Illness occurred in 6 of 50 (12%) households that used milk from the implicated plant and in none of 50 that did not use its milk ( $p=0.02$ , Fisher's exact test). Of those individuals who drank milk from that plant, 8.7% had a yersiniosis-like illness. Based on a census of 20,115 and the number of the Memphis plant milk drinkers in Greenwood, it was estimated that 857 cases (95% confidence limits 363.5-1,351.7) may have occurred in Greenwood where only 3.9% of the plant's milk is sold. The total number of cases in all three states, therefore, would appear to be higher than the 172 cases reported.

The outbreak appeared to end spontaneously. Milk from suspected lots was not available for culture, and Y. enterocolitica was not isolated from subsequent lots. A Food and Drug Administration laboratory isolated Y. enterocolitica of the same serotype found in the outbreak from a milk crate on a hog farm where outdated milk from the implicated plant is fed to hogs. Inspection of the plant identified neither a breach in pasteurizing technique nor an obvious source of contamination. Surveillance for new cases and surveillance of milk for Yersinia have continued.

Editorial Note: In this investigation pasteurized milk was epidemiologically implicated as the vehicle of transmission of Y. enterocolitica. The temporal and geographic clustering of cases and the negative cultures of subsequent lots of milk are consistent with contamination of a single lot. The mechanism of contamination is unknown.

Y. enterocolitica may be found in raw milk (1,2); contaminated raw milk was responsible for an outbreak of yersiniosis among children in Montreal (3). The organism has also been found in pasteurized milk (1,4) although not associated with illness. Y. enterocolitica generally does not survive standard pasteurization (5); however, if present in large enough numbers, viable Yersinia may persist after pasteurization (4-6). Once present in a pasteurized product, the organism grows well at refrigeration temperature (7). Therefore, pasteurization and proper handling of pasteurized milk may not ensure against enteric disease due to Y. enterocolitica.

Only two other well documented food-borne outbreaks of Y. enterocolitica enteritis have been reported in the United States: one in New York state in 1976 caused by contaminated chocolate milk (8) and one in Washington state in 1982 caused by tofu (9). Food-borne transmission of yersiniosis has also been suspected in other

outbreaks (10-12). This is the largest outbreak of yersiniosis ever reported in the United States.

#### References

1. Schiemann DA. Association of Yersinia enterocolitica with the manufacture of cheese and occurrence in pasteurized milk. *Appl Env Microbiol* 1978;36:274-7.
2. Vidon DJ, Delmas CL. Incidence of Yersinia enterocolitica in raw milk in eastern France. *Appl Environ Microbiol* 1981;41:355-9.
3. Yersinia enterocolitica gastroenteritis outbreak--Montreal. *Canada Diseases Weekly Report* 1976;2:41-4.
4. Hughes D. Isolation of Yersinia enterocolitica from milk and a dairy farm in Australia. *J Appl Bacteriol* 1979;46:125-30.
5. Francis DW, Spaulding PL, Lovett J. Enterotoxin production and thermal resistance of Yersinia enterocolitica in milk. *Appl Environ Microbiol* 1980;40:174-6.
6. Schiemann DA, Toma S. Isolation of Yersinia enterocolitica from raw milk. *Appl Environ Microbiol* 1978;35:54-8.
7. Morris GK, Feeley JC, Martin WT, Wells JG. Isolation and identification of Yersinia enterocolitica. *The Public Health Laboratory* 1977;35:217-35.
8. Black RE, Jackson RJ, Tsai T, et al. Epidemic Yersinia enterocolitica infection due to contaminated chocolate milk. *N Engl J Med* 1978; 298:76-9.
9. Washington State Department of Social and Health Services. Unpublished report.
10. Olsovsky Z, Olsakova V, Chobot S, Sviridov V. Mass occurrence of Yersinia enterocolitica in two establishments of collective care of children. *J Hyg Epidemiol Microbiol Immunol (Praha)* 1975;19:22-9.
11. Asakawa Y, Akahane S, Kagata N, Noguchi M, Sakazaki R, Tamura K. Two community outbreaks of human infection with Yersinia enterocolitica. *J Hyg (Lond)* 1973;71:715-23.
12. Shayegani M, Morse D, DeForge I, Root T, Parsons LM, Maupin P. Foodborne outbreak of Yersinia enterocolitica in Sullivan County, New York, with pathologic studies of the isolates. Abstracts of the Annual Meeting of the American Society of Microbiology, 1982, No. C175.

## J. Bibliography

### GENERAL

1. Bryan FL. Emerging foodborne diseases. I. Their surveillance and epidemiology. II. Factors that contribute to outbreaks and their control. *J Milk Food Technol* 1972;35:618-25, 632-8.
2. Bryan FA. Factors that contribute to outbreaks of foodborne disease. *J Food Protection* 1978;41:816-27.
3. Bryan FL. Foodborne diseases in the United States associated with meat and poultry. *J Food Protection* 1980;43:140-50.
4. Horwitz MA. Specific diagnosis of foodborne disease. *Gastroenterology* 1977;73:375-81.
5. Riemann H, ed. Foodborne infections and intoxications. New York: Academic Press, 1969.
6. Sours HE, Smith DG. Outbreaks of foodborne disease in the United States, 1972-1978. *J Infect Dis* 1980;142:122-5.
7. Potter ME, Kaufmann AF, Blake PA, Feldman RA. Unpasteurized milk. The hazards of a health fetish. *JAMA* 1984;252:2048-52
8. Barker WH, Sagerson JC, Hall CVH, Anderson HW, Francis BJ. Foodborne Surveillance--Washington State. *Am J Public Health* 1974;64:853-9.

### BACTERIAL

#### Bacillus cereus

1. Giannella RA, Brasile L. A hospital foodborne outbreak of diarrhea caused by Bacillus cereus: Clinical, epidemiologic, and microbiologic studies. *J Infect Dis* 1979;139:366-70.
2. Mortimer PR, McCann G. Food poisoning episodes associated with Bacillus cereus in fried rice. *Lancet* 1974;1:1043-5.
3. Terranova W, Blake PA. Bacillus cereus food poisoning. *N Engl J Med* 1978;298:143-4.
4. Turnbull PCB, Kramer JM, Torgensen K, Gilbert RJ, Melling J. Properties and production characteristics of vomiting, diarrheal, and necrotizing toxins of Bacillus cereus. *Am J Clin Nutr* 1979;32:219-28.

#### Brucella

1. Fox MD, Kaufman AF. Brucellosis in the United States, 1965-1974. *J Infect Dis* 1977;136:312-6.
2. Young EJ, Suvannaparrat U. Brucellosis outbreak attributed to ingestion of unpasteurized goat cheese. *Arch Intern Med* 1975;135:240-3.
3. Young, EJ. Human Brucellosis. *Rev Infect Dis* 1983;5:821-842.
4. Centers for Disease Control. Brucellosis--Texas. *MMWR* 1983;32:548,553.

#### Campylobacter

1. Blaser MJ, Reller LB. Campylobacter enteritis. *N Engl J Med* 1981;305:1444-52.
2. Blaser MJ, Checko P, Bopp C, Bruce A, Hughes JM. Campylobacter enteritis associated with foodborne transmission. *Am J Epidemiol* 1982;116:886-94.
3. Blaser MJ, Taylor DN, Feldman RA. Epidemiology of Campylobacter jejuni infections. *Epidemiol Rev* 1983;5:157-76.
4. Hopkins RS, Olmsted R, Istre GR. Endemic Campylobacter jejuni infection in Colorado: Identified risk factors. *Am J Public Health* 1984;74:249-50.
5. Istre GR, Blaser MJ, Shillam P, Hopkins RS. Campylobacter enteritis associated with undercooked barbecued chicken. *Am J Public Health* 1984;74:1265-67.

6. Blaser MJ, Cravens J, Powers BW, LaForce FM, Wang WLL. Campylobacter enteritis associated with unpasteurized milk. Am J Med 1979;67:715-18.

7. Taylor, Porter BW, Williams CA, Miller HG, Bopp CA, Blake PA. Campylobacter enteritis: A large outbreak traced to commercial raw milk. West J Med 1982;137:365-9.

#### Clostridium botulinum

1. Center for Disease Control. Botulism in the United States, 1899-1977. Handbook for Epidemiologists, Clinicians, and Laboratory Workers. Atlanta: CDC, May 1979.

2. Blake PA, Horwitz MA, Hopkins L, Lombard GL, McCroan JE, Prucha JC, Merson MH. Type A botulism from commercially canned beef stew. South Med J 1977;70:5-7.

3. Hughes JM, Blumenthal JR, Merson MH, Lombard GL, Dowell VR, Gangarosa EJ. Clinical features of types A and B food-borne botulism. Ann Intern Med 1981;95:442-5.

4. Seals JE, Snyder JD, Edell TA, Hatheway CL, Johnson CJ, Swanson RC, Hughes JM. Restaurant-associated type A botulism: Transmission by potato salad. Am J Epidemiol 1981;113:436-44.

5. Tacket CO, Shandera WX, Mann JM, Hargrett NT, Blake PA. Equine antitoxin use and other factors that predict outcome in type A foodborne botulism. Am J Med 1984;76:794-798.

6. Terranova W, Breman JG, Locey RP, Speck S. Botulism type B: Epidemiologic aspects of an extensive outbreak. Am J Epidemiol 1978;108:150-6.

7. Mann JM, Hatheway CL, Gardiner TM. Laboratory diagnosis in a large outbreak of type A botulism. Am J Epidemiol 1982;115:598-605.

8. MacDonald KL, Spengler RF, Hatheway CL, Hargrett NT, Cohen ML. Type A botulism from sauteed onions: Clinical and epidemiologic observations. JAMA 1985;253:1275-1278.

#### Clostridium perfringens

1. Bryan FL. What the sanitarian should know about Clostridium foodborne illness. J Milk Food Technol 1969;32:381-9.

2. Lowenstein MS. Epidemiology of Clostridium perfringens food poisoning. N Engl J Med 1972;286(19):1026-7.

3. Shandera WX, Tacket CO, Blake PA. Food poisoning due to Clostridium perfringens in the United States. J Infect Dis 1983;143:167-170.

4. Stringer MF, Turnbull PCB, Gilbert RJ. Application of serological typing to the investigation of outbreaks of Clostridium perfringens food poisoning, 1970-1978. J Hyg (Camb) 1980;84:443-56.

#### Escherichia coli

1. Marier R, Wells JG, Swanson RC, Callahan W, Mehlman IJ. An outbreak of enteropathogenic Escherichia coli foodborne disease traced to imported French cheese. Lancet 1973;2:1376-8.

2. Taylor WR, Schell WL, Wells JG, Choi K, Kinnunen DE, Heiser PT, Helstad AG. A foodborne outbreak of enterotoxigenic Escherichia coli diarrhea. N Engl J Med 1982;306:1093-95.

3. Gangarosa EJ. Epidemiology of Escherichia coli in the U.S. J Infect Dis 1978;137:634-8.

4. Lumish RM, Ryder RW, Anderson DC, Wells JG, Puhr ND. Heat-labile enterotoxigenic Escherichia coli induced diarrhea aboard a Miami-based cruise ship. Am J Epidemiol 1980;111:432-36.

5. Riley LW, Remis RS, Helgerson SD, McGee HB, Wells JG, Davis BR, Hebert RJ, Olcott ES, Johnson LM, Hargrett NT, Blake PA, Cohen ML. Hemorrhagic colitis associated with a rare Escherichia coli serotype. N Engl J Med 1983;308:681-85.

6. MacDonald KL, Eidson M, Strohmeyer C, Levy ME, Wells JG, Puhr ND, Wachsmuth K, Hargrett NT, Cohen ML. A multistate outbreak of gastrointestinal illness caused by enterotoxigenic Escherichia coli in imported semisoft cheese. *J Infect Dis* 1985; 151:716-720.

#### Listeria monocytogenes

1. Schlech WF III, Lavigne PM, Bortolussi RA, Allen AC, Haldane EV, Wort AJ, Hightower AW, Johnson SE, King SH, Nicholls ES, Broome CV. Epidemic listeriosis--evidence for transmission by food. *N Engl J Med* 1983;308:203-206.

2. Fleming DW, Cochi SL, MacDonald KL, Brondum J, Hayes PS, Plikaytis BD, Holmes MB, Audurier A, Broome CV, Reingold AL. Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. *N Engl J Med* 1985;312:404-407.

#### Salmonella

1. Cohen ML, Blake PA. Trends in foodborne salmonellosis outbreaks: 1963-1975. *J Food Protection* 1977;40:798-800.

2. Blaser MJ, Rafuse EM, Wells JG, Pollard RA, Feldman RA. An outbreak of salmonellosis involving multiple vehicles. *Am J Epidemiol* 1981;114:663-670.

3. Blaser MJ, Newman LS. A review of human salmonellosis. I. Infective dose. *Rev Infect Dis* 1982;4:1096-106.

4. Buchwald DS, Blaser MJ. A review of human salmonellosis: II. Duration of excretion following infection with non-typhi Salmonella. *Rev Infect Dis* 1982;6:345-356.

5. Cohen ML, Fontaine RE, Pollard RA, VonAllmen SD, Vernon TM, Gangarosa EJ. An assessment of patient-related economic costs in an outbreak of salmonellosis. *N Engl J Med* 1978;299:459-60.

6. Fontaine RE, Arnon S, Martin WT, Vernon TM Jr, Gangarosa EJ, Farmer JJ, Moran AB, Silliker JH, Decker DL. Raw hamburger: an interstate common source of human salmonellosis. *Am J Epidemiol* 1978;107:36-45.

7. Gunn RA, Markakis G. Salmonellosis associated with homemade ice cream. An outbreak report and summary of outbreaks in the U.S. in 1966 to 1976. *JAMA* 1978;240:1885-6.

8. Holmberg SD, Osterholm MT, Senger KA, Cohen ML. Drug-resistant Salmonella from animals fed antimicrobials. *N Engl J Med* 1984;311:617-22.

9. Holmberg SD, Wells JG, Cohen ML. Animal-to-man transmission of antimicrobial-resistant Salmonella: Investigations of U.S. outbreaks, 1971-1983. *Science* 1984;225:833-835.

10. Taylor DN, Bied JM, Munro JS, Feldman RA. Salmonella dublin infections in the United States, 1979-80. *J Infect Dis* 1982;146:322-327.

11. Klotz SA, Jorgensen JH, Buckwold FJ, Craven PC. Typhoid fever. An epidemic with remarkably few clinical signs and symptoms. *Arch Intern Med* 1984;144:533-37.

12. Tacket CO, Dominguez LB, Fisher HJ, Cohen ML. An outbreak of multiple-drug-resistant Salmonella enteritis from raw milk. *JAMA* 1985;253:2058-2060.

#### Shigella

1. Black RE, Craun GF, Blake PA. Epidemiology of common-source outbreaks of shigellosis in the United States, 1961-1975. *Am J Epidemiol* 1978;108:47-52.

2. Donadio JA, Gangarosa EJ. Foodborne shigellosis. *J Infect Dis* 1969;119:666-8.

3. Blaser MJ, Pollard RA, Feldman RA. Shigella infections in the United States, 1974-1980. *J Infect Dis* 1983;147:771-75.

#### Staphylococcus

1. Bryan FL. What the sanitarian should know about Salmonellae and Staphylococci in non-dairy foods. I. Staphylococci. *J Milk Food Technol* 1968;31:110-16.

2. Holmberg SC, Blake PA. Staphylococcal food poisoning in the United States. New facts and old misconceptions. JAMA 1984;251:487-89.

3. Merrill GA, Werner SB, Bryant RG, Fredson D, Kelly K. Staphylococcal food poisoning associated with an Easter egg hunt. JAMA 1984;252:1019-22.

#### Streptococcus

1. Hill HR, Zimmerman RA, Reid GVK, Wilson E, Kitton RM. Foodborne epidemic of streptococcal pharyngitis at the United States Air Force Academy. N Engl J Med 1969;280:917-21.

2. McCormick JB, Kay D, Hayes M, Feldman RA. Epidemic streptococcal sore throat following a community picnic. JAMA 1976;236:1039-41.

#### Vibrio cholerae O1

1. Blake PA, Allegra DT, Snyder JD, Barrett TJ, McFarland L, Caraway CT, Feeley JC, Craig JP, Lee JV, Puhr ND, Feldman RA. Cholera--A possible endemic focus in the United States. N Engl J Med 1980;302:305-309.

2. Blake PA. Prevention of food-borne diseases caused by Vibrio species. In: Colwell R, ed. Vibrios in the environment. John Wiley & Sons, 1984:579-591.

3. Blake PA. Vibrios on the half-shell: What the walrus and the carpenter didn't know. Ann Intern Med 1983;99:558-9.

4. Holmberg SD, Harris JR, Kay DE, Hargrett NT, Parker RD, Kansou N, Rao NU, Blake PA. Foodborne transmission of cholera in Micronesian households. Lancet 1984;325-28.

#### Vibrio cholerae Non-O1

1. Blake PA, Weaver RE, Hollis DG. Diseases of humans (other than cholera) caused by Vibrios. Ann Rev Microbiol 1980;34:341-67.

2. Hughes JM, Hollis DG, Gangarosa EJ, Weaver RE. Non-cholera vibrio infections in the United States--Clinical, epidemiologic, and laboratory features. Ann Intern Med 1978;88:602-6.

3. Morris JG, Wilson R, Davis BR, Wachsmuth IK, Conradine FR, Riddle BA, Wathen HG, Pollard RA, Blake PA. Non-O Group 1 Vibrio cholerae gastroenteritis in the United States. Ann Intern Med 1981;94:656-8.

4. Wilson R, Lieb S, Robers A, Stryker S, Janowski H, Gunn R, Davis B, Riddle CF, Barrett T, Morris JG, Blake PA. Non-O group 1 Vibrio cholerae gastroenteritis associated with eating raw oysters. Am J Epidemiol 1981;114:293-8.

#### Vibrio parahaemolyticus

1. Barker WH. Vibrio parahaemolyticus outbreaks in the United States. Lancet 1974;1:551-4.

2. Lawrence DN, Blake PA, Yashuk JC, Wells JG, Creech WB, Hughes JH. Vibrio parahaemolyticus gastroenteritis outbreaks aboard two cruise ships. Am J Epidemiol 1979;10:71-80.

#### Yersinia enterocolitica

1. Black RE, Jackson RJ, Tsai T, Medvesky M, Shayegani M, Feeley JC, MacLeod KIE, Wakelee AM. Epidemic Yersinia enterocolitica infection due to contaminated chocolate milk. N Engl J Med 1978;298:76-9.

2. Tacket CO, Narain JP, Sattin R, Lofgren JP, Konigsberg C, Rendtorff RC, Rausa A, Davis BR, Cohen ML. A multistate outbreak of infections caused by Yersinia enterocolitica transmitted by pasteurized milk. JAMA 1984;251:483-6.

3. Tacket CO, Ballard J, Harris N, Allard J, Nolan C, Quan T, Cohen ML. An outbreak of Yersinia enterocolitica infections caused by contaminated tofu (soybean curd). Am J Epidemiol 1985;121:705-711.

## CHEMICAL

### Heavy Metals

#### Cadmium

1. Baker TD, Hafnew WG. Cadmium poisoning from a refrigerator shelf used as an improvised barbecue grill. Public Health Rep 1961;76:543-4.

#### Copper

1. Hopper SH, Adams HS. Copper poisoning from vending machines. Public Health Rep 1958;73:910-4.

2. Semple AB, Parry WH, Phillips DE. Acute copper poisoning: An outbreak traced to contaminated water from a corroded geyser. Lancet 1960;2:700-1.

#### Tin

1. Barker WH, Runte V. Tomato juice-associated gastroenteritis. Washington and Oregon, 1969. Am J Epidemiol 1972;96:219-26.

#### Zinc

1. Brown MA, Thom JV, Orth GL, et al. Food poisoning involving zinc contamination. Arch Environ Health 1964;8:657-60.

### Ciguatoxin

1. Halstead BW. Poisonous and venomous marine animals of the world. Princeton: Darwin Press, 1978, pp 325-402.

2. Lawrence DN, Enriquez MB, Lumish RM, Maceo A. Ciguatera fish poisoning in Miami. JAMA 1980;244:254-8.

3. Engleberg NC, Morris JG, Lewis J, McMillen JP, Pollard RA, Blake PA. Ciguatera fish poisoning: A major common-source outbreak in the U.S. Virgin Islands. Ann Intern Med 1983;19:336-337.

4. Morris JG, Lewin P, Smith CW, Blake PA, Schneider R. Ciguatera fish poisoning: Epidemiology of the disease on St. Thomas, U.S. Virgin Islands. Am J Trop Med Hyg 1982;31:574-8.

5. Morris JG, Lewin P, Hargrett NT, Smith CW, Blake PA, Schneider R. Clinical features of ciguatera fish poisoning: A study of the disease in the U.S. Virgin Islands. Arch Intern Med 1982;142:1090-2.

### Puffer Fish (tetrodotoxin)

1. Halstead BW. Poisonous and venomous marine animals of the world. Princeton: Darwin Press, 1978, pp 437-548.

### Scombrototoxin

1. Arnold SH, Brown WD. Histamine toxicity from fish products. Adv Food Res 1978;24:113-54.

2. Gilbert RJ, Hobbs G, Murray CK, Cruickshank JG, Young SEJ. Scombrototoxic fish poisoning: Features of the first 50 incidents to be reported in Britain (1976-1979). Br Med J 1980;281:71-2.

3. Halstead BW. Poisonous and venomous marine animals of the world. Princeton: Darwin Press, 1978, pp 417-35.

4. Hughes JM, Merson MH. Fish and shellfish poisoning. N Engl J Med 1976;295:1117-20.

5. Merson MH, Baine WB, Gangarosa EJ, Swanson RC. Scombroid fish poisoning: Outbreak traced to commercially canned tuna fish. JAMA 1974;228:1268-9.

### Monosodium Glutamate

1. Schaumburg HH, Byck R, Gerstl R, Mashman JH. Monosodium L-glutamate; its pharmacology and role in the Chinese restaurant syndrome. Science 1969;163:826-8.

### Mushroom Poison

1. Hanrahan JP, Gordon MA. Mushroom Poisoning: The reports and a review of therapy. JAMA 1984;251:1057-61.
2. Mitchel DH. Amanita mushroom poisoning. Ann Rev Med 1980;31:51-7.
3. Tyler VE. Poisonous mushrooms: Progress in chemical toxicology. Vol 1, Stolman A (ed). New York: Academic Press, 1963, pp 339-84.

### Paralytic and Neurotoxic Shellfish Poison

1. Halstead BW. Poisonous and venomous marine animals of the world. Princeton: Darwin Press, 1978, pp 43-78.
2. Hughes JM, Merson MH. Fish and shellfish poisoning. N Engl J Med 1976;295:1117-20.
3. Music SI, Howell JT, Brumback CL. Red tide: its public health implications. J Fla Med Assoc 1973;60:27-9.

### PARASITIC

1. Osterholm MT, Forfang JC, Ristinen TL, Dean AG, Washburn JW, Godes JR, Rude RA, McCullough JG. An outbreak of foodborne giardiasis. N Engl J Med 1981;304:24-28.
2. Chitwood MD. Nematodes of medical significance found in market fish. Am J Trop Med Hyg 1970;19:599-602.
3. Gould SE. Trichinosis in man and animals. Springfield, Ill.: Charles C. Thomas, 1970.
4. Zimmerman WJ, Steele JH, Kagan IG. Trichinosis in the U.S. population 1966-1970--Prevalence and epidemiologic factors. Health Services Rep 1973;88:606-23.

### VIRAL

#### Hepatitis A

1. Carl M, Francis DP, Maynard JE. Food-borne hepatitis A: Recommendations for control. J Infec Dis 1983;148:1133-35.
2. Osterholm MT, Kantor RJ, Bradley DW, Hall WN, Francis DP, Aaron HC, Washburn JW, Velde D. Immunoglobulin M-specific serologic testing in an outbreak of foodborne viral hepatitis, type A. Am J Epidemiol 1980;112:8-16.

#### Norwalk Virus

1. Kuritsky JN, Osterholm MT, Greenberg HB, Korlath JA, Godes JR, Hedberg CW, Forfang JC, Kapikian AZ, McCullough JC, White KE. Norwalk gastroenteritis: A community outbreak associated with bakery product consumption. Ann Intern Med 1984;100:519-21.
2. Kaplan JE, Gary GW, Baron RC, Singh N, Schonberger LB, Feldman R, Greenberg HB. Epidemiology of Norwalk gastroenteritis and the role of Norwalk virus in outbreaks of acute non-bacterial gastroenteritis. Ann Intern Med 1982;96:756-61.

7/15/85

## STATE AND TERRITORIAL EPIDEMIOLOGISTS AND STATE PUBLIC HEALTH LABORATORY DIRECTORS

The State and Territorial Epidemiologists are the key to all disease surveillance activities, and their contributions to this report are gratefully acknowledged. In addition, valuable contributions are made by State Laboratory Directors.

<u>State</u>	<u>State Epidemiologist</u>	<u>State Laboratory Director</u>
Alabama	Wallace E. Birch, DVM	James L. Holston, Jr., DrPH
Alaska	John P. Middaugh, MD	Harry J. Colvin, PhD
Arizona	Norman J. Petersen, SM	Jon M. Counts, DrPH
Arkansas	Thomas C. McChesney, DVM, Acting	Robert I. Horn
California	James Chin, MD	Thaddeus F. Midura, Acting
Colorado	Stanley W. Ferguson, PhD	Robt. J. Barr, Co-Acting Dir
Connecticut	James L. Hadler, MD	Jesse Tucker, PhD
Delaware	Paul R. Silverman, DrPH	Mahadeo P. Verma, PhD
District of Columbia	Martin E. Levy, MD	James R. Thomas, PhD, Acting
Florida	John J. Witte, MD, Acting	Eldert C. Hartwig, Jr., ScD
Georgia	R. Keith Sikes, DVM	Frank M. Rumph, MD
Guam	Robert L. Haddock, DVM	Luis P. Flores
Hawaii	Arthur P. Liang, MD	Glenn Kobayashi
Idaho	Charles D. Brokopp, DrPH	D. W. Brock, DrPH
Illinois	Byron J. Francis, MD	Harry C. Rostick
Indiana	Charles L. Barrett, MD	T. L. Eddleman
Iowa	Laverne A. Wintermeyer, MD	W. J. Hausler, Jr., PhD
Kansas	Robert French, Acting	Roger H. Carlson, PhD
Kentucky	M. Ward Hinds, MD	B. F. Brown, MD
Louisiana	Louise McFarland, Dr.PH, Acting	Henry Bradford, PhD
Maine	Kathleen F. Gensheimer, MD	Philip W. Haines, DrPH
Maryland	Ehenezer Israel, MD	J. Mehnen Joseph, PhD
Massachusetts	George F. Grady, MD	George F. Grady, MD
Michigan	Kenneth R. Wilcox, Jr., MD	George R. Anderson, DVM
Micronesia*	Eliuel K. Pretrick, MD	Vacant
Minnesota	Michael Osterholm, PhD, MPH	C. Dwayne Morse, DrPH
Mississippi	Fred E. Thompson, MD	R. H. Andrews, MS
Missouri	H. Denny Donnell, Jr., MD	Elmer R. Spurrier, DrPH
Montana	Judith K. Gedrose, RN	Douglas Abbott, PhD
Nebraska	Paul A. Stoesz, MD	John Blosser
Nevada	Geo. E. Reynolds, MD, Acting	George Reynolds, MD
New Hampshire	Eugene Schwartz, MD	Mrs. Veronica C. Malmberg
New Jersey	William E. Parkin, DVM	Bernard F. Taylor, PhD
New Mexico	Harry F. Hull, MD	Loris W. Hughes, PhD
New York State	Richard Rothenberg, MD	David O. Carpenter, MD
New York City	Stephen Schultz, MD	Vacant
North Carolina	J. N. MacCormack, MD	Mildred A. Kerbaugh
North Dakota	James L. Pearson, DrPH	A. A. Gustafson
Northern Mariana Islands**	Jose T. Villagomez, MD	Vacant
Ohio	Thomas J. Halpin, MD	Garv D. Davidson, DrPH
Oklahoma	Gregory R. Istre, MD	Garry L. McKee, PhD
Oregon	John A. Googins, MD	Michael R. Skeels
Palau*	Anthony H. Polloi, MD, Acting	Vacant
Pennsylvania	Ernest J. Witte, VMD	Vern Pidcoe, DrPH
Puerto Rico	Jose G. Rigau, MD	Jose L. Villamil
Rhode Island	Richard A. Keenlyside, MBBS	Raymond G. Lundgren, PhD
Somoa	Tofiga Lisiga, MD	Vacant
South Carolina	Richard L. Parker, DVM	Arthur F. DiSalvo, MD
South Dakota	Kenneth A. Senger	A. Richard Melton, DrPH
Tennessee	Robert H. Hutcheson, Jr., MD	Michael W. Kimberly, DrPH
Texas	Charles E. Alexander, MD	Charles F. Sweet, DrPH
Utah	Craig R. Nichols, MPA	Francis M. Urry, PhD
Vermont	Richard L. Vogt, MD	Katherine A. Kelley
Virginia	Grayson B. Miller, Jr., MD	Frank W. Lambert, Jr., DrPH
Virgin Islands	John N. Lewis, MD	Norbert Mantor, PhD
Washington	John M. Kobavashi, MD	Jack Allard, PhD
West Virginia	Loretta E. Haddy, MS	John W. Brough, DrPH
Wisconsin	Jeffrey P. Davis, MD	Ronald H. Laessig, PhD
Wyoming	Harry C. Crawford, MD	Donald T. Lee, PhD

\*Formerly Trust Territory of the Pacific Islands

HHS, PHS, CDC

**DEPARTMENT OF  
HEALTH & HUMAN SERVICES**

Public Health Service  
Centers for Disease Control  
Atlanta GA 30333

---

**Official Business**  
Penalty for Private Use \$300



Postage and Fees Paid  
U.S. Dept. of H.H.S.  
HHS 396