

Influenza A (H1N1) 2009 Monovalent Vaccine Dosage Chart

Inactivated, Injectable Influenza Vaccine				
Manufacturer	Age	Dose–Presentation	Number of Doses	Route–Site
sanofi pasteur	6 through 35 months ¹	0.25 mL–prefilled syringe ¹	2 ²	Intramuscular ³
	36 months and older	0.5 mL–prefilled syringe	1 or 2 ²	
	6 months and older	Dose per age–multidose vial		
Novartis Vaccine	4 years and older	5.0 mL–multidose vial	1 or 2 ²	Intramuscular ³
		0.5 mL–prefilled syringe		
CSL	6 months and older	0.5 mL–prefilled syringe	1 or 2 ²	Intramuscular ³
		5.0 mL–multidose vial		

- Children age 6 through 35 months should receive 0.25 mL vaccine per dose. Children age 36 months through adults should receive 0.5 mL vaccine per dose. See footnote 2 to determine number of doses.
- Based on currently available information, children 6 months through 9 years who are receiving injectable influenza A (H1N1) 2009 Monovalent vaccine should receive two doses of vaccine separated by 28 days.
- Children 6 months through 2 years of age should be vaccinated in the anterolateral aspect of the thigh. Older children and adults should be vaccinated in the deltoid muscle if muscle mass is adequate. The anterolateral aspect of the thigh may be used as an alternate.

Live Attenuated Nasal Spray Influenza Vaccine (LAIV)				
Manufacturer	Age	Dose–Presentation	Number of Doses	Route
MedImmune	2 through 49 years if healthy and non-pregnant	0.2 mL–Spray ½ of dose into each nostril as indicated on the syringe.	1 or 2 ⁴	Intranasal

- Based on currently available information, healthy children 2 through 9 years of age who are receiving live attenuated influenza A (H1N1) 2009 Monovalent vaccine should receive two doses of vaccine separated by 28 days.

For more information call **800-CDC-INFO** (800-232-4636) Website www.cdc.gov/flu

