Preventing Diarrhea After A Disaster

Protect yourself and your family:



Drink and use safe water.



Wash your hands often.



Do not poop in any body of water.



Eat safe food. Boil it, cook it, peel it, or throw it away.



Clean up safely.



Avoid areas with standing water.

Drink and use safe water.


- Listen to local officials to find out if your water is safe.
- Use bottled water for drinking, washing and preparing food, making ice, and brushing your teeth.
- If you do not have bottled water, boil or disinfect your water to make it safe.

How to make your water safe:

- If boiling, bring your water to a complete boil and keep boiling for at least 1 minute.
- To clean your water, use unscented household liquid chlorine bleach. If your water is clear, add 8 drops to 1 gallon of water. If your water is cloudy, use 16 drops to 1 gallon of water. Wait 30 minutes before drinking.

Wash your hands often with soap and safe water.

- Before you eat or prepare food.
- Before feeding your children.
- Before and after treating wounds or taking care of someone who is sick.
- After going to the bathroom, changing diapers, or cleaning a child after they have gone to the bathroom.



Eat safe food.

- Boil it, cook it, peel it, or throw it away.
- Avoid meat and dairy products that have not been refrigerated.
- Cook food well. Eat it hot and keep it covered.
- Avoid raw foods other than fruits and vegetables you have peeled yourself.

Clean up safely.

- Clean food preparation areas and kitchenware with soap and safe water and let dry completely before reuse.
- Wash yourself, your children, diapers, and clothes 100 feet away away from drinking water sources.

Avoid areas with standing water.

- Wash your hands with soap and water after contact with dirty water.
- Do not allow children to play in standing water.
- Do not allow children to play with toys that have touched dirty water and have not been cleaned.

What to do if you or your family are sick with diarrhea:

- Stay hydrated by drinking safe water.
- Visit the nearest health facility. Continue to keep hydrated while at home and while you travel to get treatment.
- Continue to breastfeed your baby if they have watery diarrhea, even when traveling to get treatment.





For more information about preventing illness after a disaster, visit: www.cdc.gov/disasters/disease/facts.html.