

## Commercially-bottled Water\*

Etiology	Ground Water				Unknown Water			
	Outbreaks	Cases	Hospitalized	Died	Outbreaks	Cases	Hospitalized	Died
<b>Bacteria, non-<i>Legionella</i> spp.</b>	0	0	0	0	1	22	0	0
<i>E. coli</i> O157:H7	0	0	0	0	0	0	0	0
<i>Pantoea agglomerans</i>	0	0	0	0	0	0	0	0
<i>Shigella sonnei</i>	0	0	0	0	1	22	0	0
<b>Bacteria, <i>Legionella</i> spp.</b>	0	0	0	0	0	0	0	0
<i>Legionella</i> spp.	0	0	0	0	0	0	0	0
<b>Virus</b>	0	0	0	0	0	0	0	0
Norovirus	0	0	0	0	0	0	0	0
<b>Parasite</b>	0	0	0	0	0	0	0	0
<i>Giardia</i>	0	0	0	0	0	0	0	0
<b>Chemical</b>	0	0	0	0	0	0	0	0
Suspected propylene glycol	0	0	0	0	0	0	0	0
<b>Multiple<sup>†</sup></b>	0	0	0	0	0	0	0	0
<b>Unidentified</b>	1	3	0	0	0	0	0	0
<b>Total</b>	1	3	0	0	1	22	0	0

\*No reported outbreaks in bottled water systems had mixed water or surface water source.

<sup>†</sup>One outbreak had multiple etiologic agent types. The etiologies were *Campylobacter* spp. (i.e., bacterium) and *Giardia intestinalis* (i.e., parasite).

