

Non-community Water Systems*

Etiology	Ground Water				Mixed Water [†]			
	Outbreaks	Cases	Hospitalized	Died	Outbreaks	Cases	Hospitalized	Died
Bacteria, non-<i>Legionella</i> spp.	1	56	2	0	0	0	0	0
<i>E. coli</i> O157:H7	1	56	2	0	0	0	0	0
<i>Pantoea agglomerans</i>	0	0	0	0	0	0	0	0
<i>Shigella sonnei</i>	0	0	0	0	0	0	0	0
Bacteria, <i>Legionella</i> spp.	0	0	0	0	0	0	0	0
<i>Legionella</i> spp.	0	0	0	0	0	0	0	0
Virus	2	138	0	0	0	0	0	0
Norovirus	2	138	0	0	0	0	0	0
Parasite	0	0	0	0	1	21	0	0
<i>Giardia</i>	0	0	0	0	1	21	0	0
Chemical	0	0	0	0	0	0	0	0
Suspected propylene glycol	0	0	0	0	0	0	0	0
Multiple[§]	0	0	0	0	0	0	0	0
Unidentified	1	7	0	0	0	0	0	0
Total	4	201	2	0	1	21	0	0

*No reported outbreaks in public non-community water systems had surface water or unknown water source.

[†]Includes outbreaks with mixed water sources (i.e. ground water and surface water). Two giardiasis outbreaks were associated with mixed source community.

[§]One outbreak had multiple etiologic agent types. The etiologies were *Campylobacter* spp. (i.e., bacterium) and *Giardia intestinalis* (i.e., parasite).

