

GIARDIASIS

SUMMARY REPORT



2013

NATIONAL NOTIFIABLE DISEASES
SURVEILLANCE SYSTEM, UNITED STATES



Centers for Disease
Control and Prevention
National Center for Emerging and
Zoonotic Infectious Diseases

Suggested Citation

Centers for Disease Control and Prevention (CDC). Giardiasis Summary Report — National Notifiable Diseases Surveillance System, United States, 2013. Atlanta, Georgia: U.S. Department of Health and Human Services, CDC, 2019.

Report Compiled By

Michelle E. Gleason, MPH; Ariana Perez, MPH; Gabriella Veytsel, MPH; Sarah A. Collier, MPH; Zachary A. Marsh, MPH; Michele C. Hlavsa, RN, MPH; Jennifer R. Cope, MD, MPH; Kathleen E. Fullerton, MPH; Kathy Benedict, DVM, PhD

Centers for Disease Control and Prevention

National Center for Emerging and Zoonotic Infections
Division of Foodborne, Waterborne, and Environmental Diseases
Waterborne Disease Prevention Branch
1600 Clifton Road, Atlanta, GA 30329
E-mail: healthywater@cdc.gov

Data are presented for cases of giardiasis for the year 2013 reported to CDC through October 9, 2018.

Findings and conclusions from this report do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Contents

Background.....	4
Surveillance Overview: National Giardiasis Case Surveillance	4
Methods.....	4
Case Definition	4
Analysis	4
Acknowledgements.....	5
References.....	6
Tables and Figures.....	7
Figure 1. Incidence of giardiasis cases, by year and case classification — National Notifiable Diseases Surveillance System, United States, 1995–2013 (n = 389,713).....	7
Table 1. Number, percentage, and incidence of giardiasis cases, by region and jurisdiction — National Notifiable Diseases Surveillance System, United States, 2013 (n = 15,267).....	8
Figure 2. Incidence of giardiasis cases, by reporting jurisdiction — National Notifiable Diseases Surveillance System, United States, 2013 (n=15,267)	10
Table 2. Number and percentage of giardiasis cases, by selected patient demographic characteristics — National Notifiable Diseases Surveillance System, United States, 2013 (n = 15,267).....	11
Figure 3. Incidence of giardiasis cases, by age group — National Notifiable Diseases Surveillance System, United States, 2013 (n = 14,882).....	12
Figure 4. Incidence of giardiasis cases, by sex and age group — National Notifiable Diseases Surveillance System, United States, 2013 (n = 14,807)	13
Figure 5. Number of giardiasis cases, by date of symptom onset — National Notifiable Diseases Surveillance System, United States, 2013 (n = 7,936)	14
Supplemental Table	15
Table S1. Giardiasis reporting and non-reporting states (gray shading) — National Notifiable Diseases Surveillance System, United States, 1995–2013	15

Background

Surveillance Overview: National Giardiasis Case Surveillance

Giardiasis is an illness caused by the protozoan parasite *Giardia duodenalis* (formerly called *G. lamblia* or *G. intestinalis*), which causes gastrointestinal symptoms such as diarrhea, abdominal cramps, bloating, weight loss, or malabsorption (1, 2). Each year in the United States, it is estimated that *Giardia* causes more than 1.2 million illnesses (3).

Giardiasis is a [nationally notifiable disease](#); the first full year of reporting was 1993. National data are collected through passive surveillance. Healthcare providers and laboratories that diagnose confirmed giardiasis cases report to the local or state health departments. State and territorial health departments, the District of Columbia (DC), and the New York City health departments, in turn, voluntarily notify CDC of cases via the [National Notifiable Disease Surveillance System \(NNSS\)](#). The number of health departments submitting can vary from year to year depending on which states have designated giardiasis as reportable in their jurisdictions.

State, DC, US territory, and Freely Associated State public health agencies voluntarily notify CDC of giardiasis outbreaks via the [National Outbreak Reporting System \(NORS\)](#). NORS data are not presented here; however, [summaries of data on waterborne disease outbreaks](#) are published elsewhere.

Methods

Case Definition

The [definition](#) of a confirmed case of giardiasis has changed over time; the [first national case definition](#) was published in 1997 (4), and [a revised case definition](#) was published in 2011 (5). The current (2011) case definition differs from the 1997 definition in clarifying that clinical symptoms are necessary for categorizing giardiasis cases as confirmed.

A confirmed case of giardiasis is defined as a case that meets the clinical description and the criteria for laboratory confirmation. Laboratory-confirmed giardiasis is defined as the detection of *Giardia* organisms, antigen, or DNA in stool, intestinal fluid, tissue samples, biopsy specimens, or other biological samples (5). Nonconfirmed cases of giardiasis include probable, suspected, and unknown cases. A probable case of giardiasis meets the clinical description and is epidemiologically linked to a confirmed case. A national case definition for suspected cases of giardiasis does not exist; the definition varies by state. Cases not classified as confirmed, probable, or suspect are classified as unknown.

Analysis

National giardiasis surveillance data for 2013 were analyzed using R version 3.5.1 and SAS 9.4. Data cleaning processes included case deduplication and the verification of case status (e.g., confirmed, nonconfirmed). Numbers, percentages, and incidence (cases per 100,000 population) of giardiasis were calculated in aggregate for the United States and separately for each reporting jurisdiction. Rates were calculated by dividing the number of giardiasis cases by each year's mid-year census estimates (6, 7) and multiplying by 100,000. Region and total population estimates included only jurisdictions that reported (Supplemental Table 1). In addition to analyzing data nationally and by reporting jurisdiction, data were analyzed by region (Northeast, Midwest, South, and West regions), as defined by the U.S. Census Bureau (8). To account for differences in the seasonal use of recreational water, the West region was further subdivided into Northwest and Southwest.

To examine reporting over time, giardiasis rates were calculated by year (1995 to 2013) and case status (confirmed or nonconfirmed). Average annual giardiasis rates were calculated by demographic variables (e.g., age and sex) and jurisdiction. Rates were not calculated for race, ethnicity, or month of onset due to large proportion of missing data for these variables (i.e., 39.1%, 44.9%, and 48.0%, respectively). Cases reported by territories were excluded from the analysis, because detailed demographic census data are not available to calculate rates by age and sex.

Acknowledgements


The authors gratefully acknowledge Julia Painter and Jonathan Yoder for their assistance in summarizing the annual reporting of giardiasis by state through previous work on the Domestic Epidemiology Team, Waterborne Disease Prevention Branch, CDC. This report is based on contributions by state and local epidemiologists and microbiologists.

References

1. Huang DB, White AC. An updated review on *Cryptosporidium* and *Giardia*. *Gastroenterol Clin North Am*. 2006;35(2):291-314, viii. DOI: <https://www.sciencedirect.com/science/article/pii/S0889855306000252?via%3Dihub>.
2. Eberhard M, Gabrielli A, Montresor A, et al. Giardiasis (*Giardia enteritis*). In: Control of communicable diseases manual, 20th Edition. Heymann DL, Ed. Washington, DC; 2014:234-36.
3. Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 2011;17:7–15. DOI: [10.3201/eid1701.P11101](https://doi.org/10.3201/eid1701.P11101).
4. CDC. Giardiasis: 1997 case definition. Atlanta, GA: US Department of Health and Human Services, CDC; 1997. Available at <http://wwwn.cdc.gov/nndss/script/casedef.aspx?CondYrID=683&DatePub=1997-01-01>.
5. CDC. Giardiasis: 2011 case definition. Atlanta, GA: US Department of Health and Human Services, CDC; 2011. Available at <http://wwwn.cdc.gov/nndss/script/casedef.aspx?CondYrID=685&DatePub=1/1/2011>.
6. United States Department of Health and Human Services (US DHHS), Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Bridged-Race Population Estimates, United States July 1st resident population by state, county, age, sex, bridged-race, and Hispanic origin. Compiled from 1990-1999 bridged-race intercensal population estimates (released by NCHS on 7/26/2004); revised bridged-race 2000-2009 intercensal population estimates (released by NCHS on 10/26/2012); and bridged-race Vintage 2017 (2010-2017) postcensal population estimates (released by NCHS on 6/27/2018). Available on CDC WONDER Online Database. Accessed at <http://wonder.cdc.gov/bridged-race-v2017.html> on Apr 18, 2019.
7. US Census Bureau. Annual Estimates of the Resident Population for Incorporated Places of 50,000 or More, Ranked by July 1, 2017 Population: April 1, 2010 to July 1, 2017. Washington, DC: US Census Bureau; Release Date: May 2018. Available at <https://www.census.gov/content/census/en/data/tables/time-series/demo/pepstat/2010s-total-cities-and-towns.html>.
8. US Census Bureau. Census regions and divisions of the United States. Washington, DC: US Census Bureau. Available at https://www2.census.gov/geo/pdfs/maps-data/maps/reference/us_regdiv.pdf.

Tables and Figures

Figure 1. Incidence* of giardiasis cases, by year and case classification — National Notifiable Diseases Surveillance System, United States, 1995–2013 (n = 389,713) - Download raw data.



* Cases per 100,000 population per year

[§] Probable, suspect, and unknown cases

Since 2011, the incidence of reported, confirmed giardiasis cases has remained < 7.0 cases per 100,000 population. In 2013, there were 15,267 reported giardiasis cases in the United States (97.6% confirmed and 2.4% nonconfirmed).

Table 1. Number, percentage*, and incidence[§] of giardiasis cases, by region and jurisdiction — National Notifiable Diseases Surveillance System, United States, 2013 (n = 15,267)

Region/Jurisdiction	No.	%	Incidence	No. of outbreak associated cases
Northeast	4,421	29	7.9	100
Connecticut	230	1.5	6.4	
Maine	218	1.4	16.4	
Massachusetts	664	4.3	9.9	3
New Hampshire	117	0.8	8.8	
New Jersey	336	2.2	3.8	
New York City [¶]	764	5.0	9.0	
New York State [¶]	1,113	7.3	9.9	59
Pennsylvania	765	5.0	6.0	38
Rhode Island	41	0.3	3.9	
Vermont	173	1.1	27.6	
Midwest	3,551	23.3	5.3	35
Illinois	311	2.0	2.4	18
Indiana	229	1.5	3.5	2
Iowa	273	1.8	8.8	
Kansas	102	0.7	3.5	1
Michigan	547	3.6	5.5	
Minnesota	622	4.1	11.5	
Missouri	244	1.6	4.0	
Nebraska	169	1.1	9.0	
North Dakota	44	0.3	6.1	
Ohio	513	3.4	4.4	14
South Dakota	111	0.7	13.2	
Wisconsin	386	2.5	6.7	
South	3,097	20.3	4.8	96
Alabama	174	1.1	3.6	
Arkansas	119	0.8	4.0	
Delaware	20	0.1	2.2	
District of Columbia	84	0.6	12.9	
Florida	1,114	7.3	5.7	96
Georgia	641	4.2	6.4	
Kentucky	NR			
Louisiana	255	1.7	5.5	
Maryland	228	1.5	3.8	
Mississippi	NR			
North Carolina	NR			
Oklahoma	NR			
South Carolina	133	0.9	2.8	
Tennessee	NR			
Texas	NR			
Virginia	279	1.8	3.4	
West Virginia	50	0.3	2.7	
Northwest	1,256	8.2	8.5	7
Alaska	82	0.5	11.1	
Idaho	139	0.9	8.6	6
Montana	91	0.6	9.0	

Oregon	364	2.4	9.3	1
Washington	548	3.6	7.9	
Wyoming	32	0.2	5.5	
Southwest	2,942	19.3	5.0	5
Arizona	115	0.8	1.7	
California	1,991	13.0	5.2	
Colorado	355	2.3	6.7	5
Hawaii	60	0.4	4.3	
Nevada	91	0.6	3.3	
New Mexico	102	0.7	4.9	
Utah	228	1.5	7.9	
Total	15,267	100	5.8	243

Abbreviation NR = Not Reportable


* Percentages might not total 100% because of rounding

§ Cases per 100,000 population

¶ New York State and New York City data are mutually exclusive

By jurisdiction, giardiasis incidence ranged from 1.7 per 100,000 population in Arizona to 27.6 per 100,000 population in Vermont. By region, incidence of reported giardiasis cases ranged from 4.8 cases per 100,000 population in the South to 8.5 in the Northwest. Differences in incidence might reflect differences in risk factors or mode of transmission of *Giardia*; the magnitude of outbreaks; or the capacity or requirements to detect, investigate, and report cases.

Figure 2. Incidence* of giardiasis cases, by reporting jurisdiction — National Notifiable Diseases Surveillance System, United States, 2013 (n=15,267[§]) - Download raw data.



* Cases per 100,000 population

[§] Non-reporting states included Kentucky, Mississippi, North Carolina, Oklahoma, Tennessee, and Texas

[¶] New York State and New York City data are mutually exclusive

Giardiasis is geographically widespread across the United States. Although incidence rates appear to be consistently higher in the northern states, differences in incidence might reflect differences in risk factors or modes of transmission of *Giardia*; the magnitude of outbreaks; or the capacity or requirements to detect, investigate, and report cases.


Table 2. Number and percentage* of giardiasis cases, by selected patient demographic characteristics — National Notifiable Diseases Surveillance System, United States, 2013 (n = 15,267)

Characteristic	No.	%
Sex		
Male	9,125	59.8
Female	6,061	39.7
Missing	81	0.5
Race		
American Indian/Alaska Native	52	0.3
Asian/Pacific Islander	698	4.6
Black	1,112	7.3
White	6,888	45.1
Other	545	3.6
Missing	5,972	39.1
Ethnicity		
Hispanic	1,064	7
Non-Hispanic	7,344	48.1
Missing	6,859	44.9
Total	15,267	100.0

* Percentages might not total 100% because of rounding

During 2013, a total of 9,125 patients were male (59.8%) and 6,061 (39.7%) were female; 81 (0.5%) were missing data on sex. The majority of cases for whom data on race were available occurred among the classifications white (74.1%), black (12.0%), and Asian/Pacific Islander (7.5%). Data on race were not included for 39.1% of total annual case reports, and data on ethnicity were missing for 44.9% of case reports. The majority of patients for whom data on ethnicity were available were non-Hispanic (87.3%).

Figure 3. Incidence* of giardiasis cases, by age group — National Notifiable Diseases Surveillance System, United States, 2013 (n = 14,882[§]) - Download raw data.




* Cases per 100,000 population

[§] Age data missing for 385 patients

In 2013, the incidence of reported giardiasis cases was highest among patients aged 1-4 years, 5-9 years, and 30-34 years (Incidence = 13.7, 7.6, and 6.0 cases per 100,000 population, respectively).

Figure 4. Incidence* of giardiasis cases, by sex and age group — National Notifiable Diseases Surveillance System, United States, 2013 (n = 14,807[§]) - Download raw data.




* Cases per 100,000 population

[§] Age or sex data missing for 460 patients

Rates were highest among males in every age group. The difference was most pronounced between sexes aged 45-49 years.

Figure 5. Number of giardiasis cases, by date of symptom onset — National Notifiable Diseases Surveillance System, United States, 2013 (n = 7,936*) - Download raw data.



* Date of symptom onset data missing for 7,331 patients

In 2013, the majority of cases by symptom onset occurred between July and October, with a peak in July (n=1,029).

Supplemental Table

Table S1. Giardiasis reporting and non-reporting states (gray shading) — National Notifiable Diseases Surveillance System, United States, 1995–2013

State	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
AK																				
AL																				
AR																				
AZ																				
CA																				
CO																				
CT																				
DE																				
FL																				
GA																				
HI																				
IA																				
ID																				
IL																				
IN																				
KS																				
KY																				
LA																				
MA																				
MD																				
ME																				
MI																				
MN																				
MO																				
MS																				
MT																				
NC																				
ND																				
NE																				
NH																				
NJ																				
NM																				
NV																				
NY																				
OH																				
OK																				
OR																				
PA																				
RI																				
SC																				
SD																				
TN																				
TX																				
UT																				
VA																				
VT																				
WA																				
WI																				
WV																				
WY																				