

Number of Waterborne Disease Outbreaks Associated with Recreational Water, by Predominant Illness* and Type of Water Exposure — Waterborne Disease and Outbreak Surveillance System, 2011–2012

Predominant Illness	Type of Water Exposure															
	Treated				Untreated				Total							
	Outbreaks	Cases	Hospitalized	Died	Outbreaks	Cases	Hospitalized	Died	Outbreaks	%	Cases	%	Hospitalized	%	Died	%
AGI	44	1,064	54	0	15	374	20	0	59	65.6%	1,438	80.4%	74	77.9%	0	0.0%
ARI	13	97	19	1	0	0	0	0	13	14.4%	97	5.4%	19	20.0%	1	100.0%
ARI and Eye	1	3	0	0	0	0	0	0	1	1.1%	3	0.2%	0	0.0%	0	0.0%
ARI, Eye, and Skin	1	11	0	0	0	0	0	0	1	1.1%	11	0.6%	0	0.0%	0	0.0%
Eye	0	0	0	0	2	40	1	0	2	2.2%	40	2.2%	1	1.1%	0	0.0%
Eye and Skin	2	46	0	0	0	0	0	0	2	2.2%	46	2.6%	0	0.0%	0	0.0%
Skin	8	88	0	0	4	65	1	0	12	13.3%	153	8.6%	1	1.1%	0	0.0%
Total (%)	69 (76.7%)	1,309 (73.2%)	73 (76.8%)	1 (100.0%)	21 (23.3%)	479 (26.8%)	22 (23.2%)	0 (0.0%)	90	100.0%	1788	100.0%	95	100.0%	1	100.0%

Abbreviations: AGI = acute gastrointestinal illness; ARI = acute respiratory illness; Eye = illnesses, conditions, or symptoms related to the eyes; Skin = illnesses, conditions, or symptoms - related to the skin. -

*n=90; The category of illness reported by 50% of ill respondents. All legionellosis outbreaks were categorized as ARI. -