

<b>Adolescent Behaviors and Experiences Survey</b>				
	<b>Female</b>		<b>Male</b>	
<b>Mental Health</b>	<b>Percentage</b>	<b>Confidence Interval</b>	<b>Percentage</b>	<b>Confidence Interval</b>
Who felt sad or hopeless (almost every day for two or more weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	56.5	53.4 - 59.5	31.4	29.1 - 33.7
Seriously considered attempting suicide (during the 12 months before the survey)	26.0	23.4 - 28.6	13.6	12.0 - 15.4
Made a plan about how they would attempt suicide (during the 12 months before the survey)	20.5	18.0 - 23.2	10.0	8.6 - 11.5
Actually attempted suicide (one or more times during the 12 months before the survey)	12.4	10.5 - 14.5	5.3	4.2 - 6.6
Had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.6	1.8 - 3.6	1.2	0.8 - 1.8
Reported that their mental health was most of the time or always not good (including stress, anxiety, and depression, during the 30 days before the survey)	41.6	38.4 - 44.9	19.6	17.6 - 21.8