

This slide set contains information about and data from the School Health Profiles or Profiles. It includes state data from 2014.

What is the School Health Profiles (Profiles)?

- Profiles is a system of surveys assessing school health policies and practices in states, territories, and large urban school districts
- Profiles surveys are conducted biennially (during even-numbered years) by state, territorial, and local education and health agencies with technical assistance from CDC
- Profiles questionnaires are administered to middle and high school principals and lead health education teachers

What is the School Health Profiles (Profiles)?

- Profiles is a system of surveys assessing school health policies and practices in states, territories, and large urban school districts
- Profiles surveys are conducted biennially (during even-numbered years) by state, territorial, and local education and health agencies with technical assistance from CDC
- Profiles questionnaires are administered to middle and high school principals and lead health education teachers

Topics Monitored by Profiles

- School health education requirements and content
- Physical education and physical activity
- Practices related to bullying and sexual harassment
- School health policies related to tobacco-use prevention and nutrition
- School-based health services
- Family engagement and community involvement
- School health coordination

Topics Monitored by Profiles

-
-
-
-
-
-
-


Profiles Participation, 1998 – 2014

	1998	2000	2002	2004	2006	2008	2010	2012	2014
# of states	38	38	43	41	44	50	50	49	50
# of large urban school districts	10	13	14	13	16	21	20	19	21
# of territories	--	--	--	--	--	5	6	5	3
# of tribal governments	--	--	--	--	--	2	2	2	--
Total # of sites	48	51	57	54	60	78	78	75	74
Total % of sites with weighted data	77%	75%	77%	70%	82%	94%	99%	92%	96%

This slide contains state, large urban school district, territory, and tribal government Profiles participation data from 1998 through 2012.

- In 1998, 38 states and 10 cities (including Washington, D.C.) participated in Profiles, for a total number of 48 states and cities. The percent of sites with weighted data in 1998 was 77%.
- In 2000, 38 states and 13 cities (including Washington, D.C.) participated in Profiles, for a total number of 51 states and cities. The percent of sites with weighted data in 2000 was 75%.
- In 2002, 43 states and 14 cities (including Washington, D.C.) participated in Profiles, for a total number of 57 states and cities. The percent of sites with weighted data in 2002 was 77%.
- In 2004, 41 states and 13 cities (including Washington, D.C.) participated in Profiles, for a total number of 54 states and cities. The percent of sites with weighted data in 2004 was 70%.
- In 2006, 44 states and 16 cities (including Washington, D.C.) participated in Profiles, for a total number of 60 states and cities. The percent of sites with weighted data in 2006 was 82%.
- In 2008, 50 states, 21 cities (including Washington, D.C.), 5 territories, and 2 tribes participated in Profiles, for a total number of 78 sites. The percent of sites with weighted data in 2008 was 94%.
- In 2010, 50 states, 20 cities (including Washington, D.C.), 6 territories, and 2 tribes participated in Profiles, for a total number of 78 sites. The percent of sites with weighted data in 2010 was 99%.
- In 2012, 49 states, 19 cities (including Washington, D.C.), 5 territories, and 2 tribes participated in Profiles, for a total number of 75 sites. The percent of sites with weighted data in 2012 was 92%.
- In 2014, 50 states, 21 cities (including Washington, D.C.), and 3 territories participated in Profiles, for a total number of 74 sites. The percent of sites with weighted data in 2014 was 96%.

State and District Participation, Profiles 2014



This map illustrates states and large urban school districts that participated in the School Health Profiles in 2014. States and large urban school districts with weighted and unweighted results are shown.

States with weighted results are: Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming

States with unweighted results are: Louisiana and New Mexico

Surveyed large urban school districts with weighted results are: Baltimore, MD; Boston, MA; Broward County, FL; Chicago, IL; Cleveland, OH; DeKalb County, GA; Detroit, MI; District of Columbia; Duval County, FL; Fort Worth, TX; Houston, TX; Los Angeles, CA; Miami-Dade County, FL; New York City, NY; Oakland, CA; Orange County, FL; Palm Beach County, FL; Philadelphia, PA; San Diego, CA; San Francisco, CA; Shelby County, TN

Weighted data means that at least 70% of the principals or lead health education teachers in the sample completed the survey. Weighted data represent the state, large urban school district, territory, or tribal government. With weighted data, it is possible to say, for example, "X% of schools in our state offer students intramural activities or physical activity clubs."

Unweighted data represent only the schools that completed the questionnaire.

Uses of Profiles Data

Education and health officials use Profiles data to

- Describe school health policies and practices and compare them across jurisdictions
- Identify professional development needs
- Plan and monitor programs
- Support health-related policies and legislation
- Seek funding
- Garner support for future surveys

Uses of Profiles Data

Education and health officials use Profiles data to

- Describe school health policies and practices and compare them across jurisdictions
- Identify professional development needs
- Plan and monitor programs
- Support health-related policies and legislation
- Seek funding
- Garner support for future surveys

Examples of Uses of Profiles Data

- **Montana**
 - Used Profiles results to develop customized technical assistance plans for selected schools to increase attention towards nutrition physical activity, and employee wellness
- **Florida**
 - Used Profiles results to develop the agenda for their Healthy Schools Summer Academy professional development event
- **Alaska**
 - Use Profiles data to help write a grant to support rural school health services

Examples of Uses of Profiles Data

Montana used Profiles results to develop customized technical assistance plans for selected schools to increase attention towards nutrition physical activity, and employee wellness.

Florida used Profiles results to develop the agenda for their Healthy Schools Summer Academy professional development event.

Alaska used Profiles data to help write a grant to support rural school health services.

Profiles Information

<http://www.cdc.gov/schoolhealthprofiles>

- Overview of the School Health Profiles
- Questionnaires and item rationales
- Comprehensive results
- Fact sheet
- Publications and journal articles
- Participation history
- Frequently asked questions about requesting data files
- Data file request form


For more information about Profiles go to <http://www.cdc.gov/schoolhealthprofiles>.

This web site contains


- Overview of the School Health Profiles
- Questionnaires and item rationales
- Comprehensive results
- Fact sheet
- Publications and journal articles
- Participation history
- Frequently asked questions about requesting data files
- Data file request form

HIV, STD, and Pregnancy Prevention


HIV, STD, and Pregnancy Prevention




This slide shows the percentage of secondary schools that taught 11 key HIV, STD, and pregnancy prevention topics in a required course during grades 6, 7, or 8 and during grades 9, 10, 11, or 12. The values range from 16% to 69%. Alaska, Arizona, Nebraska, North Dakota, South Dakota, range from 0% to 24%. Alabama, Arkansas, California, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, Ohio, Oregon, Pennsylvania, South Carolina, Tennessee, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. Delaware, Maine, Maryland, New Hampshire, New Jersey, New York, North Carolina, Rhode Island, West Virginia, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that assessed the ability of students to do 7 skills in a required course during grades 6, 7, or 8 and during grades 9, 10, 11, or 12. The values range from 22% to 80%. Arizona, range from 0% to 24%. Alaska, California, Iowa, Kansas, Massachusetts, Missouri, Nebraska, North Dakota, Ohio, Pennsylvania, South Dakota, Tennessee, range from 25% to 49%. Alabama, Arkansas, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kentucky, Maine, Michigan, Minnesota, Mississippi, Montana, Nevada, New Hampshire, New Jersey, Oregon, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Maryland, New York, North Carolina, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools in which those who teach sexual health education are provided with key materials for teaching sexual health education. The values range from 29% to 73%. Alaska, Arizona, Arkansas, Delaware, Hawaii, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Minnesota, Montana, Nebraska, North Dakota, Ohio, Oregon, South Dakota, Utah, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Alabama, California, Connecticut, Florida, Georgia, Idaho, Maryland, Massachusetts, Michigan, Mississippi, Missouri, Nevada, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on 10 key HIV prevention topics. The values range from 0% to 13%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that taught about 7 contraceptives in a required course during grades 9, 10, 11, or 12. The values range from 16% to 87%. Alabama, Arizona, Arkansas, Georgia, Indiana, North Dakota, South Dakota, range from 0% to 24%. Alaska, Florida, Idaho, Illinois, Iowa, Kansas, Kentucky, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, Ohio, South Carolina, Tennessee, Virginia, Wyoming, range from 25% to 49%. California, Connecticut, Hawaii, Maine, Maryland, Massachusetts, North Carolina, Oregon, Pennsylvania, Rhode Island, Washington, West Virginia, Wisconsin, range from 50% to 74%. Delaware, New Hampshire, New Jersey, New York, Vermont, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that tried to increase student knowledge on HIV prevention in a required course. The values range from 40% to 97%. Arizona, range from 25% to 49%. Alaska, Tennessee, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on human sexuality in a required course. The values range from 36% to 96%. Arizona, range from 25% to 49%. Alabama, Alaska, Arkansas, California, Florida, Kentucky, Mississippi, Missouri, Ohio, South Dakota, Tennessee, range from 50% to 74%. Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Maryland, Massachusetts, Michigan, Minnesota, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on pregnancy prevention in a required course. The values range from 36% to 94%. Arizona, range from 25% to 49%. Alabama, Alaska, California, Connecticut, Florida, South Dakota, Tennessee, range from 50% to 74%. Arkansas, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on STD prevention in a required course. The values range from 39% to 96%. Arizona, range from 25% to 49%. Alaska, South Dakota, Tennessee, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that taught how HIV and other STDs are transmitted in a required course. The values range from 38% to 97%. Arizona, range from 25% to 49%. Alaska, South Dakota, Tennessee, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught health consequences of HIV, other STDs, and pregnancy in a required course. The values range from 38% to 97%. Arizona, range from 25% to 49%. Alaska, South Dakota, Tennessee, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught the benefits of being sexually abstinent in a required course. The values range from 37% to 98%. Arizona, range from 25% to 49%. Alaska, South Dakota, Tennessee, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught how to access valid and reliable health information, products, and services related to HIV, other STDs, and pregnancy in a required course. The values range from 31% to 92%. Arizona, South Dakota, range from 25% to 49%. Alabama, Alaska, California, Connecticut, Florida, Iowa, Kansas, Kentucky, Massachusetts, Nebraska, Ohio, Tennessee, range from 50% to 74%. Arkansas, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught the influences of family, peers, media, technology, and other factors on sexual risk behaviors in a required course. The values range from 35% to 96%. Arizona, range from 25% to 49%. Alaska, California, Kentucky, South Dakota, Tennessee, range from 50% to 74%. Alabama, Arkansas, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught communication and negotiation skills related to eliminating or reducing risk for HIV, other STDs, and pregnancy in a required course. The values range from 33% to 94%. Arizona, South Dakota, range from 25% to 49%. Alabama, Alaska, California, Connecticut, Florida, Kentucky, Massachusetts, Nebraska, Tennessee, range from 50% to 74%. Arkansas, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught goal-setting and decision-making skills related to eliminating or reducing risk for HIV, other STDs, and pregnancy in a required course. The values range from 33% to 94%. Arizona, South Dakota, range from 25% to 49%. Alabama, Alaska, California, Florida, Iowa, Kansas, Kentucky, Massachusetts, Nebraska, Tennessee, range from 50% to 74%. Arkansas, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught influencing and supporting others to avoid or reduce sexual risk behaviors in a required course. The values range from 34% to 94%. Arizona, South Dakota, range from 25% to 49%. Alabama, Alaska, California, Connecticut, Florida, Iowa, Kentucky, Massachusetts, Nebraska, Tennessee, range from 50% to 74%. Arkansas, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught efficacy of condoms in a required course. The values range from 27% to 87%. Arizona, South Dakota, Tennessee, range from 25% to 49%. Alabama, Alaska, Arkansas, California, Connecticut, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, North Dakota, Ohio, Pennsylvania, South Carolina, Virginia, West Virginia, Wyoming, range from 50% to 74%. Delaware, Hawaii, Maine, Maryland, New Hampshire, New Jersey, New York, North Carolina, Oregon, Rhode Island, Vermont, Washington, Wisconsin, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught the importance of using condoms consistently and correctly in a required course. The values range from 23% to 83%. Arizona, range from 0% to 24%. Alabama, Georgia, Indiana, Kansas, Kentucky, Montana, Nebraska, South Dakota, Tennessee, range from 25% to 49%. Alaska, Arkansas, California, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Iowa, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nevada, New Hampshire, New Jersey, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Maine, New York, North Carolina, Vermont, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught how to obtain condoms in a required course. The values range from 17% to 75%. Arizona, South Dakota, range from 0% to 24%. Alabama, Alaska, California, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Michigan, Mississippi, Missouri, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, South Carolina, Tennessee, Virginia, Wyoming, range from 25% to 49%. Arkansas, Connecticut, Delaware, Hawaii, Iowa, Maine, Maryland, Massachusetts, Minnesota, Nevada, New Hampshire, New Jersey, North Carolina, Oregon, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, range from 50% to 74%. New York, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught how to correctly use a condom in a required course. The values range from 15% to 67%. Arizona, Georgia, South Dakota, range from 0% to 24%. Alabama, Alaska, Arkansas, California, Connecticut, Florida, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, North Dakota, Ohio, Pennsylvania, South Carolina, Tennessee, Virginia, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Delaware, Hawaii, Maine, Massachusetts, New Hampshire, New Jersey, New York, North Carolina, Oregon, Rhode Island, Vermont, Washington, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught the importance of using a condom at the same time as another form of contraception to prevent both STDs and pregnancy in a required course. The values range from 23% to 80%. Arizona, range from 0% to 24%. Georgia, Indiana, Kentucky, South Dakota, Tennessee, range from 25% to 49%. Alabama, Alaska, Arkansas, California, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Iowa, Kansas, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Maine, New York, North Carolina, Vermont, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught how to create and sustain healthy and respectful relationships in a required course. The values range from 37% to 96%. Arizona, range from 25% to 49%. Alaska, California, Florida, South Dakota, Tennessee, range from 50% to 74%. Alabama, Arkansas, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught the importance of limiting the number of sexual partners in a required course. The values range from 30% to 91%. Arizona, South Dakota, range from 25% to 49%. Alabama, Alaska, California, Connecticut, Florida, Georgia, Kansas, Kentucky, Massachusetts, Missouri, Montana, Nebraska, South Carolina, Tennessee, Virginia, Wyoming, range from 50% to 74%. Arkansas, Delaware, Hawaii, Idaho, Illinois, Indiana, Iowa, Maine, Maryland, Michigan, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that taught preventive care that is necessary to maintain reproductive and sexual health in a required course. The values range from 29% to 92%. Arizona, South Dakota, range from 25% to 49%. Alabama, Alaska, California, Connecticut, Florida, Georgia, Hawaii, Illinois, Iowa, Kansas, Kentucky, Maine, Massachusetts, Michigan, Mississippi, Missouri, Montana, Nebraska, Ohio, Pennsylvania, Tennessee, Virginia, Wyoming, range from 50% to 74%. Arkansas, Delaware, Idaho, Indiana, Maryland, Minnesota, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oregon, Rhode Island, South Carolina, Vermont, Washington, West Virginia, Wisconsin, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools with a policy on students or staff who have HIV infection or AIDS that addresses attendance of students with HIV infection, procedures to protect HIV-infected students and staff from discrimination, and maintaining confidentiality of HIV-infected students and staff. The values range from 21% to 84%. Kentucky, range from 0% to 24%. Alaska, Arizona, Arkansas, Colorado, Delaware, Georgia, Hawaii, Iowa, Maryland, Michigan, Minnesota, Mississippi, North Carolina, Ohio, South Dakota, Texas, range from 25% to 49%. Alabama, California, Connecticut, Florida, Idaho, Illinois, Indiana, Kansas, Maine, Massachusetts, Missouri, Montana, Nebraska, Nevada, New Jersey, New York, North Dakota, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. New Hampshire, Oregon, Vermont, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on HIV prevention. The values range from 5% to 53%. Alaska, Arizona, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, West Virginia, Wyoming, range from 0% to 24%. Alabama, Arkansas, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Maine, Massachusetts, Mississippi, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, South Carolina, Tennessee, Utah, Vermont, Virginia, Washington, Wisconsin, range from 25% to 49%. Maryland, Michigan, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.



This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on human sexuality. The values range from 6% to 62%. Alabama, Alaska, Arizona, Arkansas, California, Illinois, Indiana, Iowa, Kansas, Kentucky, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, West Virginia, Wyoming, range from 0% to 24%. Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Maine, Maryland, Massachusetts, Michigan, Mississippi, Montana, New Jersey, New York, North Carolina, South Carolina, Vermont, Virginia, Washington, Wisconsin, range from 25% to 49%. Nevada, New Hampshire, Utah, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on pregnancy prevention. The values range from 5% to 42%. Alabama, Alaska, Arizona, Arkansas, California, Connecticut, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Minnesota, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Delaware, Florida, Hawaii, Maine, Maryland, Massachusetts, Michigan, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, South Carolina, Utah, range from 25% to 49%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on STD prevention. The values range from 5% to 48%. Alabama, Alaska, Arizona, Arkansas, Illinois, Iowa, Kansas, Kentucky, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Virginia, West Virginia, Wyoming, range from 0% to 24%. California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Indiana, Maine, Maryland, Massachusetts, Michigan, Mississippi, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, South Carolina, Tennessee, Utah, Vermont, Washington, Wisconsin, range from 25% to 49%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




Practices Related to Safe and Supportive Environments




This slide shows the percentage of secondary schools that provide curricula or supplementary materials that include HIV, STD, or pregnancy prevention information that is relevant to LGBTQ youth . The values range from 11% to 56%. Alabama, Alaska, Arizona, Arkansas, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, North Dakota, Ohio, South Carolina, South Dakota, Tennessee, Virginia, Wyoming, range from 0% to 24%. California, Connecticut, Delaware, Florida, Hawaii, Iowa, Maine, Maryland, Massachusetts, Nevada, New Hampshire, New Jersey, North Carolina, Oregon, Pennsylvania, Rhode Island, Washington, West Virginia, Wisconsin, range from 25% to 49%. New York, Vermont, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools with a gay/straight alliance or similar club. The values range from 13% to 56%. Alabama, Alaska, Arizona, Arkansas, Georgia, Idaho, Illinois, Kansas, Kentucky, Mississippi, Missouri, Montana, Nebraska, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, West Virginia, Wyoming, range from 0% to 24%. California, Colorado, Connecticut, Delaware, Florida, Hawaii, Indiana, Iowa, Maryland, Michigan, Minnesota, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Texas, Vermont, Virginia, Washington, Wisconsin, range from 25% to 49%. Maine, Massachusetts, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that implement parent engagement strategies for all students. The values range from 25% to 66%. Alabama, Alaska, Arizona, Colorado, Hawaii, Idaho, Illinois, Kansas, Missouri, Montana, Nebraska, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Dakota, Washington, Wyoming, range from 25% to 49%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, New Hampshire, New Jersey, New York, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, West Virginia, Wisconsin, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that implement school connectedness strategies . The values range from 61% to 93%. Alaska, Arizona, California, Hawaii, Maine, Montana, North Dakota, Ohio, Oklahoma, South Dakota, Wyoming, range from 50% to 74%. Alabama, Arkansas, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that prevent bullying and sexual harassment, including electronic aggression, among all students. The values range from 27% to 76%. Alabama, Alaska, Arizona, Colorado, Georgia, Hawaii, Idaho, Iowa, Kansas, Maine, Michigan, Minnesota, Missouri, Montana, Nebraska, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Dakota, Wisconsin, Wyoming, range from 25% to 49%. Arkansas, California, Connecticut, Delaware, Florida, Illinois, Indiana, Kentucky, Maryland, Massachusetts, Mississippi, Nevada, New Hampshire, New York, North Dakota, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, range from 50% to 74%. New Jersey, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that provide curricula or supplementary materials and engage in 5 practices related to LGBTQ youth. The values range from 2% to 27%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Massachusetts, New Hampshire, New York, Vermont, range from 25% to 49%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that identify “safe spaces” where LGBTQ youth can receive support from administrators, teachers, or other school staff. The values range from 37% to 85%. Alaska, Arizona, Mississippi, Nebraska, North Dakota, South Dakota, Wyoming, range from 25% to 49%. Alabama, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Michigan, Minnesota, Missouri, Montana, Nevada, New Hampshire, New Jersey, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Massachusetts, New York, Rhode Island, range from 75% to 100%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that prohibit harassment based on a student's perceived or actual sexual orientation or gender identity. The values range from 73% to 97%. Mississippi, South Dakota, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.



This slide shows the percentage of secondary schools that encourage staff to attend professional development on safe and supportive school environments for all students, regardless of sexual orientation or gender identity. The values range from 39% to 82%. Alaska, Arizona, Georgia, Kansas, Mississippi, North Dakota, South Dakota, range from 25% to 49%. Alabama, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Massachusetts, New Hampshire, New Jersey, New York, Vermont, range from 75% to 100%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that facilitate access to providers not on school property who have experience in providing health services to LGBTQ youth. The values range from 30% to 69%. Alabama, Alaska, Arizona, Arkansas, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Virginia, Wisconsin, Wyoming, range from 25% to 49%. California, Colorado, Connecticut, Iowa, Maine, Massachusetts, Minnesota, Nevada, New Hampshire, New Jersey, New York, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that facilitate access to providers not on school property who have experience in providing social and psychological services to LGBTQ youth. The values range from 30% to 73%. Alabama, Alaska, Arizona, Arkansas, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, North Dakota, Ohio, South Carolina, South Dakota, Tennessee, Texas, Virginia, range from 25% to 49%. California, Colorado, Connecticut, Delaware, Florida, Iowa, Maine, Maryland, Massachusetts, Minnesota, New Hampshire, New Jersey, New York, North Carolina, Oklahoma, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




School Health Services




This slide shows the percentage of secondary schools that have a full-time registered nurse who provides health services to students . The values range from 5% to 99%. Alaska, California, Hawaii, Idaho, Michigan, Montana, North Dakota, Oklahoma, Oregon, South Dakota, Utah, Washington, Wisconsin, range from 0% to 24%. Arizona, Colorado, Kansas, Kentucky, Mississippi, Nebraska, Nevada, North Carolina, Ohio, West Virginia, Wyoming, range from 25% to 49%. Florida, Georgia, Illinois, Indiana, Iowa, Maine, Minnesota, Tennessee, range from 50% to 74%. Alabama, Arkansas, Connecticut, Delaware, Maryland, Massachusetts, Missouri, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Texas, Vermont, Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that provide students with on-site services or referrals to healthcare providers for 7 sexual health services. The values range from 17% to 52%. Arizona, Georgia, Kansas, Mississippi, Nebraska, South Carolina, range from 0% to 24%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maine, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide HIV testing to students. The values range from 0% to 22%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide HIV treatment to students. The values range from 0% to 9%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide STD testing to students. The values range from 0% to 24%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide STD treatment to students. The values range from 0% to 19%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide pregnancy testing to students. The values range from 0% to 24%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide condoms to students. The values range from 0% to 30%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. New York, range from 25% to 49%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide condom-compatible lubricants to students. The values range from 0% to 19%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide contraceptives other than condoms to students. The values range from 0% to 13%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.



This slide shows the percentage of secondary schools that provide prenatal care to students. The values range from 1% to 12%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide HPV vaccine administration to students. The values range from 0% to 21%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.




This slide shows the percentage of secondary schools that provide students with referrals* for HIV testing. The values range from 27% to 68%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Virginia, range from 25% to 49%. Delaware, Indiana, Iowa, Maine, Massachusetts, Minnesota, New Hampshire, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for HIV treatment. The values range from 26% to 66%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, range from 25% to 49%. Delaware, Iowa, Maine, Massachusetts, Minnesota, New Hampshire, New York, Oregon, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for STD testing. The values range from 27% to 69%. Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Virginia, range from 25% to 49%. California, Delaware, Indiana, Iowa, Maine, Massachusetts, Minnesota, New Hampshire, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for STD treatment. The values range from 26% to 67%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Virginia, range from 25% to 49%. Delaware, Iowa, Maine, Massachusetts, Minnesota, New Hampshire, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for pregnancy testing. The values range from 30% to 71%. Alabama, Alaska, Arizona, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Nebraska, Nevada, New Jersey, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Virginia, range from 25% to 49%. Arkansas, California, Delaware, Indiana, Iowa, Maine, Massachusetts, Minnesota, Montana, New Hampshire, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for condoms. The values range from 19% to 56%. Arizona, Georgia, Kansas, Mississippi, Nebraska, South Carolina, range from 0% to 24%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maine, New York, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for condom-compatible lubricants. The values range from 18% to 53%. Arizona, Georgia, Kansas, Mississippi, Nebraska, South Carolina, range from 0% to 24%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 25% to 49%. Maine, New York, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for contraceptives other than condoms. The values range from 17% to 56%. Arizona, Georgia, Mississippi, Nebraska, South Carolina, range from 0% to 24%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nevada, New Hampshire, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, Wisconsin, range from 25% to 49%. Maine, New York, West Virginia, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for prenatal care. The values range from 26% to 69%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, range from 25% to 49%. Delaware, Indiana, Iowa, Maine, Massachusetts, Minnesota, New Hampshire, New York, North Carolina, Oregon, Pennsylvania, Rhode Island, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that provide students with referrals* for HPV vaccine administration. The values range from 24% to 63%. Arizona, Georgia, range from 0% to 24%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Idaho, Illinois, Indiana, Kansas, Kentucky, Maryland, Michigan, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Wisconsin, range from 25% to 49%. Iowa, Maine, Massachusetts, Minnesota, New Hampshire, New York, Oregon, Rhode Island, Washington, West Virginia, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results. Utah did not ask this question.

*To any organizations or health care professionals not on school property.




This slide shows the percentage of secondary schools that identify and track students with chronic conditions* that may require daily or emergency management. The values range from 81% to 100%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.


*For example, asthma or food allergies




This slide shows the percentage of secondary schools that have protocols that ensure students with a chronic condition that may require daily or emergency management are enrolled into private, state, or federally funded insurance programs if eligible. The values range from 46% to 84%. Alaska, North Dakota, South Dakota, range from 25% to 49%. Alabama, Arizona, Arkansas, California, Colorado, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New York, North Carolina, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Connecticut, Delaware, Maryland, Massachusetts, New Hampshire, New Jersey, Pennsylvania, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that provide referrals to community-based medical care providers for students identified with chronic conditions or at risk for activity, diet, and weight-related chronic conditions. The values range from 34% to 100%. Alabama, Arizona, Florida, Georgia, Hawaii, Idaho, Kansas, Michigan, Mississippi, North Dakota, Oklahoma, Tennessee, range from 25% to 49%. Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Utah, Vermont, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




School Health Coordination




This slide shows the percentage of secondary schools that currently have someone who oversees or coordinates school health and safety programs and activities . The values range from 61% to 96%. Alaska, Arizona, North Dakota, South Dakota, range from 50% to 74%. Alabama, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that had a school health council, committee, or team that offers guidance on the development of policies or coordinates activities on health topics . The values range from 29% to 76%. Alabama, Alaska, Arizona, California, Georgia, Idaho, Illinois, Nevada, North Dakota, Ohio, Oregon, South Dakota, Utah, Washington, range from 25% to 49%. Arkansas, Colorado, Connecticut, Delaware, Florida, Hawaii, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. New Hampshire, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that have a school health council, committee, or team with representation from at least 6 of 8 groups. The values range from 10% to 64%. Alaska, Arizona, California, Hawaii, Idaho, Maryland, Nevada, Ohio, Oregon, Utah, Washington, range from 0% to 24%. Alabama, Colorado, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Jersey, New York, North Carolina, North Dakota, Pennsylvania, Rhode Island, South Carolina, South Dakota, Vermont, Virginia, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Arkansas, New Hampshire, Oklahoma, Tennessee, Texas, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools that used the School Health Index or other self-assessment tool to assess their policies, activities, and programs in HIV, STD, and teen pregnancy prevention; physical activity; nutrition; and tobacco-use prevention. The values range from 17% to 53%. Alaska, Arizona, Colorado, Connecticut, Delaware, Idaho, Iowa, Kansas, Maryland, Nevada, Ohio, Pennsylvania, South Dakota, Utah, Virginia, Washington, Wyoming, range from 0% to 24%. Alabama, California, Florida, Georgia, Hawaii, Illinois, Indiana, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Oklahoma, Oregon, Rhode Island, South Carolina, Texas, Vermont, West Virginia, Wisconsin, range from 25% to 49%. Arkansas, Tennessee, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that include at least 1 health and safety objective in their school improvement plan and have completed a self-assessment of school health policies and practices and have reviewed health and safety data during the past year as part of their school improvement planning process. The values range from 10% to 53%. Alabama, Alaska, Arizona, Colorado, Connecticut, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Minnesota, Missouri, Nebraska, Nevada, New Hampshire, New Jersey, New York, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Utah, Virginia, Wyoming, range from 0% to 24%. California, Delaware, Florida, Georgia, Kentucky, Maryland, Massachusetts, Michigan, Mississippi, Montana, North Carolina, North Dakota, Oklahoma, South Carolina, Tennessee, Texas, Vermont, Washington, West Virginia, Wisconsin, range from 25% to 49%. Arkansas, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that have a school health team that performs 5 actions to help plan and implement school health programs. The values range from 6% to 32%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Massachusetts, New Hampshire, Tennessee, Vermont, range from 25% to 49%. Louisiana and New Mexico did not have weighted results.




Health Education




This slide shows the percentage of secondary schools with a health education curriculum that addresses all 8 national standards for health education. The values range from 31% to 87%. Arizona, California, Delaware, Hawaii, Iowa, Ohio, range from 25% to 49%. Alabama, Alaska, Connecticut, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, New York, North Dakota, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Arkansas, Maryland, Mississippi, New Hampshire, New Jersey, North Carolina, Rhode Island, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.



This slide shows the percentage of secondary schools that required students to take 2 or more health education courses. The values range from 15% to 91%. Alabama, Arizona, California, Florida, Nevada, range from 0% to 24%. Alaska, Georgia, Hawaii, Iowa, Kansas, Kentucky, Michigan, Mississippi, Nebraska, North Carolina, Ohio, South Carolina, South Dakota, Tennessee, Utah, Vermont, Washington, range from 25% to 49%. Arkansas, Connecticut, Delaware, Idaho, Illinois, Indiana, Maine, Maryland, Massachusetts, Missouri, New Hampshire, New York, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Minnesota, Montana, New Jersey, North Dakota, Oregon, Pennsylvania, Rhode Island, Virginia, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the pmong secondary schools that required a health education course, percentage that required students who fail the course to repeat it. The values range from 38% to 81%. California, Connecticut, Massachusetts, New Jersey, Rhode Island, South Carolina, Utah, Virginia, range from 25% to 49%. Arizona, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New York, North Carolina, North Dakota, Oregon, Pennsylvania, Tennessee, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Alabama, Alaska, Arkansas, Hawaii, Idaho, Ohio, South Dakota, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.



This slide shows the percentage of secondary schools in which the lead health education teacher had professional preparation in health education or in health and physical education combined. The values range from 15% to 87%. Alaska, Arizona, California, range from 0% to 24%. Alabama, Florida, Hawaii, Illinois, Iowa, Maine, Maryland, Massachusetts, Mississippi, Missouri, Nebraska, New Hampshire, New York, Oregon, South Carolina, Utah, Vermont, Washington, range from 25% to 49%. Arkansas, Connecticut, Georgia, Idaho, Indiana, Kansas, Kentucky, Michigan, Montana, Nevada, New Jersey, North Carolina, North Dakota, Ohio, Rhode Island, South Dakota, Tennessee, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Delaware, Minnesota, Pennsylvania, Virginia, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.


Tobacco-Use Prevention

Tobacco-Use Prevention




This slide shows the percentage of secondary schools that prohibited all tobacco use at all times in all locations*. The values range from 34% to 77%. Delaware, Idaho, Kansas, Kentucky, Missouri, Nebraska, Ohio, South Dakota, range from 25% to 49%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Illinois, Indiana, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Montana, Nevada, New Hampshire, New Jersey, New York, North Dakota, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. North Carolina, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.


*Prohibited the use of all tobacco, including cigarettes, smokeless tobacco, cigars, and pipes, by students, faculty and school staff, and visitors, in school buildings, outside on school grounds, on school buses or other vehicles used to transport students, and at off-campus, school-sponsored events, during school hours and non-school hours.




This slide shows the percentage of secondary schools that taught 18 key tobacco-use prevention topics in a required course. The values range from 19% to 72%. Arizona, range from 0% to 24%. Alaska, California, Connecticut, Delaware, Florida, Hawaii, Illinois, Iowa, Kansas, Maine, Massachusetts, Michigan, Minnesota, Missouri, Nebraska, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Utah, Vermont, Washington, Wisconsin, Wyoming, range from 25% to 49%. Alabama, Arkansas, Georgia, Idaho, Indiana, Kentucky, Maryland, Mississippi, Montana, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, South Dakota, Virginia, West Virginia, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that provided tobacco cessation services for students, faculty, and staff at school or through arrangements with providers not on school property. The values range from 9% to 55%. Alabama, Arizona, Connecticut, Georgia, Hawaii, Idaho, Illinois, Iowa, Kansas, Kentucky, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New York, Ohio, Oklahoma, South Dakota, Tennessee, Texas, range from 0% to 24%. Alaska, Arkansas, California, Colorado, Delaware, Florida, Indiana, Maine, Maryland, Montana, New Hampshire, North Carolina, North Dakota, Oregon, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. West Virginia, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on tobacco-use prevention. The values range from 10% to 48%. Alabama, Arizona, California, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Massachusetts, Minnesota, Missouri, Nebraska, New Jersey, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Washington, Wyoming, range from 0% to 24%. Alaska, Arkansas, Florida, Kentucky, Maryland, Michigan, Mississippi, Montana, Nevada, New Hampshire, New York, North Carolina, Tennessee, Utah, Vermont, Virginia, West Virginia, Wisconsin, range from 25% to 49%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.



This slide shows the percentage of secondary schools that posted signs marking a tobacco-free school zone. The values range from 42% to 97%. Connecticut, range from 25% to 49%. Hawaii, Idaho, Illinois, Kentucky, Maryland, Massachusetts, Michigan, Missouri, Nebraska, Nevada, New Jersey, Ohio, Pennsylvania, Rhode Island, South Carolina, Utah, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Delaware, Florida, Georgia, Indiana, Iowa, Kansas, Maine, Minnesota, Mississippi, Montana, New Hampshire, New York, North Carolina, North Dakota, Oklahoma, Oregon, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




Nutrition




This slide shows the percentage of secondary schools that did not sell less healthy foods and beverages* in vending machines or at the school store, canteen, or snack bar. The values range from 19% to 88%. Idaho, Montana, Nevada, South Carolina, Utah, range from 0% to 24%. Alabama, California, Colorado, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Michigan, Minnesota, Missouri, Nebraska, New Jersey, New York, North Carolina, North Dakota, Oklahoma, Pennsylvania, South Dakota, Virginia, Washington, Wisconsin, Wyoming, range from 25% to 49%. Alaska, Arizona, Arkansas, Delaware, Maine, Mississippi, New Hampshire, Ohio, Oregon, Rhode Island, Tennessee, Texas, Vermont, range from 50% to 74%. Connecticut, Hawaii, Massachusetts, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.


*Baked goods not low in fat, salty snacks not low in fat, chocolate candy, other kinds of candy, soda pop or fruit drinks that are not 100% juice, and sports drinks



This slide shows the percentage of secondary schools in which students could not purchase chocolate candy from vending machines or at the school store, canteen, or snack bar. The values range from 34% to 99%. Utah, range from 25% to 49%. Georgia, Idaho, Indiana, Michigan, Minnesota, Missouri, Montana, Nebraska, North Carolina, Oklahoma, South Carolina, Washington, Wyoming, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Illinois, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase other kinds of candy from vending machines or at the school store, canteen, or snack bar. The values range from 30% to 99%. Idaho, South Carolina, Utah, range from 25% to 49%. Georgia, Illinois, Indiana, Kansas, Kentucky, Maryland, Michigan, Minnesota, Missouri, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Iowa, Maine, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New York, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, Tennessee, Texas, Vermont, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase salty snacks that are not low in fat* from vending machines or at the school store, canteen, or snack bar. The values range from 39% to 95%. Utah, range from 25% to 49%. Alaska, Arizona, Colorado, Florida, Georgia, Idaho, Illinois, Indiana, Kentucky, Maryland, Michigan, Minnesota, Missouri, Montana, Nebraska, New Jersey, North Carolina, Oklahoma, Pennsylvania, South Carolina, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Alabama, Arkansas, California, Connecticut, Delaware, Hawaii, Iowa, Kansas, Maine, Massachusetts, Mississippi, Nevada, New Hampshire, New York, North Dakota, Ohio, Oregon, Rhode Island, South Dakota, Tennessee, Texas, Vermont, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.


*Such as regular potato chips.




This slide shows the percentage of secondary schools in which students could not purchase cookies, crackers, cakes, pastries, or other baked goods that are not low in fat from vending machines or at the school store, canteen, or snack bar. The values range from 43% to 98%. Utah, range from 25% to 49%. Arizona, Colorado, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Maryland, Michigan, Minnesota, Missouri, Montana, Nebraska, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, South Carolina, South Dakota, Virginia, Washington, Wisconsin, Wyoming, range from 50% to 74%. Alabama, Alaska, Arkansas, California, Connecticut, Delaware, Hawaii, Iowa, Kentucky, Maine, Massachusetts, Mississippi, Nevada, New Hampshire, North Dakota, Ohio, Oregon, Rhode Island, Tennessee, Texas, Vermont, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools in which students could not purchase 2% or whole milk from vending machines or at the school store, canteen, or snack bar. The values range from 63% to 96%. California, Florida, Georgia, Idaho, Illinois, Indiana, Maryland, Michigan, Minnesota, Missouri, Nevada, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Texas, Utah, Virginia, Washington, Wisconsin, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Delaware, Hawaii, Iowa, Kansas, Kentucky, Maine, Massachusetts, Mississippi, Montana, Nebraska, North Dakota, Ohio, Oklahoma, Oregon, South Dakota, Tennessee, Vermont, West Virginia, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase soda pop or fruit drinks that are not 100% juice from vending machines or at the school store, canteen, or snack bar. The values range from 50% to 97%. Idaho, range from 25% to 49%. Alabama, Alaska, Georgia, Illinois, Indiana, Kentucky, Maryland, Michigan, Minnesota, Missouri, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, South Carolina, South Dakota, Utah, Virginia, Washington, Wyoming, range from 50% to 74%. Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Iowa, Kansas, Maine, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New York, Ohio, Oregon, Pennsylvania, Rhode Island, Tennessee, Texas, Vermont, West Virginia, Wisconsin, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase sports drinks* from vending machines or at the school store, canteen, or snack bar. The values range from 29% to 92%. Alabama, Georgia, Idaho, Indiana, Kansas, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, North Dakota, South Carolina, South Dakota, Utah, Washington, Wisconsin, Wyoming, range from 25% to 49%. Alaska, Arizona, California, Colorado, Delaware, Florida, Illinois, Iowa, Kentucky, Maine, Maryland, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Tennessee, Texas, Vermont, Virginia, range from 50% to 74%. Arkansas, Connecticut, Hawaii, Massachusetts, Rhode Island, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Such as Gatorade.




This slide shows the percentage of secondary schools in which students could not purchase energy drinks* from vending machines or at the school store, canteen, or snack bar. The values range from 93% to 100%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.


*Such as Red Bull or Monster.



This slide shows the percentage of secondary schools in which students could not purchase foods or beverages containing caffeine from vending machines or at the school store, canteen, or snack bar. The values range from 50% to 98%. Idaho, range from 25% to 49%. Georgia, Illinois, Indiana, Kentucky, Michigan, Minnesota, Missouri, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wyoming, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Hawaii, Iowa, Kansas, Maine, Maryland, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New York, Ohio, Oregon, Pennsylvania, Rhode Island, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that allowed students to purchase fruits and vegetables from vending machines or at the school store, canteen, or snack bar. The values range from 1% to 39%. Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Delaware, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Massachusetts, Mississippi, Missouri, Montana, Nebraska, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. California, Florida, Maryland, Michigan, Minnesota, New Hampshire, New Jersey, New York, Rhode Island, range from 25% to 49%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that prohibited all forms of advertising and promotion of candy, fast food restaurants, or soft drinks in all locations*. The values range from 27% to 78%. Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Missouri, Montana, Nebraska, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Virginia, Wisconsin, Wyoming, range from 25% to 49%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Hawaii, Maryland, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New York, Oregon, Texas, Utah, Vermont, Washington, West Virginia, range from 50% to 74%. Maine, Massachusetts, Rhode Island, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.


*In school buildings; on school grounds, including on the outside of the school building, on playing fields, or other areas of the campus; on school buses or other vehicles used to transport students; and in school publications.




This slide shows the percentage of secondary schools that priced nutritious foods and beverages at a lower cost while increasing the price of less nutritious foods and beverages. The values range from 4% to 26%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Delaware, range from 25% to 49%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that provided information to students or families on the nutrition and caloric content of foods available. The values range from 30% to 69%. Alabama, Alaska, Arizona, Arkansas, Georgia, Hawaii, Idaho, Kansas, Kentucky, Mississippi, Nebraska, Nevada, North Carolina, North Dakota, Oklahoma, Oregon, South Dakota, Utah, Washington, Wyoming, range from 25% to 49%. California, Colorado, Connecticut, Delaware, Florida, Illinois, Indiana, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that placed fruits and vegetables near the cafeteria cashier, where they are easy to access. The values range from 48% to 90%. Alaska, Hawaii, Nevada, range from 25% to 49%. Alabama, Arizona, Arkansas, Idaho, Illinois, Iowa, Kansas, Mississippi, Missouri, Montana, Nebraska, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Wisconsin, Wyoming, range from 50% to 74%. California, Colorado, Connecticut, Delaware, Florida, Georgia, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Oregon, Pennsylvania, Rhode Island, Texas, Utah, Vermont, Virginia, Washington, West Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that allowed students to have access to drinking water. The values range from 42% to 88%. Connecticut, Delaware, Maine, Pennsylvania, Rhode Island, Vermont, range from 25% to 49%. Alabama, Alaska, Arkansas, Colorado, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, South Carolina, South Dakota, Tennessee, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Arizona, California, Florida, Hawaii, Montana, Nebraska, Nevada, Oklahoma, Texas, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that offer fruits or non-fried vegetables when foods or beverages are offered at school celebrations. The values range from 17% to 59%. Illinois, Indiana, Iowa, Kansas, Kentucky, Nebraska, North Dakota, South Dakota, range from 0% to 24%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Jersey, New York, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. Maine, New Hampshire, Rhode Island, Vermont, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that prohibited school staff from giving students food or food coupons as a reward for good behavior or good academic performance. The values range from 10% to 72%. Alaska, Colorado, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Missouri, Montana, Nebraska, North Dakota, Ohio, Oregon, South Carolina, South Dakota, Utah, Virginia, Washington, Wisconsin, Wyoming, range from 0% to 24%. Alabama, Arizona, Arkansas, California, Connecticut, Hawaii, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, Rhode Island, Tennessee, Texas, Vermont, range from 25% to 49%. West Virginia, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools that prohibited less nutritious foods and beverages from being sold for fundraising purposes. The values range from 11% to 64%. Alaska, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Montana, Nebraska, South Carolina, South Dakota, Utah, Washington, Wyoming, range from 0% to 24%. Alabama, Arizona, Arkansas, California, Colorado, Delaware, Florida, Maine, Maryland, Minnesota, Mississippi, Missouri, Nevada, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Tennessee, Texas, Vermont, Virginia, Wisconsin, range from 25% to 49%. Connecticut, Hawaii, Massachusetts, New Hampshire, Rhode Island, West Virginia, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on nutrition and dietary behavior. The values range from 14% to 71%. Alaska, California, Georgia, Hawaii, Indiana, Oregon, range from 0% to 24%. Alabama, Arizona, Arkansas, Connecticut, Delaware, Florida, Idaho, Illinois, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, New York, North Carolina, North Dakota, Ohio, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 25% to 49%. New Hampshire, range from 50% to 74%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.




This slide shows the percentage of secondary schools that taught 20 key nutrition and dietary behavior topics in a required course. The values range from 28% to 79%. Alaska, Arizona, California, Connecticut, Delaware, Hawaii, Iowa, Maine, Massachusetts, Michigan, Ohio, Vermont, Washington, range from 25% to 49%. Alabama, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Kentucky, Maryland, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Dakota, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Virginia, Wisconsin, Wyoming, range from 50% to 74%. Arkansas, North Carolina, West Virginia, range from 75% to 100%. Colorado, Louisiana, New Mexico, Oklahoma, and Texas did not have weighted results.

Physical Education And Physical Activity

Physical Education and Physical Activity




This slide shows the percentage of secondary schools that have established, implemented, or evaluated a Comprehensive School Physical Activity Program. The values range from 0% to 10%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 0% to 24%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 6th grade*. The values range from 72% to 100%. Michigan, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 7th grade*. The values range from 47% to 100%. Nevada, range from 25% to 49%. Michigan, Oklahoma, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 8th grade*. The values range from 58% to 100%. Arizona, Hawaii, Kentucky, Michigan, Oklahoma, range from 50% to 74%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Maine, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 9th grade*. The values range from 28% to 100%. Oklahoma, range from 25% to 49%. Arizona, Idaho, range from 50% to 74%. Alabama, Alaska, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 10th grade*. The values range from 15% to 100%. Kansas, Kentucky, Oklahoma, range from 0% to 24%. Arizona, Georgia, Michigan, Nebraska, North Carolina, South Carolina, South Dakota, Wyoming, range from 25% to 49%. Alabama, Alaska, Colorado, Florida, Hawaii, Idaho, Indiana, Maryland, Minnesota, Missouri, New Hampshire, North Dakota, Ohio, Oregon, West Virginia, range from 50% to 74%. Arkansas, California, Connecticut, Delaware, Illinois, Iowa, Maine, Massachusetts, Mississippi, Montana, Nevada, New Jersey, New York, Pennsylvania, Rhode Island, Tennessee, Texas, Utah, Vermont, Virginia, Washington, Wisconsin, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 11th grade*. The values range from 9% to 100%. Arizona, Delaware, Kansas, Kentucky, Minnesota, Montana, North Carolina, Oklahoma, South Dakota, Virginia, range from 0% to 24%. Alabama, California, Georgia, Hawaii, Idaho, Indiana, Maine, Maryland, Michigan, Nebraska, Nevada, New Hampshire, North Dakota, Ohio, Oregon, South Carolina, West Virginia, Wyoming, range from 25% to 49%. Alaska, Colorado, Connecticut, Florida, Missouri, Tennessee, Texas, Utah, Vermont, Washington, range from 50% to 74%. Arkansas, Illinois, Iowa, Massachusetts, Mississippi, New Jersey, New York, Pennsylvania, Rhode Island, Wisconsin, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which a required physical education course is taught in 12th grade*. The values range from 10% to 100%. Arizona, Kansas, Kentucky, Minnesota, Montana, North Carolina, North Dakota, Oklahoma, South Dakota, Virginia, range from 0% to 24%. Alabama, California, Colorado, Delaware, Georgia, Hawaii, Idaho, Indiana, Maine, Maryland, Michigan, Nebraska, Nevada, New Hampshire, Ohio, Oregon, South Carolina, Utah, West Virginia, Wyoming, range from 25% to 49%. Alaska, Connecticut, Florida, Massachusetts, Missouri, Tennessee, Texas, Vermont, Washington, Wisconsin, range from 50% to 74%. Arkansas, Illinois, Iowa, Mississippi, New Jersey, New York, Pennsylvania, Rhode Island, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*Among schools with students in that grade.




This slide shows the percentage of secondary schools in which at least 1 physical education teacher or specialist received professional development on physical education*. The values range from 38% to 97%. Alaska, range from 25% to 49%. Arizona, California, Hawaii, Idaho, Indiana, Iowa, Oklahoma, Oregon, South Dakota, range from 50% to 74%. Alabama, Arkansas, Colorado, Connecticut, Delaware, Florida, Georgia, Illinois, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.

*During the year before the survey.




This slide shows the percentage of secondary schools in which students participate in physical activity breaks in classrooms during the school day*. The values range from 24% to 72%. Alabama, range from 0% to 24%. Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Utah, Virginia, Washington, Wisconsin, range from 25% to 49%. Alaska, Arizona, Colorado, Hawaii, Maine, New Hampshire, Tennessee, Vermont, West Virginia, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.


*Outside of physical education.



This slide shows the percentage of secondary schools that offered opportunities for all students to participate in intramural sports programs or physical activity clubs. The values range from 31% to 86%. Arkansas, Iowa, Kansas, Nebraska, North Dakota, South Dakota, Wyoming, range from 25% to 49%. Alabama, Arizona, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kentucky, Maine, Michigan, Minnesota, Mississippi, Missouri, Montana, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Utah, Virginia, Washington, West Virginia, Wisconsin, range from 50% to 74%. Alaska, California, Colorado, Connecticut, Hawaii, Maryland, Massachusetts, Nevada, New Hampshire, New Jersey, New York, Vermont, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that offer interscholastic sports to students. The values range from 72% to 97%. Hawaii, Maryland, range from 50% to 74%. Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools that offer opportunities for students to participate in physical activity before the school day*. The values range from 19% to 66%. Delaware, Kentucky, Mississippi, Ohio, Virginia, range from 0% to 24%. Alabama, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Maine, Maryland, Massachusetts, Michigan, Missouri, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Vermont, Washington, West Virginia, range from 25% to 49%. Alaska, Arizona, Hawaii, Iowa, Minnesota, Montana, Nebraska, North Dakota, Texas, Utah, Wisconsin, Wyoming, range from 50% to 74%. Louisiana and New Mexico did not have weighted results.

*Through organized physical activities or access to facilities or equipment for physical activity.



This slide shows the percentage of secondary schools that have a joint use agreement for shared use of physical activity facilities. The values range from 43% to 88%. Arkansas, Ohio, Oklahoma, range from 25% to 49%. Alabama, Alaska, Arizona, California, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Dakota, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 50% to 74%. Colorado, Maryland, North Carolina, Utah, Virginia, range from 75% to 100%. Louisiana and New Mexico did not have weighted results.