

Youth Risk Behavior Surveillance System (YRBSS)



Interpretation of YRBS Trend Data

August 2020

Where can I get more information? Visit www.cdc.gov/yrbss or call 800-CDC-INFO (800-232-4636).

National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention
Division of Adolescent and School Health



Introduction

Purpose The purpose of this document is to provide background and to describe how to interpret the results of the trend analyses conducted for the YRBS Trend Fact Sheets.

Contents

Introduction

Purpose

Contents

Other Resources

Summary

Why Percent Change Is Not A Good Test of Difference

Introduction

Two Scenarios

What Statistical Tests Provide An Appropriate Test Of Difference Or Trends Over Time

Introduction

Statistically Significant Linear Trend

Statistically Significant Linear and Quadratic Trends

How To Interpret The YRBS Trend Fact Sheets

Long-term Change

Change from 2017-2019

Why Public Health Significance Also Should Be Considered

Other Resources

Readers unfamiliar with YRBS data should review:

- the latest [national data documentation](#),
 - [Methodology of the Youth Risk Behavior Surveillance System—2013](#),
 - [Software for Analysis of YRBS Data](#),
 - [Combining YRBS Data Across Years and Sites](#),
 - [Conducting Trend Analyses of YRBS Data](#), and
 - the [Frequently Asked Questions](#) page on the YRBS website.
-

Summary

The Youth Risk Behavior Survey (YRBS) trend fact sheets provide information about whether the percentage of high school students who are engaging in specific health risk behaviors is increasing, decreasing, or staying the same over time. YRBS data are collected

Interpretation of YRBS Trend Data

from a sample of the population of students in each jurisdiction. Consequently, it is not possible to determine whether the percentage of students who have engaged in a particular risk behavior has actually changed over time by simply looking at the prevalence estimates for each year on the trend fact sheets. The best way to determine whether an actual change has occurred over time is to conduct a statistical test that takes into account the YRBS sample design, changes in population characteristics over time, and all possible data points in each trend line.

The Centers for Disease Control and Prevention (CDC) has conducted appropriate statistical tests to determine whether changes have actually occurred in the prevalence of specific health risk behaviors since 1991 and between 2017 and 2019. Any changes detected by these tests are indicated in the last two columns of the trend fact sheets. Statistical tests must be used to determine if the prevalence estimates are actually different. Unless a change is indicated in the final two columns of the fact sheets, do not assume a change has occurred. If you are interested in changes during a different time period than the ones presented in these fact sheets (e.g., 2005–2019), additional statistical tests can be conducted with our web-based dissemination tool, Youth Online:

<https://nccd.cdc.gov/youthonline/App/Default.aspx>.

Why Percent Change Is Not A Good Test Of Difference

Introduction It is not possible to determine if two prevalence estimates based on a sample of the population are different or represent an actual trend by calculating percent change. Percent change is strongly dependent on the prevalence estimate's magnitude. Therefore, using the percent change can be misleading, especially if the absolute difference in the prevalence estimates is not presented in conjunction with the percent change.

Two Scenarios For example, consider these two scenarios:

Scenario 1: 1% of students used XYZ drug in 2017 and 2% did so in 2019. The absolute difference during the 2-year period is 1% ($2\% - 1\% = 1\%$). The percent increase is 100% $\{[(2-1)/1] * 100\% = 100\%\}$.

Scenario 2: 47% of students used XYZ drug in 2017 and 48% did so in 2019. The absolute difference during the 2-year period is 1% ($48\% - 47\% = 1\%$). The percent increase is $\{[(48-47)/47] * 100\% = 2\%\}$.

In both scenarios, the absolute difference is 1%. However, the percent increase is 100% in Scenario 1 but only 2% in Scenario 2. Clearly, using percent change alone to describe differences can be very misleading if the magnitude of the prevalence estimates is not simultaneously considered. Further, percent change calculations do not take into consideration the variance associated with each prevalence estimate, or changes in population characteristics (e.g., differences in racial/ethnic distributions) over time. We recommend performing statistical tests to detect differences between prevalence estimates.

Statistical Tests to Examine Difference or Trends Over Time

Introduction To accurately determine if prevalence estimates based on a sample of the population are different, or represent a trend, it is not enough to compare only the prevalence estimates. One must also consider the variance of each prevalence estimate. For example, a simple and conservative test that considers the prevalence estimate and the variance involves examining 95% confidence intervals. A good rule of thumb is that if two confidence intervals do not overlap (or have values in common), then the prevalence estimates can be considered statistically different. A more sophisticated statistical test that CDC uses to analyze YRBS data is a t-test. A t-test is a more accurate way to compare two prevalence estimates than examining confidence intervals.

If there are more than two prevalence estimates (as occurs when the same behavior is measured at least three times), logistic regression analyses (or similar analyses) consider *all* data points (not just the starting and ending prevalence estimates) and simultaneously adjust for changes in underlying population characteristics such as sex, age, or race/ethnicity. These adjustments are necessary if many years of data are being examined. Logistic regression analysis is the most effective way to determine if a statistically significant change has occurred across three or more prevalence estimates.

CDC analyzes long-term trends in risk behaviors using logistic regression analyses that assess linear and quadratic trends and control for changes in sex, race/ethnicity, and grade in the population during the same time period. Statistically significant linear trends indicate an increase or decrease during the entire time period. Statistically significant quadratic trends indicate changes in direction of segments of data over time, and provide an inflection point (point at which a change in direction occurs). The following pictures illustrate how linear and quadratic trends are interpreted.

How To Interpret the YRBS Trend Fact Sheets

Long-term Change

The YRBS trend fact sheet column titled “Long-term Change” describes statistically significant linear and quadratic changes over time from 1991 (or the first year in which the data were collected) through 2019 based on logistic regression analyses for each selected behavior. If there is a statistically significant linear trend then this situation is described as “Increased, 1991–2019” or “Decreased, 1991–2019.” If there is a statistically significant quadratic trend, both parts of the quadratic trend are described. For example, “Decreased, 1991–1999 and No change, 1999–2019.” If there is no statistically significant linear or quadratic trend then this situation is described as “No change, 1991–2019.” If you are interested in whether a statistically significant linear or quadratic change occurred during a different period of time (e.g., 2003–2019) than is represented in the YRBS fact sheets, additional statistical tests must be conducted. It is NOT possible to determine if a statistically significant trend exists by simply looking at the prevalence estimates.

Change from 2017–2019

The YRBS trend fact sheet column titled “Change from 2017–2019” describes whether the 2017 and 2019 prevalence estimates for each selected behavior are statistically different based on t-test analyses. If there is a statistically significant difference, then this situation is described as “Increased” or “Decreased.” If there is not a statistically significant difference, then this situation is described as “No change.” If you are interested in whether any other two prevalence estimates (besides the 2017 and 2019 estimates) are different, a new statistical test must be conducted. It is NOT possible to determine if two estimates are statistically different by simply looking at them.
