

TABLES

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010

Site	Principal surveys		Teacher surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
STATE SURVEYS				
Alabama	246	72	243	71
Alaska	190	75	185	73
Arizona	307	79	297	76
Arkansas	233	78	230	77
California	386	78	378	77
Colorado	233	70	NA	NA
Connecticut	229	71	232	72
Delaware*	67	80	65	77
Florida	358	82	361	82
Georgia	259	72	257	71
Hawaii*	82	73	80	71
Idaho*	255	75	249	73
Indiana	306	73	299	71
Iowa	254	73	249	72
Kansas	276	75	267	73
Kentucky	266	75	264	74
Louisiana	231	74	225	72
Maine*	259	80	241	74
Maryland	241	73	233	71
Massachusetts*	613	80	625	82
Michigan	308	76	302	75
Minnesota	306	73	301	72
Mississippi	218	76	205	71
Missouri	310	79	302	77
Montana*	211	73	209	72
Nebraska	210	72	212	73
Nevada*	161	90	131	73
New Hampshire*	182	85	183	86
New Jersey	308	73	306	72
New Mexico	191	70	NA	NA
New York	694	75	677	73
North Carolina	358	84	338	79
North Dakota	173	72	171	71
Ohio	338	73	335	73
Oklahoma	327	77	315	74
Oregon	275	72	270	70
Pennsylvania	342	74	327	71
Rhode Island*	95	85	95	85
South Carolina*	335	74	342	76
South Dakota	188	71	189	72
Tennessee	288	78	285	78
Texas	398	78	384	75
Utah*	183	71	189	73
Vermont*	128	84	116	76
Virginia	359	86	359	86
Washington	231	76	217	72
West Virginia	156	78	147	74
Wisconsin	411	88	385	83
Wyoming*	131	86	115	75

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Principal surveys		Teacher surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
LOCAL SURVEYS				
Albuquerque*	37	86	32	74
Baltimore*	83	75	78	70
Broward County*	66	84	64	81
Charlotte-Mecklenberg County*	53	82	56	86
Chicago	104	86	98	81
Detroit*	89	99	85	94
District of Columbia*	36	95	34	89
Fresno*	25	96	21	81
Houston*	81	95	81	95
Los Angeles*	124	91	125	91
Memphis*	58	87	58	87
Miami-Dade County*	110	93	110	93
Milwaukee*	73	88	64	77
New York City	277	71	271	70
Newark*	51	89	45	79
Orange County*	46	82	43	77
Philadelphia	145	84	146	84
San Diego*	67	100	60	90
San Francisco*	33	80	38	93
TERRITORIAL SURVEYS				
American Samoa*	26	90	26	90
Guam*	13	100	13	100
Marshall Islands*	68	80	NA	NA
Northern Mariana Islands*	8	100	8	100
Palau*	11	100	11	100
TRIBAL SURVEYS				
Cherokee Nation*	115	76	108	71
Nez Perce*	7	100	7	100

NA=Data not available.

*Sample included a census of secondary schools.

TABLE 2. Percentage of Secondary Schools That Required Health Education Instruction in Any of Grades 6–12, the Percentage That Required Students to Take Only 1 Health Education Course or 2 or More Courses, and Among Schools That Required a Health Education Course, the Percentage That Required Students Who Fail Such a Course to Repeat It, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Required health education instruction	Required only 1 health education course	Required 2 or more health education courses	Required students who fail a required health education course to repeat it*
STATE SURVEYS				
Alabama	80.8	63.9	12.4	80.7
Alaska	84.3	40.2	44.7	84.8
Arizona	52.4	35.9	18.5	51.6
Arkansas	98.3	41.7	55.2	79.5
California	72.8	45.4	21.9	45.2
Connecticut	85.4	27.2	63.4	50.8
Delaware	95.2	46.1	52.3	55.5
Florida	73.4	45.6	22.5	63.3
Georgia	87.4	49.2	40.7	56.8
Hawaii	91.8	53.9	39.2	72.3
Idaho	98.1	44.9	53.8	70.4
Indiana	96.1	32.4	63.0	61.5
Iowa	85.8	32.0	49.3	57.1
Kansas	88.9	45.8	42.0	61.5
Kentucky	81.7	42.6	43.9	64.4
Louisiana	88.2	57.2	32.3	63.5
Maine	94.5	31.8	62.4	47.6
Maryland	89.0	43.8	48.1	51.0
Massachusetts	84.8	26.5	59.7	41.3
Michigan	84.5	49.4	38.9	68.3
Minnesota	91.4	18.7	74.2	64.3
Mississippi	86.3	64.2	22.9	73.7
Missouri	97.3	33.1	62.4	67.1
Montana	98.4	7.0	92.4	69.8
Nebraska	92.7	34.4	59.4	60.7
Nevada	92.9	72.3	22.8	63.7
New Hampshire	89.6	30.9	63.1	51.6
New Jersey	95.6	17.3	78.3	39.2
New York	94.4	35.9	59.9	67.5
North Carolina	89.6	41.7	49.2	50.1
North Dakota	96.8	20.8	75.4	65.8
Ohio	89.3	37.5	51.7	63.6
Oklahoma	39.2	31.0	16.1	21.2
Oregon	95.2	20.1	75.1	56.7
Pennsylvania	91.4	16.0	76.9	56.8
Rhode Island	96.8	17.5	78.9	51.2
South Carolina	80.1	38.1	43.4	36.1
South Dakota	85.1	36.9	33.6	69.6
Tennessee	72.5	43.6	30.8	58.0
Texas	83.6	62.4	22.2	74.8
Utah	95.1	49.3	47.4	49.8
Vermont	89.0	25.6	60.7	54.4
Virginia	92.8	13.5	78.7	42.7
Washington	90.3	37.6	52.5	60.0
West Virginia	98.7	32.2	67.8	48.8
Wisconsin	96.9	31.8	65.5	72.2
Wyoming	90.4	33.9	60.6	66.8
State Median	89.6	36.9	52.3	60.7
State Range	39.2-98.7	7.0-72.3	12.4-92.4	21.2-84.8

TABLE 2. Percentage of Secondary Schools That Required Health Education Instruction in Any of Grades 6–12, the Percentage That Required Students to Take Only 1 Health Education Course or 2 or More Courses, and Among Schools That Required a Health Education Course, the Percentage That Required Students Who Fail Such a Course to Repeat It, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Required health education instruction	Required only 1 health education course	Required 2 or more health education courses	Required students who fail a required health education course to repeat it*
LOCAL SURVEYS				
Albuquerque	96.8	80.6	19.4	36.8
Baltimore	69.6	53.2	15.5	54.7
Broward County	65.2	52.5	9.8	71.8
Charlotte-Mecklenberg County	93.2	40.0	56.6	48.4
Chicago	59.2	37.3	27.9	35.3
Detroit	61.6	55.7	10.7	52.3
District of Columbia	82.5	39.6	45.5	68.6
Fresno	68.0	48.2	14.5	45.7
Houston	90.0	64.8	30.2	47.1
Los Angeles	99.2	57.4	41.7	47.8
Memphis	67.2	50.5	27.5	59.8
Miami-Dade County	62.1	32.2	20.7	55.0
Milwaukee	85.8	44.9	37.7	38.4
New York City	87.9	64.1	26.3	56.6
Newark	81.1	32.9	51.3	38.1
Orange County	90.6	44.4	7.3	80.4
Philadelphia	70.7	36.3	32.5	49.1
San Diego	98.3	0.0	0.0	NA
San Francisco	94.6	25.4	66.4	45.3
Local Median	82.5	44.9	27.5	48.8
Local Median	59.2-99.2	0.0-80.6	0.0-66.4	35.3-80.4
TERRITORIAL SURVEYS				
American Samoa	100.0	15.4	84.6	91.3
Guam	100.0	91.7	8.3	33.3
Northern Mariana Islands	87.5	57.1	42.9	100.0
Palau	90.9	100.0	0.0	9.1
Territorial Median	95.5	74.4	25.6	62.3
Territorial Range	87.5-100.0	15.4-100.0	0.0-84.6	9.1-100.0
TRIBAL SURVEYS				
Cherokee Nation	40.3	28.3	11.0	23.8
Nez Perce	100.0	0.0	100.0	85.7

NA=Data not available.

*Among schools that required a health education course.

TABLE 3. Percentage of Secondary Schools That Taught a Required Health Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
STATE SURVEYS							
Alabama	29.1	32.2	29.5	30.2	74.7	13.3	12.4
Alaska	47.6	44.6	49.3	67.2	49.9	43.1	36.2
Arizona	30.7	34.7	34.7	23.4	31.2	13.4	14.4
Arkansas	50.5	80.9	54.8	89.6	70.0	65.4	64.5
California	27.8	52.3	32.1	40.7	20.0	8.1	8.8
Connecticut	71.6	74.2	81.3	67.2	65.6	48.0	41.8
Delaware	77.4	84.4	84.4	87.5	27.8	16.7	11.1
Florida	36.6	33.6	37.9	42.3	41.9	31.9	36.8
Georgia	75.2	74.8	77.3	77.4	16.9	10.9	11.3
Hawaii	64.8	76.2	32.8	47.0	77.5	11.8	11.8
Idaho	27.6	62.2	68.4	29.4	63.5	32.7	16.9
Indiana	68.2	79.1	80.3	49.9	73.5	10.7	10.7
Iowa	37.7	59.7	53.1	49.4	36.3	21.1	19.4
Kansas	48.0	54.2	48.7	76.2	17.2	7.4	6.4
Kentucky	57.3	69.1	59.8	68.3	30.9	6.7	7.8
Louisiana	56.0	62.7	61.4	66.9	47.4	34.9	34.0
Maine	78.8	84.0	82.5	57.1	67.2	14.2	11.0
Maryland	76.8	79.9	79.6	53.8	51.6	34.6	38.2
Massachusetts	68.8	72.2	72.3	63.1	56.3	29.6	24.5
Michigan	33.7	47.8	39.0	67.7	29.9	16.2	18.2
Minnesota	46.4	67.0	73.5	49.1	76.5	18.1	12.4
Mississippi	40.3	47.7	49.3	76.9	71.7	65.7	64.7
Missouri	63.1	81.0	81.9	76.0	40.7	24.9	25.4
Montana	75.7	93.1	92.9	95.6	92.0	8.6	5.5
Nebraska	58.8	69.3	69.4	67.4	35.7	8.7	13.3
Nevada	14.0	17.0	72.8	72.0	40.6	9.5	6.0
New Hampshire	69.6	80.9	80.5	67.2	64.7	30.4	21.1
New Jersey	88.6	91.0	91.0	87.3	90.8	96.0	99.0
New York	46.4	61.0	59.2	35.2	70.4	51.0	49.6
North Carolina	79.1	78.7	78.2	81.4	12.1	8.7	7.5
North Dakota	73.5	88.3	91.1	65.9	36.4	13.8	11.8
Ohio	34.9	49.1	49.8	63.0	43.5	9.2	8.1
Oklahoma	29.9	24.2	22.7	12.3	18.0	14.1	13.4
Oregon	70.4	81.9	77.5	61.6	64.2	46.9	27.9
Pennsylvania	66.3	66.2	66.0	60.0	60.9	39.2	23.9
Rhode Island	92.0	94.6	94.6	86.1	90.9	90.9	88.5
South Carolina	74.8	75.9	75.8	56.1	24.9	18.6	18.8
South Dakota	42.8	62.9	56.2	41.5	13.1	6.2	6.9
Tennessee	41.0	46.2	46.3	51.5	43.7	24.2	23.1
Texas	35.4	31.9	45.8	63.5	62.3	57.7	57.3
Utah	35.4	40.5	66.3	16.1	89.6	32.6	20.1
Vermont	57.2	67.1	65.6	51.3	53.1	22.3	22.3
Virginia	75.2	78.7	72.8	82.7	86.7	2.9	2.9
Washington	46.4	63.2	62.5	66.5	40.6	25.7	23.4
West Virginia	94.0	92.1	94.3	66.2	74.9	23.8	23.8
Wisconsin	62.9	64.0	65.0	65.6	39.2	15.5	5.7
Wyoming	59.4	77.5	78.4	76.2	55.6	21.5	17.6
State Median	57.3	67.1	66.3	65.6	51.6	21.1	18.2
State Range	14.0-94.0	17.0-94.6	22.7-94.6	12.3-95.6	12.1-92.0	2.9-96.0	2.9-99.0

TABLE 3. Percentage of Secondary Schools That Taught a Required Health Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
LOCAL SURVEYS							
Albuquerque	5.3	84.2	10.5	100.0	11.1	12.5	12.5
Baltimore	25.0	31.5	32.5	40.8	56.5	47.6	57.1
Broward County	33.8	32.7	32.7	45.9	65.2	63.6	72.7
Charlotte-Mecklenberg County	92.7	92.7	92.7	93.2	9.2	9.2	9.6
Chicago	46.6	48.3	49.8	34.1	21.2	14.3	14.3
Detroit	30.4	34.1	40.5	23.4	75.0	60.0	65.0
District of Columbia	68.4	63.2	66.7	39.3	100.0	23.1	23.1
Fresno	12.5	41.7	8.3	38.2	0.0	0.0	0.0
Houston	91.4	89.5	89.5	85.0	89.6	81.6	82.3
Los Angeles	54.5	100.0	4.7	96.0	12.2	14.3	14.3
Memphis	40.3	39.7	47.8	63.1	52.4	52.4	52.4
Miami-Dade County	30.6	32.4	28.6	13.3	52.6	22.9	36.4
Milwaukee	72.6	72.6	72.6	24.4	61.5	46.2	57.1
New York City	45.6	48.9	58.2	47.1	66.5	68.4	73.1
Newark	59.9	61.0	61.0	63.6	100.0	100.0	100.0
Orange County	19.2	23.1	15.4	43.5	41.7	36.4	36.4
Philadelphia	42.2	49.8	49.8	35.7	68.6	53.2	53.0
San Diego	0.0	0.0	0.0	0.0	0.0	0.0	0.0
San Francisco	79.8	74.8	74.8	67.1	53.3	50.0	56.3
Local Median	42.2	48.9	47.8	43.5	53.3	46.2	52.4
Local Range	0.0-92.7	0.0-100.0	0.0-92.7	0.0-100.0	0.0-100.0	0.0-100.0	0.0-100.0
TERRITORIAL SURVEYS							
American Samoa	100.0	100.0	100.0	85.7	100.0	66.7	66.7
Guam	37.5	75.0	12.5	75.0	75.0	25.0	25.0
Northern Mariana Islands	100.0	100.0	66.7	100.0	66.7	66.7	66.7
Palau	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	100.0	100.0	83.4	92.9	87.5	66.7	66.7
Territorial Range	37.5-100.0	75.0-100.0	12.5-100.0	75.0-100.0	66.7-100.0	25.0-100.0	25.0-100.0
TRIBAL SURVEYS							
Cherokee Nation	31.4	24.2	26.2	12.9	15.7	13.3	13.3
Nez Perce	0.0	75.0	75.0	0.0	75.0	0.0	50.0

*Among schools with students in that grade.

TABLE 4. Percentage of Secondary Schools That Provided Those Who Teach Health Education with Materials for Teaching Health Education, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Goals, objectives, and expected outcomes for health education	Chart describing annual scope and sequence of instruction for health education	Plans for how to assess student performance in health education	Written health education curriculum	All 4 types of materials
STATE SURVEYS					
Alabama	90.7	47.3	51.5	75.1	37.8
Alaska	75.2	46.0	48.0	62.6	33.0
Arizona	30.7	34.7	34.7	53.6	33.9
Arkansas	94.6	60.8	75.9	83.6	55.4
California	68.7	48.2	47.9	62.5	35.6
Connecticut	82.1	59.9	64.6	81.9	51.9
Delaware	85.9	80.9	66.7	68.3	52.4
Florida	80.1	59.7	63.4	70.1	51.1
Georgia	91.6	64.3	68.8	80.5	55.7
Hawaii	93.6	76.5	70.4	64.7	54.3
Idaho	88.4	69.5	71.4	78.6	59.1
Indiana	86.9	59.0	61.1	81.7	46.3
Iowa	83.9	63.8	62.0	73.0	50.3
Kansas	78.0	51.2	52.6	70.4	37.2
Kentucky	94.0	69.7	77.1	82.0	60.8
Louisiana	88.4	59.5	71.9	78.9	54.1
Maine	83.7	63.3	65.1	81.1	48.3
Maryland	94.0	82.8	76.0	89.6	69.8
Massachusetts	81.5	61.9	59.4	74.1	48.1
Michigan	84.0	63.7	67.0	76.7	55.4
Minnesota	72.5	56.9	55.8	66.2	41.2
Mississippi	93.2	72.7	84.4	91.3	69.7
Missouri	94.3	70.6	76.6	87.2	63.3
Montana	86.4	65.9	66.5	84.8	53.5
Nebraska	79.3	53.5	58.4	76.8	41.0
Nevada	91.8	73.6	63.7	81.1	54.5
New Hampshire	77.0	59.6	63.6	69.2	43.4
New Jersey	95.3	76.7	80.1	94.4	69.0
New York	85.8	73.9	73.9	79.5	62.9
North Carolina	94.4	59.9	62.5	86.9	47.9
North Dakota	81.0	55.1	58.3	62.8	42.5
Ohio	78.5	54.4	60.1	70.0	44.0
Oklahoma	77.4	44.3	60.7	57.8	37.9
Oregon	89.8	68.8	57.7	73.4	46.3
Pennsylvania	87.3	72.4	71.2	80.0	57.3
Rhode Island	89.0	74.8	75.9	77.2	62.8
South Carolina	88.6	62.1	67.8	74.8	54.7
South Dakota	76.3	44.6	59.7	56.7	35.2
Tennessee	86.7	59.8	72.1	80.6	56.0
Texas	87.6	70.3	73.3	77.3	59.5
Utah	90.7	58.9	59.3	84.6	46.0
Vermont	79.7	56.8	55.5	64.0	44.9
Virginia	93.6	72.5	70.4	88.9	60.0
Washington	79.4	62.1	64.7	63.1	45.6
West Virginia	97.8	65.9	72.6	82.2	57.1
Wisconsin	85.5	67.5	66.1	81.0	55.3
Wyoming	90.3	75.7	86.2	79.7	67.6
State Median	86.7	62.1	65.1	77.3	52.4
State Range	68.7-97.8	42.5-82.8	47.9-86.2	53.6-94.4	33.0-69.8

TABLE 4. Percentage of Secondary Schools That Provided Those Who Teach Health Education with Materials for Teaching Health Education, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Goals, objectives, and expected outcomes for health education	Chart describing annual scope and sequence of instruction for health education	Plans for how to assess student performance in health education	Written health education curriculum	All 4 types of materials
LOCAL SURVEYS					
Albuquerque	87.5	54.7	59.3	53.1	41.9
Baltimore	77.2	59.8	54.5	68.6	43.5
Broward County	87.3	72.5	79.9	85.9	66.7
Charlotte-Mecklenberg County	96.6	64.8	59.1	94.5	48.7
Chicago	75.7	52.1	67.2	62.3	46.4
Detroit	80.3	74.3	72.0	72.7	67.2
District of Columbia	97.1	75.7	79.4	85.0	63.6
Fresno	62.2	42.8	42.8	67.8	33.1
Houston	91.4	80.2	85.3	86.5	77.8
Los Angeles	87.4	65.2	63.3	70.5	49.2
Memphis	93.1	87.2	84.3	87.7	76.4
Miami-Dade County	83.9	67.2	69.5	72.4	58.4
Milwaukee	90.1	78.3	80.7	86.5	70.3
New York City	81.9	65.5	74.0	76.3	58.6
Newark	81.2	74.0	71.8	82.6	67.2
Orange County	100.0	78.2	73.7	90.3	64.0
Philadelphia	83.4	73.2	68.8	65.9	51.0
San Diego	100.0	98.3	1.7	100.0	0.0
San Francisco	94.7	76.5	81.6	86.9	71.2
Local Median	87.4	73.2	71.8	82.6	58.6
Local Range	62.2-100.0	42.8-98.3	1.7-85.3	53.1-100.0	0.0-77.8
TERRITORIAL SURVEYS					
American Samoa	100.0	92.3	92.3	100.0	92.3
Guam	84.6	53.8	46.2	69.2	38.5
Northern Mariana Islands	100.0	50.0	87.5	87.5	50.0
Palau	100.0	100.0	90.9	100.0	90.9
Territorial Median	100.0	73.1	89.2	93.8	70.5
Territorial Range	84.6-100.0	50.0-100.0	46.2-92.3	69.2-100.0	38.5-92.3
TRIBAL SURVEYS					
Cherokee Nation	78.9	52.3	65.1	62.3	49.3
Nez Perce	100.0	71.4	57.1	71.4	57.1

TABLE 5a. Percentage of Secondary Schools in Which Teachers Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Alcohol- or other drug-use prevention	Asthma	Emotional and mental health	Foodborne illness prevention	HIV* prevention	Human sexuality	Injury prevention and safety
STATE SURVEYS							
Alabama	97.1	62.2	75.0	71.1	86.4	75.5	80.1
Alaska	89.2	35.1	81.0	60.1	68.2	69.7	76.2
Arizona	75.9	38.2	62.9	50.0	52.8	48.5	66.8
Arkansas	97.6	80.7	94.9	83.8	93.7	85.6	96.3
California	87.9	42.3	67.6	52.6	87.2	80.8	66.3
Connecticut	91.6	44.6	88.8	65.1	87.2	87.8	78.7
Delaware	98.4	51.6	96.8	65.5	91.7	91.7	82.3
Florida	87.3	50.0	73.4	58.7	82.0	77.1	80.3
Georgia	91.2	49.2	86.3	67.5	89.7	75.7	87.3
Hawaii	97.7	47.0	93.2	66.7	93.8	95.2	92.7
Idaho	98.0	59.4	97.2	78.1	94.6	88.2	93.8
Indiana	97.0	69.3	91.8	84.7	96.1	92.4	91.8
Iowa	92.0	45.1	84.6	70.9	88.5	86.7	80.5
Kansas	94.5	45.3	86.9	67.9	88.7	91.9	82.7
Kentucky	94.8	67.7	90.8	79.9	89.7	80.3	89.7
Louisiana	96.5	48.8	84.7	66.1	72.8	57.6	87.1
Maine	97.7	46.7	96.1	73.2	92.8	89.7	86.4
Maryland	94.6	62.1	93.8	73.0	88.6	87.1	88.1
Massachusetts	91.5	40.2	90.5	58.7	85.3	84.7	79.7
Michigan	91.4	41.3	86.8	62.0	88.7	80.7	77.9
Minnesota	92.9	43.9	89.1	63.3	91.2	93.2	78.5
Mississippi	98.0	67.6	91.1	81.5	87.2	81.3	92.0
Missouri	98.3	72.2	94.1	83.7	93.4	84.6	94.4
Montana	99.1	60.6	90.3	77.7	90.0	85.0	91.9
Nebraska	96.7	58.0	90.9	70.4	89.9	87.3	84.7
Nevada	93.7	56.4	87.0	68.3	95.5	94.5	86.6
New Hampshire	97.2	48.5	93.8	74.4	91.1	91.2	87.3
New Jersey	96.3	67.4	96.4	72.2	94.2	95.0	93.6
New York	97.3	71.2	95.9	76.7	97.2	93.7	89.7
North Carolina	95.7	67.6	89.3	70.5	87.5	78.6	88.6
North Dakota	98.6	62.3	93.9	84.9	92.3	84.5	93.8
Ohio	93.2	57.9	87.1	69.9	87.1	84.7	81.5
Oklahoma	81.3	45.3	60.4	54.9	74.4	59.2	66.0
Oregon	95.5	47.8	92.3	73.9	93.2	93.2	82.9
Pennsylvania	94.3	59.9	91.8	70.2	92.3	89.2	86.7
Rhode Island	98.9	49.0	94.8	52.7	93.8	88.4	89.1
South Carolina	92.4	58.5	88.7	68.3	91.0	89.2	84.4
South Dakota	92.4	42.6	85.7	59.3	77.8	74.1	82.5
Tennessee	89.9	62.2	85.4	72.4	82.9	77.6	86.8
Texas	94.5	61.3	87.9	78.1	85.3	80.1	90.7
Utah	97.3	53.5	95.1	72.1	96.7	92.7	89.1
Vermont	96.7	35.5	89.9	69.8	86.5	90.8	85.3
Virginia	97.7	69.0	91.4	78.7	91.3	84.7	94.6
Washington	93.7	44.4	88.9	69.5	95.4	86.9	82.9
West Virginia	98.7	75.1	96.0	86.4	93.6	89.6	95.3
Wisconsin	98.3	50.1	96.5	72.3	94.9	93.5	89.4
Wyoming	96.4	46.4	94.0	75.2	87.3	82.3	92.2
State Median	95.7	51.6	90.5	70.5	89.9	86.7	86.8
State Range	75.9-99.1	35.1-80.7	60.4-97.2	50.0-86.4	52.8-97.2	48.5-95.2	66.0-96.3

TABLE 5a. Percentage of Secondary Schools in Which Teachers Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Alcohol- or other drug-use prevention	Asthma	Emotional and mental health	Foodborne illness prevention	HIV* prevention	Human sexuality	Injury prevention and safety
LOCAL SURVEYS							
Albuquerque	96.9	46.9	100.0	71.8	96.8	96.8	90.3
Baltimore	80.6	60.5	78.9	56.4	72.1	62.4	70.9
Broward County	86.0	67.7	72.4	66.3	94.1	95.2	75.5
Charlotte-Mecklenberg County	96.7	68.5	85.7	59.4	92.6	88.7	83.6
Chicago	81.2	67.0	67.9	57.7	65.4	67.7	77.8
Detroit	81.7	64.9	72.6	57.2	63.0	67.6	75.0
District of Columbia	88.2	54.6	78.8	67.7	91.2	87.9	84.8
Fresno	77.5	47.3	60.6	33.1	87.3	87.3	54.1
Houston	96.3	76.5	91.5	70.0	88.6	83.2	95.2
Los Angeles	100.0	74.4	97.7	80.9	99.2	97.6	90.4
Memphis	91.5	75.8	79.9	76.7	86.0	78.0	87.8
Miami-Dade County	90.8	52.6	73.9	64.3	87.4	83.7	87.6
Milwaukee	93.5	72.5	86.6	63.8	81.0	90.2	78.5
New York City	92.9	74.7	90.1	70.1	93.6	86.6	80.2
Newark	87.4	76.6	88.8	72.9	82.6	78.9	88.5
Orange County	90.6	64.8	78.8	61.6	100.0	100.0	71.8
Philadelphia	80.5	61.8	78.2	56.8	76.9	76.3	76.5
San Diego	100.0	10.0	98.4	13.4	100.0	100.0	96.5
San Francisco	97.4	70.2	97.4	58.7	97.4	89.6	84.2
Local Median	90.8	67.0	79.9	63.8	88.6	87.3	83.6
Local Range	77.5-100.0	10.0-76.6	60.6-100.0	13.4-80.9	63.0-100.0	62.4-100.0	54.1-96.5
TERRITORIAL SURVEYS							
American Samoa	100.0	76.9	88.5	84.6	100.0	96.2	100.0
Guam	84.6	61.5	76.9	61.5	91.7	58.3	84.6
Northern Mariana Islands	100.0	75.0	87.5	50.0	100.0	100.0	87.5
Palau	100.0	18.2	100.0	27.3	100.0	54.5	100.0
Territorial Median	100.0	68.3	88.0	55.8	100.0	77.3	93.8
Territorial Range	84.6-100.0	18.2-76.9	76.9-100.0	27.3-84.6	91.7-100.0	54.5-100.0	84.6-100.0
TRIBAL SURVEYS							
Cherokee Nation	85.3	44.6	71.8	61.1	76.4	68.4	72.7
Nez Perce	100.0	57.1	85.7	71.4	100.0	100.0	100.0

*Human immunodeficiency virus.

TABLE 5b. Percentage of Secondary Schools in Which Teachers Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
STATE SURVEYS							
Alabama	96.5	99.0	85.3	89.4	76.7	97.9	82.5
Alaska	87.6	94.2	64.4	68.8	65.4	86.8	81.5
Arizona	83.2	90.7	46.6	49.6	43.1	75.1	74.2
Arkansas	98.8	99.6	89.4	91.8	86.8	97.9	94.9
California	86.4	98.2	73.8	85.9	48.9	88.4	78.0
Connecticut	92.3	95.1	78.0	87.0	70.8	91.5	88.0
Delaware	98.4	98.4	89.8	91.7	68.8	98.4	96.7
Florida	89.4	95.4	71.2	77.9	61.3	86.6	86.2
Georgia	91.4	95.0	88.1	89.8	65.8	91.2	82.2
Hawaii	98.8	97.4	94.0	91.3	80.0	96.5	95.4
Idaho	98.7	99.5	83.6	89.2	85.4	97.8	93.7
Indiana	96.5	99.0	91.4	97.1	78.7	97.6	89.0
Iowa	91.9	97.9	81.8	86.3	66.3	93.0	83.7
Kansas	97.0	100.0	82.8	85.7	63.5	94.8	89.8
Kentucky	95.9	96.7	83.8	88.0	74.2	94.7	93.6
Louisiana	95.9	98.7	64.9	69.9	71.2	97.4	90.0
Maine	99.0	98.8	86.8	91.1	71.2	97.1	88.8
Maryland	96.4	97.7	85.3	88.8	83.9	95.0	94.3
Massachusetts	93.8	96.5	78.3	85.3	68.4	91.5	92.5
Michigan	93.9	96.1	73.8	84.1	60.5	92.1	85.6
Minnesota	94.3	96.2	84.3	90.7	77.4	94.6	83.0
Mississippi	99.4	99.5	85.7	87.7	79.2	97.4	96.0
Missouri	99.0	100.0	86.6	91.9	79.2	97.9	91.4
Montana	99.5	99.6	82.3	88.1	76.8	98.2	90.2
Nebraska	96.1	98.0	83.3	93.0	70.7	97.1	89.5
Nevada	95.9	96.5	89.0	92.3	80.9	95.9	92.9
New Hampshire	98.1	99.4	84.1	88.4	74.1	96.6	88.9
New Jersey	98.3	99.6	89.5	94.6	82.5	97.9	94.0
New York	98.7	99.5	89.2	94.1	82.5	97.2	94.4
North Carolina	96.6	97.3	82.0	86.1	72.4	95.3	92.9
North Dakota	99.4	100.0	75.0	85.0	83.4	99.2	93.9
Ohio	93.9	94.6	83.7	86.3	76.3	92.6	87.7
Oklahoma	79.1	84.0	62.1	67.5	48.6	79.2	73.3
Oregon	96.1	98.1	90.0	93.2	78.1	95.7	91.8
Pennsylvania	98.1	98.7	82.2	90.4	68.1	93.8	89.2
Rhode Island	97.8	98.9	79.7	88.6	71.1	92.5	95.8
South Carolina	95.7	98.2	87.5	91.6	59.4	91.4	86.1
South Dakota	88.7	93.2	68.3	76.2	68.8	90.6	87.5
Tennessee	91.6	97.5	78.9	82.6	75.3	88.7	88.5
Texas	96.3	98.8	82.3	86.8	73.8	94.0	89.8
Utah	97.8	99.5	82.4	93.0	88.6	97.8	92.6
Vermont	96.6	99.2	77.4	83.9	73.4	95.7	92.4
Virginia	98.2	98.8	84.6	89.5	73.4	96.8	95.2
Washington	98.0	97.9	81.6	91.9	73.0	94.5	89.1
West Virginia	100.0	100.0	87.9	91.5	81.1	99.4	97.8
Wisconsin	98.3	99.1	90.5	94.0	90.8	97.4	91.8
Wyoming	98.3	99.1	78.7	88.2	82.9	99.1	89.5
State Median	96.5	98.4	83.3	88.6	73.8	95.7	89.8
State Range	79.1-100.0	84.0-100.0	46.6-94.0	49.6-97.1	43.1-90.8	75.1-99.4	73.3-97.8

TABLE 5b. Percentage of Secondary Schools in Which Teachers Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
LOCAL SURVEYS							
Albuquerque	100.0	100.0	96.8	100.0	87.4	100.0	96.9
Baltimore	89.4	90.5	58.3	63.6	53.4	77.6	81.5
Broward County	84.1	96.8	81.3	95.5	62.9	89.1	88.0
Charlotte-Mecklenberg County	98.3	98.3	93.1	96.5	80.1	96.7	93.3
Chicago	88.3	93.3	61.1	59.6	44.3	77.6	83.6
Detroit	86.9	92.6	57.1	63.4	53.5	77.6	82.0
District of Columbia	93.9	96.8	76.0	85.0	57.7	87.8	79.3
Fresno	76.5	94.1	81.0	100.0	55.0	76.2	77.5
Houston	97.6	100.0	82.2	89.8	75.2	93.7	95.1
Los Angeles	100.0	99.2	97.6	99.2	81.8	100.0	97.6
Memphis	89.5	91.2	75.1	82.6	73.2	82.7	89.6
Miami-Dade County	98.1	98.1	73.5	84.8	71.8	90.3	89.7
Milwaukee	93.7	98.3	83.2	81.3	74.7	89.7	93.4
New York City	97.1	99.2	81.4	89.1	71.9	92.1	90.3
Newark	91.1	98.0	85.0	85.7	65.4	90.5	93.6
Orange County	92.7	97.6	100.0	100.0	76.4	88.2	90.6
Philadelphia	92.7	95.6	68.1	75.6	55.3	78.6	83.7
San Diego	51.6	100.0	100.0	100.0	20.7	98.3	98.3
San Francisco	97.3	100.0	87.1	92.3	87.1	100.0	97.3
Local Median	92.7	98.0	81.4	89.1	71.8	89.7	90.3
Local Range	51.6-100.0	90.5-100.0	57.1-100.0	59.6-100.0	20.7-87.4	76.2-100.0	77.5-98.3
TERRITORIAL SURVEYS							
American Samoa	100.0	100.0	92.3	100.0	92.3	100.0	100.0
Guam	91.7	100.0	75.0	91.7	53.8	84.6	84.6
Northern Mariana Islands	100.0	100.0	100.0	100.0	87.5	100.0	87.5
Palau	100.0	100.0	63.6	100.0	100.0	100.0	100.0
Territorial Median	100.0	100.0	83.7	100.0	89.9	100.0	93.8
Territorial Range	91.7-100.0	100.0-100.0	63.6-100.0	91.7-100.0	53.8-100.0	84.6-100.0	84.6-100.0
TRIBAL SURVEYS							
Cherokee Nation	81.4	85.6	62.8	70.2	49.8	77.0	76.3
Nez Perce	100.0	100.0	71.4	85.7	71.4	100.0	100.0

*Sexually transmitted disease.

TABLE 6. Percentage of Secondary Schools with a Health Education Curriculum That Addressed Specific Skills, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Comprehending concepts related to health promotion and disease prevention to enhance health	Analyzing the influence of family, peers, culture, media, technology, and other factors on health behaviors	Accessing valid information and products and services to enhance health	Using interpersonal communication skills to enhance health and avoid or reduce health risks	Using decision-making skills to enhance health	Using goal-setting skills to enhance health	Practicing health-enhancing behaviors to avoid or reduce risks	Advocating for personal, family, and community health	All 8 skills
STATE SURVEYS									
Alabama	92.3	77.2	78.3	78.6	83.7	83.9	79.7	75.7	60.1
Alaska	86.5	84.5	75.8	84.9	87.1	84.8	86.0	78.8	48.9
Arizona	68.6	64.1	59.3	62.3	68.4	64.1	67.9	58.9	37.2
Arkansas	96.6	95.5	93.5	95.1	97.5	96.6	97.1	94.4	77.4
California	78.4	76.6	65.0	74.5	79.0	75.5	76.5	73.0	47.5
Connecticut	86.6	85.9	80.2	85.5	87.8	85.5	86.3	81.4	68.7
Delaware	92.1	92.1	88.9	92.1	92.1	92.1	93.5	90.3	63.9
Florida	83.1	81.1	73.4	83.0	86.3	82.1	81.9	77.0	52.4
Georgia	94.7	94.6	86.4	91.4	95.4	94.2	93.9	89.2	69.8
Hawaii	96.6	95.4	97.7	96.6	96.6	97.7	97.7	94.1	62.8
Idaho	96.9	95.6	89.5	95.3	96.9	93.9	94.0	90.9	68.6
Indiana	98.0	97.4	89.6	95.6	98.3	94.5	97.4	92.6	71.2
Iowa	97.1	95.9	87.2	93.2	96.8	93.8	96.4	89.2	61.8
Kansas	85.6	83.0	78.6	82.4	87.8	81.7	84.5	79.4	55.3
Kentucky	95.3	94.2	89.9	94.5	96.8	96.2	95.4	91.0	74.2
Louisiana	89.3	90.3	85.0	87.9	92.1	89.6	92.4	87.2	65.5
Maine	97.7	96.6	90.9	97.3	97.1	94.1	97.0	91.2	70.0
Maryland	95.4	94.9	90.0	93.1	94.4	93.1	93.9	91.3	82.8
Massachusetts	90.4	90.3	78.0	90.5	93.4	88.5	91.7	82.2	59.4
Michigan	87.8	87.4	77.7	87.5	88.1	83.5	86.8	82.0	58.4
Minnesota	90.2	91.5	81.6	89.9	93.8	86.6	90.7	84.6	52.5
Mississippi	94.3	94.4	88.7	92.3	94.9	94.3	95.3	92.9	83.0
Missouri	98.0	95.9	91.0	93.2	98.3	96.4	97.7	93.9	75.7
Montana	97.6	95.6	88.6	93.3	98.5	92.7	97.5	92.0	69.8
Nebraska	94.2	92.3	84.6	87.8	92.8	89.1	92.8	84.7	60.1
Nevada	99.1	97.4	90.2	96.1	99.1	95.8	98.3	94.0	70.6
New Hampshire	91.6	94.9	86.3	93.7	95.0	90.5	93.9	85.1	58.9
New Jersey	98.3	96.7	91.3	97.0	98.7	96.3	98.4	95.1	82.8
New York	96.2	94.4	90.3	95.1	96.0	94.0	95.4	92.8	72.5
North Carolina	95.6	95.2	84.6	94.4	96.7	93.2	95.2	91.2	73.6
North Dakota	92.4	92.4	89.4	92.7	94.7	89.7	93.4	89.2	56.4
Ohio	88.8	88.7	80.0	86.6	90.7	86.7	88.7	83.0	57.7
Oklahoma	73.1	68.4	65.3	69.0	74.8	71.9	73.9	68.8	47.1
Oregon	95.6	94.4	87.5	95.7	97.2	92.6	97.7	90.1	64.9
Pennsylvania	92.5	91.5	83.6	90.9	93.4	91.2	93.1	88.4	67.6
Rhode Island	95.7	94.7	90.6	94.7	95.7	93.7	95.6	88.2	66.2
South Carolina	91.2	89.1	82.7	88.3	90.0	90.9	90.0	87.4	66.6
South Dakota	80.3	79.1	72.2	79.4	81.8	75.5	78.2	76.7	47.4
Tennessee	84.7	83.9	78.2	83.9	86.4	85.3	86.1	82.2	68.0
Texas	91.1	90.0	84.7	90.5	92.4	91.6	91.2	89.2	68.6
Utah	98.4	97.4	86.8	96.6	97.9	95.7	97.9	93.2	69.0
Vermont	94.2	93.3	83.8	92.5	92.4	88.2	93.4	85.1	56.4
Virginia	95.2	94.9	88.9	92.0	95.5	93.4	94.8	93.8	81.0
Washington	93.6	89.3	81.0	89.6	92.2	93.4	93.8	82.0	50.7
West Virginia	98.0	98.7	93.2	98.7	98.7	95.7	98.7	95.3	77.2
Wisconsin	95.8	94.4	89.4	94.6	96.3	94.2	95.8	88.7	68.6
Wyoming	96.5	92.0	91.3	93.0	96.5	93.1	95.7	88.8	71.8
State Median	94.2	93.3	86.4	92.3	94.7	92.6	93.9	88.8	66.6
State Range	68.6-99.1	64.1-98.7	59.3-97.7	62.3-98.7	68.4-99.1	64.1-97.7	67.9-98.7	58.9-95.3	37.2-83.0

TABLE 6. Percentage of Secondary Schools with a Health Education Curriculum That Addressed Specific Skills, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Comprehending concepts related to health promotion and disease prevention to enhance health	Analyzing the influence of family, peers, culture, media, technology, and other factors on health behaviors	Accessing valid information and products and services to enhance health	Using interpersonal communication skills to enhance health and avoid or reduce health risks	Using decision-making skills to enhance health	Using goal-setting skills to enhance health	Practicing health-enhancing behaviors to avoid or reduce risks	Advocating for personal, family, and community health	All 8 skills
LOCAL SURVEYS									
Albuquerque	100.0	96.9	96.9	93.7	96.9	96.9	100.0	93.7	50.0
Baltimore	85.5	82.7	71.4	78.0	83.2	77.8	81.6	76.7	54.5
Broward County	90.6	83.6	84.3	90.9	92.6	86.2	87.4	82.3	71.5
Charlotte-Mecklenberg County	98.2	96.6	96.7	100.0	100.0	100.0	100.0	98.3	89.1
Chicago	74.3	75.3	66.8	72.2	75.3	71.2	77.4	70.8	46.0
Detroit	78.4	67.2	66.6	75.6	79.5	77.1	79.5	69.7	50.7
District of Columbia	94.2	91.1	84.4	94.0	94.0	94.0	94.0	84.9	71.1
Fresno	71.9	77.5	62.2	70.2	77.5	71.9	70.2	70.2	44.4
Houston	95.1	92.7	86.5	93.9	93.9	92.7	95.1	90.2	82.5
Los Angeles	99.2	100.0	96.0	99.2	100.0	97.6	99.2	96.9	67.1
Memphis	91.9	83.1	84.3	89.9	93.3	93.3	89.5	86.1	73.8
Miami-Dade County	82.9	81.7	74.3	81.0	85.8	80.9	84.0	78.9	57.5
Milwaukee	88.7	88.7	80.4	87.2	90.1	88.2	87.9	84.9	67.8
New York City	92.0	87.2	80.6	90.2	92.0	89.3	90.8	84.3	63.0
Newark	86.7	86.0	80.0	88.5	86.7	82.2	88.5	83.4	72.8
Orange County	90.3	92.9	90.6	95.3	95.3	95.3	92.9	92.9	78.2
Philadelphia	83.7	79.1	73.8	75.9	80.8	79.2	81.4	80.7	54.0
San Diego	100.0	98.3	100.0	100.0	100.0	98.3	100.0	98.3	98.3
San Francisco	97.4	94.7	92.2	89.4	94.7	94.7	97.4	94.7	76.3
Local Median	90.6	87.2	84.3	89.9	92.6	89.3	89.5	84.9	67.8
Local Range	71.9-100.0	67.2-100.0	62.2-100.0	70.2-100.0	75.3-100.0	71.2-100.0	70.2-100.0	69.7-98.3	44.4-98.3
TERRITORIAL SURVEYS									
American Samoa	100.0	92.3	88.5	96.2	100.0	100.0	96.2	92.3	88.5
Guam	84.6	84.6	69.2	84.6	84.6	76.9	84.6	61.5	53.8
Northern Mariana Islands	100.0	100.0	75.0	100.0	100.0	87.5	100.0	100.0	75.0
Palau	100.0	90.9	90.9	81.8	100.0	100.0	100.0	90.9	63.6
Territorial Median	100.0	91.6	81.8	90.4	100.0	93.8	98.1	91.6	69.3
Territorial Range	84.6-100.0	84.6-100.0	69.2-90.9	81.8-100.0	84.6-100.0	76.9-100.0	84.6-100.0	61.5-100.0	53.8-88.5
TRIBAL SURVEYS									
Cherokee Nation	71.0	70.9	68.4	70.2	73.7	72.7	74.8	68.2	56.3
Nez Perce	100.0	100.0	83.3	100.0	100.0	85.7	100.0	100.0	57.1

TABLE 7a. Percentage of Secondary Schools in Which Teachers Taught Specific Tobacco-Use Prevention Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Identifying tobacco products and the harmful substances they contain	Identifying short- and long-term health consequences of tobacco use	Identifying legal, social, economic, and cosmetic consequences of tobacco use	Understanding the addictive nature of nicotine	Effects of tobacco use on athletic performance	Effects of second-hand smoke and benefits of a smoke-free environment	Understanding social influences on tobacco use, including media, family, peers, and culture
STATE SURVEYS							
Alabama	96.0	89.0	75.9	79.4	79.0	83.7	79.8
Alaska	71.9	77.6	65.1	73.7	69.7	70.5	69.4
Arizona	64.6	69.3	58.6	64.9	60.9	64.9	61.4
Arkansas	97.5	97.9	92.6	95.8	94.7	96.6	95.3
California	77.6	79.0	72.5	77.4	71.8	76.9	74.5
Connecticut	84.0	86.5	79.9	85.7	78.9	83.5	82.5
Delaware	93.4	93.4	93.4	95.1	87.1	95.1	93.3
Florida	80.1	80.5	74.7	79.1	75.2	78.9	75.9
Georgia	89.3	90.2	86.5	87.7	82.9	88.5	87.3
Hawaii	91.4	92.8	85.3	87.9	87.4	91.5	90.2
Idaho	94.8	96.0	92.2	94.8	85.2	94.6	92.5
Indiana	96.9	97.2	94.2	96.6	88.8	96.5	94.6
Iowa	86.9	89.4	79.9	86.7	74.8	85.1	81.2
Kansas	89.4	91.0	85.2	90.5	80.7	88.6	86.6
Kentucky	91.0	91.4	85.5	91.3	84.8	88.7	90.3
Louisiana	93.7	92.8	88.7	91.9	91.3	90.8	90.2
Maine	93.4	95.1	87.0	92.4	82.9	90.6	87.1
Maryland	92.7	92.8	90.1	92.3	83.9	92.3	91.8
Massachusetts	84.7	86.0	76.9	85.7	76.1	82.2	81.9
Michigan	87.0	87.8	84.9	86.8	78.7	86.1	84.1
Minnesota	89.8	91.0	86.1	90.4	74.6	89.5	85.1
Mississippi	93.8	94.3	90.2	93.3	90.9	92.2	92.1
Missouri	96.0	96.9	89.2	96.0	88.6	94.6	89.3
Montana	95.1	97.2	93.2	94.6	91.4	96.6	94.3
Nebraska	93.1	94.6	88.7	90.2	83.3	92.0	91.2
Nevada	92.3	93.2	88.6	93.2	83.1	91.5	90.6
New Hampshire	92.0	92.6	88.0	90.9	85.8	91.9	88.6
New Jersey	94.1	94.5	91.8	94.8	90.4	93.8	94.8
New York	94.1	95.0	90.8	93.3	91.7	93.0	92.0
North Carolina	90.3	90.8	85.4	89.8	85.3	91.1	87.9
North Dakota	95.2	96.9	93.2	95.7	87.6	94.8	91.7
Ohio	87.4	88.5	84.3	88.4	83.6	87.9	87.2
Oklahoma	64.1	65.7	59.1	65.1	61.4	65.3	59.2
Oregon	90.0	91.9	87.8	91.4	78.8	91.0	90.0
Pennsylvania	90.1	91.3	86.6	91.0	84.0	88.6	88.2
Rhode Island	90.2	89.1	84.5	89.3	75.0	87.0	89.2
South Carolina	84.0	86.0	79.4	83.8	81.7	81.6	82.0
South Dakota	84.0	87.8	82.4	85.1	82.9	85.0	84.0
Tennessee	83.4	85.1	78.5	82.6	80.3	81.8	79.9
Texas	90.5	90.7	84.7	88.9	84.3	89.4	88.0
Utah	94.1	96.2	94.7	95.2	84.1	93.9	92.2
Vermont	84.5	89.5	78.3	87.1	80.0	85.3	85.2
Virginia	95.3	95.6	92.2	94.8	90.1	94.1	93.6
Washington	90.2	90.7	86.6	90.7	82.7	89.9	88.5
West Virginia	98.5	99.4	97.1	99.4	95.2	98.6	96.5
Wisconsin	93.0	95.0	91.2	93.2	84.3	93.2	92.5
Wyoming	93.0	94.0	88.8	92.2	86.0	90.4	92.2
State Median	91.0	91.4	86.6	90.7	83.6	90.4	88.6
State Range	64.1-98.5	65.7-99.4	58.6-97.1	64.9-99.4	60.9-95.2	64.9-98.6	59.2-96.5

TABLE 7a. Percentage of Secondary Schools in Which Teachers Taught Specific Tobacco-Use Prevention Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Identifying tobacco products and the harmful substances they contain	Identifying short- and long-term health consequences of tobacco use	Identifying legal, social, economic, and cosmetic consequences of tobacco use	Understanding the addictive nature of nicotine	Effects of tobacco use on athletic performance	Effects of second-hand smoke and benefits of a smoke-free environment	Understanding social influences on tobacco use, including media, family, peers, and culture
LOCAL SURVEYS							
Albuquerque	100.0	100.0	90.3	93.5	83.3	87.1	100.0
Baltimore	71.7	72.1	66.5	70.7	63.9	71.7	68.9
Broward County	78.5	78.5	73.9	78.5	74.6	78.5	75.0
Charlotte-Mecklenberg County	93.3	95.0	88.1	93.3	80.5	93.0	89.9
Chicago	61.2	65.3	49.9	60.2	59.0	62.4	58.2
Detroit	66.7	67.2	60.6	62.3	61.3	62.3	66.4
District of Columbia	76.8	83.9	80.1	80.7	74.1	77.5	71.9
Fresno	70.2	70.2	70.2	76.2	62.2	70.2	70.2
Houston	88.7	88.7	85.0	85.0	84.6	87.2	87.5
Los Angeles	98.4	100.0	94.3	97.6	91.8	100.0	98.4
Memphis	67.5	72.1	63.9	65.7	69.4	69.6	72.1
Miami-Dade County	79.0	79.8	68.5	75.4	74.8	76.0	73.3
Milwaukee	74.3	79.4	69.3	75.9	66.3	76.2	74.3
New York City	85.9	87.5	79.5	84.0	82.2	84.0	80.7
Newark	81.0	81.5	69.4	79.6	80.0	81.5	83.5
Orange County	83.5	90.3	87.9	87.9	83.1	87.9	87.9
Philadelphia	70.3	71.5	63.5	69.6	63.6	70.8	67.1
San Diego	55.0	61.7	56.6	58.3	51.3	63.4	53.1
San Francisco	92.2	92.2	79.0	84.3	81.6	86.9	81.6
Local Median	78.5	79.8	70.2	78.5	74.6	77.5	74.3
Local Range	55.0-100.0	61.7-100.0	49.9-94.3	58.3-97.6	51.3-91.8	62.3-100.0	53.1-100.0
TERRITORIAL SURVEYS							
American Samoa	95.8	88.0	84.0	84.0	91.7	91.7	91.7
Guam	76.9	76.9	69.2	69.2	76.9	76.9	69.2
Northern Mariana Islands	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Palau	100.0	100.0	100.0	100.0	100.0	90.9	100.0
Territorial Median	97.9	94.0	92.0	92.0	95.9	91.3	95.9
Territorial Range	76.9-100.0	76.9-100.0	69.2-100.0	69.2-100.0	76.9-100.0	76.9-100.0	69.2-100.0
TRIBAL SURVEYS							
Cherokee Nation	62.9	64.9	60.1	64.0	61.1	64.2	56.0
Nez Perce	66.7	66.7	66.7	83.3	83.3	83.3	66.7

TABLE 7b. Percentage of Secondary Schools in Which Teachers Taught Specific Tobacco-Use Prevention Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Identifying reasons why students do and do not use tobacco	Making accurate assessments of how many peers use tobacco	Using interpersonal communication skills to avoid tobacco use	Using goal-setting and decision-making skills related to not using tobacco	Finding valid information and services related to tobacco-use prevention and cessation
STATE SURVEYS					
Alabama	83.5	74.6	78.5	81.5	75.2
Alaska	69.7	52.4	69.7	67.1	60.0
Arizona	61.7	48.8	62.0	57.9	51.6
Arkansas	97.0	83.4	94.4	92.1	89.4
California	73.5	60.4	75.0	68.7	61.0
Connecticut	83.4	56.3	81.6	78.4	67.4
Delaware	95.1	81.7	93.3	95.1	86.7
Florida	77.5	63.9	74.9	72.8	67.7
Georgia	88.5	71.6	87.7	87.2	78.6
Hawaii	90.1	67.5	86.5	86.5	70.6
Idaho	93.0	73.1	92.0	91.7	79.6
Indiana	94.4	71.9	90.9	86.8	78.4
Iowa	84.2	61.9	78.1	74.3	71.2
Kansas	85.8	62.9	85.4	78.9	70.0
Kentucky	89.7	71.6	88.9	86.9	74.9
Louisiana	90.2	75.9	89.1	84.8	78.0
Maine	87.1	64.0	85.3	79.9	71.8
Maryland	91.4	76.5	91.0	91.3	81.4
Massachusetts	81.7	63.0	80.4	74.0	63.5
Michigan	84.7	66.9	81.8	79.4	71.8
Minnesota	83.8	58.3	84.6	82.8	66.3
Mississippi	92.8	80.2	90.8	91.8	82.5
Missouri	93.0	70.0	92.3	89.0	81.0
Montana	93.1	76.3	90.7	86.5	82.6
Nebraska	88.6	71.8	85.8	82.5	68.6
Nevada	88.4	70.1	88.2	86.9	82.4
New Hampshire	87.1	70.1	84.8	83.1	68.3
New Jersey	93.5	78.6	92.4	91.1	84.2
New York	93.3	76.8	91.5	89.2	83.7
North Carolina	88.9	69.8	86.6	83.0	75.2
North Dakota	94.3	73.8	88.0	86.4	80.9
Ohio	86.8	73.1	84.5	82.4	74.0
Oklahoma	60.9	52.5	57.6	58.0	52.8
Oregon	88.2	65.4	88.1	80.3	65.4
Pennsylvania	88.8	67.5	88.5	87.6	76.0
Rhode Island	88.1	68.2	84.9	81.8	71.7
South Carolina	83.1	66.3	80.7	78.6	71.9
South Dakota	86.0	61.7	80.0	79.6	64.3
Tennessee	81.1	68.1	78.6	73.4	71.7
Texas	87.9	70.9	87.4	83.4	75.9
Utah	92.9	64.6	94.1	89.0	75.4
Vermont	88.0	72.2	81.9	76.8	63.2
Virginia	93.6	73.8	90.8	89.4	82.4
Washington	87.9	66.0	85.3	77.9	71.0
West Virginia	97.8	85.4	97.8	96.5	92.3
Wisconsin	93.5	74.2	91.4	88.4	76.6
Wyoming	87.6	68.3	89.6	81.7	79.0
State Median	88.2	70.0	86.6	83.0	74.9
State Range	60.9-97.8	48.8-85.4	57.6-97.8	57.9-96.5	51.6-92.3

TABLE 7b. Percentage of Secondary Schools in Which Teachers Taught Specific Tobacco-Use Prevention Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Identifying reasons why students do and do not use tobacco	Making accurate assessments of how many peers use tobacco	Using interpersonal communication skills to avoid tobacco use	Using goal-setting and decision-making skills related to not using tobacco	Finding valid information and services related to tobacco-use prevention and cessation
LOCAL SURVEYS					
Albuquerque	96.7	81.2	100.0	90.0	77.4
Baltimore	69.0	54.8	69.1	66.8	56.0
Broward County	76.8	59.2	76.7	71.3	64.1
Charlotte-Mecklenberg County	95.1	73.5	89.9	82.6	71.7
Chicago	60.3	45.2	57.6	56.7	55.6
Detroit	62.3	53.0	65.8	66.0	59.9
District of Columbia	78.0	58.0	77.5	70.9	62.5
Fresno	70.2	30.8	76.2	51.1	40.4
Houston	89.9	80.9	86.2	86.3	82.4
Los Angeles	98.4	85.9	97.5	96.7	88.6
Memphis	64.4	52.5	63.2	64.4	65.7
Miami-Dade County	76.1	66.9	73.6	70.8	67.6
Milwaukee	79.5	58.2	77.8	73.3	68.0
New York City	83.8	68.8	80.8	78.2	73.4
Newark	76.6	68.0	78.9	68.7	69.7
Orange County	87.9	77.9	83.2	83.2	80.7
Philadelphia	67.7	56.5	68.8	66.6	53.7
San Diego	51.5	43.2	52.9	47.2	48.9
San Francisco	89.4	73.6	87.1	81.6	78.9
Local Median	76.8	59.2	77.5	70.9	67.6
Local Range	51.5-98.4	30.8-85.9	52.9-100.0	47.2-96.7	40.4-88.6
TERRITORIAL SURVEYS					
American Samoa	79.2	83.3	95.8	95.8	80.0
Guam	69.2	61.5	69.2	69.2	61.5
Northern Mariana Islands	100.0	75.0	100.0	87.5	100.0
Palau	100.0	100.0	72.7	90.9	90.9
Territorial Median	89.6	79.2	84.3	89.2	85.5
Territorial Range	69.2-100.0	61.5-100.0	69.2-100.0	69.2-95.8	61.5-100.0
TRIBAL SURVEYS					
Cherokee Nation	59.9	55.3	60.5	57.2	52.7
Nez Perce	66.7	50.0	83.3	83.3	66.7

TABLE 7c. Percentage of Secondary Schools in Which Teachers Taught Specific Tobacco-Use Prevention Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Supporting others who abstain from or want to quit using tobacco	Supporting school and community action to support a tobacco-free environment	Identifying harmful effects of tobacco use on fetal development	All 15 tobacco-use prevention topics
STATE SURVEYS				
Alabama	75.6	77.3	75.6	56.7
Alaska	59.8	64.3	63.3	36.1
Arizona	53.4	53.6	50.5	34.3
Arkansas	87.7	91.6	92.7	71.5
California	62.7	66.4	67.4	43.1
Connecticut	67.8	60.5	70.8	35.2
Delaware	85.0	79.6	91.8	69.5
Florida	66.9	69.9	69.4	50.7
Georgia	76.0	75.4	83.1	57.0
Hawaii	75.3	72.1	73.9	49.1
Idaho	82.0	80.2	87.5	55.6
Indiana	83.8	81.5	91.3	56.5
Iowa	72.8	73.7	75.6	46.3
Kansas	71.9	75.5	82.5	45.7
Kentucky	79.0	77.3	80.3	57.3
Louisiana	78.3	79.6	80.2	57.9
Maine	69.1	66.9	76.6	43.3
Maryland	80.7	78.0	84.9	57.0
Massachusetts	63.3	60.3	69.7	40.3
Michigan	73.4	70.4	76.6	49.8
Minnesota	67.1	66.4	80.7	36.9
Mississippi	85.6	91.3	89.5	72.2
Missouri	81.8	82.4	87.8	58.3
Montana	78.8	88.7	85.8	55.4
Nebraska	73.8	77.1	82.8	51.6
Nevada	80.6	79.8	90.0	58.7
New Hampshire	76.0	71.4	79.0	49.0
New Jersey	83.7	83.0	90.0	64.4
New York	83.3	81.2	88.1	61.8
North Carolina	78.5	83.1	79.8	54.1
North Dakota	79.3	81.9	81.6	53.0
Ohio	75.9	74.3	79.5	52.6
Oklahoma	54.1	56.7	52.4	38.9
Oregon	74.5	70.5	79.8	41.4
Pennsylvania	74.1	71.3	82.0	49.3
Rhode Island	65.1	64.4	76.7	48.2
South Carolina	74.1	73.0	80.8	56.1
South Dakota	68.4	69.0	73.0	44.1
Tennessee	72.1	73.8	72.5	53.6
Texas	77.2	78.3	81.5	58.7
Utah	79.0	81.0	84.2	46.5
Vermont	67.7	64.5	68.5	42.0
Virginia	81.6	83.5	85.3	62.9
Washington	68.8	69.4	77.8	45.5
West Virginia	90.9	90.0	92.0	77.6
Wisconsin	80.0	77.6	87.0	52.1
Wyoming	77.2	79.7	78.0	52.8
State Median	75.9	75.5	80.3	52.6
State Range	53.4-90.9	53.6-91.6	50.5-92.7	34.3-77.6

TABLE 7c. Percentage of Secondary Schools in Which Teachers Taught Specific Tobacco-Use Prevention Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Supporting others who abstain from or want to quit using tobacco	Supporting school and community action to support a tobacco-free environment	Identifying harmful effects of tobacco use on fetal development	All 15 tobacco-use prevention topics
LOCAL SURVEYS				
Albuquerque	74.1	73.2	90.3	53.1
Baltimore	54.2	56.0	59.7	36.0
Broward County	63.0	59.2	63.0	48.0
Charlotte-Mecklenberg County	77.0	74.6	84.4	48.0
Chicago	49.9	50.9	57.3	36.0
Detroit	54.1	57.6	56.4	45.1
District of Columbia	62.5	59.2	61.5	38.8
Fresno	43.0	42.1	52.3	12.9
Houston	79.8	80.0	82.3	69.6
Los Angeles	92.8	89.5	95.1	71.0
Memphis	60.8	64.2	63.9	47.8
Miami-Dade County	69.7	74.7	69.9	51.9
Milwaukee	66.4	69.5	73.1	51.6
New York City	71.0	69.2	78.6	53.2
Newark	76.5	67.7	74.2	52.9
Orange County	83.2	76.0	83.1	68.8
Philadelphia	54.9	53.9	64.4	34.6
San Diego	49.1	51.5	53.4	36.7
San Francisco	72.9	81.6	79.0	59.4
Local Median	66.4	67.7	69.9	48.0
Local Range	43.0-92.8	42.1-89.5	52.3-95.1	12.9-71.0
TERRITORIAL SURVEYS				
American Samoa	83.3	91.7	88.0	72.0
Guam	69.2	76.9	69.2	53.8
Northern Mariana Islands	75.0	87.5	87.5	50.0
Palau	90.0	100.0	100.0	70.0
Territorial Median	79.2	89.6	87.8	61.9
Territorial Range	69.2-90.0	76.9-100.0	69.2-100.0	50.0-72.0
TRIBAL SURVEYS				
Cherokee Nation	54.3	54.3	49.3	38.2
Nez Perce	66.7	66.7	83.3	50.0

TABLE 8a. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 6, 7, or 8 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	The differences between HIV and AIDS [‡]	How HIV and other STDs are transmitted	How HIV and other STDs are diagnosed and treated	Health consequences of HIV, other STDs, and pregnancy	The relationship among HIV, other STDs, and pregnancy	The relationship between alcohol and other drug use and risk for HIV, other STDs, and pregnancy	The benefits of being sexually abstinent
STATE SURVEYS							
Alabama	84.2	81.5	72.2	73.1	68.6	74.1	74.1
Alaska	42.7	43.1	40.1	41.6	36.2	41.7	44.6
Arizona	32.1	33.7	24.6	31.1	28.5	30.0	31.7
Arkansas	81.7	81.8	76.4	79.3	74.9	77.7	81.2
California	81.1	81.6	72.4	79.3	75.1	73.3	78.9
Connecticut	73.0	78.8	66.3	76.8	67.0	71.0	75.8
Delaware	67.7	70.6	64.7	69.7	64.7	70.6	73.5
Florida	73.2	74.6	68.1	70.4	67.3	64.9	73.5
Georgia	85.8	87.2	82.0	84.8	83.5	86.7	88.0
Hawaii	90.1	94.8	83.0	87.6	83.0	83.0	92.4
Idaho	73.4	74.4	63.1	71.9	67.3	73.4	77.8
Indiana	90.5	91.5	84.5	90.2	88.8	90.8	93.8
Iowa	76.6	78.4	68.3	79.2	68.0	73.3	77.7
Kansas	75.4	77.2	68.7	74.1	71.6	70.4	78.4
Kentucky	78.6	81.9	67.9	77.8	70.0	78.8	79.8
Louisiana	47.9	47.6	44.1	47.1	42.2	46.9	45.3
Maine	82.1	83.9	69.4	82.7	74.2	77.9	82.3
Maryland	81.4	81.5	77.1	77.5	71.8	76.0	79.4
Massachusetts	72.4	75.1	69.6	71.3	63.3	70.2	74.4
Michigan	76.2	75.8	73.4	73.0	66.6	69.3	77.5
Minnesota	80.2	82.5	76.8	75.5	70.3	78.5	87.7
Mississippi	57.0	56.4	52.7	57.2	53.0	54.9	56.5
Missouri	76.6	79.1	69.4	79.0	71.2	76.7	79.3
Montana	72.8	77.0	66.4	74.0	65.8	72.6	77.2
Nebraska	63.6	68.9	58.7	66.7	62.6	65.9	70.3
Nevada	93.6	95.5	87.7	95.5	92.3	90.7	90.4
New Hampshire	73.6	77.7	71.9	76.6	70.9	76.2	78.7
New Jersey	88.5	89.9	85.4	87.3	85.8	87.4	88.0
New York	92.8	93.4	88.4	90.9	85.4	87.7	90.9
North Carolina	73.8	75.9	66.2	72.9	68.2	71.5	77.0
North Dakota	75.4	77.8	64.4	72.1	64.2	73.2	75.0
Ohio	71.2	73.8	65.5	70.9	70.3	72.1	74.3
Oklahoma	65.5	65.8	60.2	62.6	59.7	58.5	61.0
Oregon	81.0	85.1	67.4	79.1	77.7	76.6	84.9
Pennsylvania	78.9	78.3	72.1	76.2	73.1	73.8	79.2
Rhode Island	92.1	92.1	83.9	87.5	77.4	83.6	88.2
South Carolina	90.4	89.4	85.8	86.9	84.5	85.1	89.9
South Dakota	63.7	67.0	59.2	60.6	53.5	60.9	66.2
Tennessee	63.6	65.8	58.5	63.0	57.5	61.2	64.5
Texas	77.9	78.7	71.7	77.8	72.9	75.2	80.7
Utah	92.8	89.7	81.4	86.1	82.8	84.0	85.6
Vermont	77.1	80.7	66.9	73.8	69.9	76.2	80.1
Virginia	83.7	84.5	80.2	82.2	78.9	79.7	85.2
Washington	91.9	91.2	85.9	88.6	81.6	82.9	86.2
West Virginia	84.2	86.2	79.9	82.9	82.6	84.6	90.1
Wisconsin	85.3	86.9	79.6	82.6	79.3	84.4	89.9
Wyoming	75.4	77.6	71.5	75.2	71.8	70.2	78.7
State Median	77.1	78.8	69.6	76.6	70.9	74.1	78.9
State Range	32.1-93.6	33.7-95.5	24.6-88.4	31.1-95.5	28.5-92.3	30.0-90.8	31.7-93.8

TABLE 8a. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 6, 7, or 8 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	The differences between HIV and AIDS‡	How HIV and other STDs are transmitted	How HIV and other STDs are diagnosed and treated	Health consequences of HIV, other STDs, and pregnancy	The relationship among HIV, other STDs, and pregnancy	The relationship between alcohol and other drug use and risk for HIV, other STDs, and pregnancy	The benefits of being sexually abstinent
LOCAL SURVEYS							
Albuquerque	90.0	95.0	90.5	95.0	95.0	89.5	90.0
Baltimore	50.9	54.3	47.5	42.4	39.1	45.9	50.6
Broward County	90.5	93.2	81.7	87.8	83.8	81.0	96.4
Charlotte-Mecklenberg County	93.2	93.4	86.9	93.4	93.2	93.4	93.4
Chicago	52.4	54.7	48.6	49.4	44.8	50.1	53.1
Detroit	51.1	51.1	47.1	48.0	43.7	39.6	51.1
District of Columbia	88.2	88.2	82.4	82.4	82.4	82.4	88.2
Fresno	90.0	80.0	70.0	70.0	70.0	77.8	70.0
Houston	83.1	80.5	76.2	80.1	77.8	75.8	80.1
Los Angeles	97.2	97.2	97.2	97.2	95.8	94.4	97.2
Memphis	68.2	64.3	60.4	60.4	56.9	62.3	64.3
Miami-Dade County	87.7	87.9	84.6	84.7	84.7	84.2	89.2
Milwaukee	76.1	80.4	80.9	80.5	80.5	73.8	85.4
New York City	87.7	87.8	85.1	84.1	81.5	79.4	80.9
Newark	75.6	80.9	75.6	77.6	70.7	67.7	73.7
Orange County	96.3	96.3	96.3	96.3	96.3	96.3	96.3
Philadelphia	66.4	64.8	61.3	65.6	62.5	63.7	64.8
San Diego	100.0	100.0	100.0	100.0	100.0	100.0	100.0
San Francisco	81.6	86.1	77.1	90.7	77.1	77.1	81.6
Local Median	87.7	86.1	80.9	82.4	80.5	77.8	81.6
Local Range	50.9-100.0	51.1-100.0	47.1-100.0	42.4-100.0	39.1-100.0	39.6-100.0	50.6-100.0
TERRITORIAL SURVEYS							
American Samoa	100.0	100.0	100.0	100.0	95.0	95.0	100.0
Guam	87.5	87.5	75.0	75.0	75.0	75.0	87.5
Northern Mariana Islands	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Palau	100.0	100.0	80.0	80.0	80.0	80.0	100.0
Territorial Median	100.0	100.0	90.0	90.0	87.5	87.5	100.0
Territorial Range	87.5-100.0	87.5-100.0	75.0-100.0	75.0-100.0	75.0-100.0	75.0-100.0	87.5-100.0
TRIBAL SURVEYS							
Cherokee Nation	70.7	70.6	63.5	70.9	67.0	67.2	70.7
Nez Perce	66.7	66.7	66.7	66.7	66.7	66.7	66.7

*Human immunodeficiency virus.

†Sexually transmitted disease.

‡Acquired immunodeficiency syndrome.

TABLE 8b. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 6, 7, or 8 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	How to prevent HIV, other STDs, and pregnancy	How to access valid and reliable information, products, and services related to HIV, other STDs, and pregnancy	The influences of media, family, and social and cultural norms on sexual behavior	Communication and negotiation skills [‡]	Goal-setting and decision-making skills [‡]	Compassion for persons living with HIV or AIDS [‡]
STATE SURVEYS						
Alabama	75.8	64.1	65.0	64.8	67.4	60.5
Alaska	44.6	41.9	38.5	39.3	38.9	32.6
Arizona	32.3	25.8	28.6	29.2	26.5	19.8
Arkansas	81.9	74.2	74.7	71.1	72.4	64.9
California	76.0	67.2	67.0	64.0	66.1	62.5
Connecticut	74.0	59.5	67.2	65.4	59.4	62.9
Delaware	72.7	57.6	66.7	63.6	69.7	59.4
Florida	70.5	60.6	65.8	66.2	68.3	60.4
Georgia	87.1	72.1	86.3	84.7	82.4	75.5
Hawaii	90.1	80.6	82.9	80.6	87.4	72.8
Idaho	71.7	59.0	69.1	67.0	66.0	50.9
Indiana	92.0	73.0	86.5	83.1	80.1	66.6
Iowa	78.3	66.0	71.4	64.8	67.2	55.4
Kansas	77.8	58.2	66.8	66.6	65.0	53.0
Kentucky	77.3	63.0	74.9	69.7	70.9	56.3
Louisiana	44.8	38.1	41.2	36.8	41.9	35.3
Maine	80.8	68.0	75.9	74.9	72.0	58.5
Maryland	78.2	70.6	73.9	73.6	76.6	65.0
Massachusetts	72.9	59.5	69.5	64.2	65.4	56.3
Michigan	71.8	57.7	65.3	63.6	66.1	56.1
Minnesota	79.2	64.8	76.3	68.1	73.3	49.7
Mississippi	55.0	51.5	50.7	49.2	53.8	45.6
Missouri	77.8	64.9	70.0	66.8	71.3	58.0
Montana	74.7	60.5	68.8	62.7	63.3	57.1
Nebraska	64.6	58.4	65.9	63.1	59.1	53.7
Nevada	95.5	82.3	82.7	88.9	92.6	70.4
New Hampshire	77.8	61.2	69.9	71.2	70.2	60.0
New Jersey	86.9	77.1	82.6	80.6	80.7	73.8
New York	90.4	81.3	84.7	84.5	83.3	80.2
North Carolina	74.9	61.7	71.5	69.7	68.5	54.6
North Dakota	72.5	57.2	71.5	64.3	64.5	60.8
Ohio	69.9	59.3	69.6	62.9	70.8	57.5
Oklahoma	63.1	54.0	52.1	53.5	53.9	52.5
Oregon	82.5	67.6	77.9	74.3	70.7	57.2
Pennsylvania	75.3	64.0	67.4	67.2	72.2	53.2
Rhode Island	88.2	75.1	79.5	75.3	71.8	70.3
South Carolina	87.5	73.7	85.9	81.1	84.6	67.5
South Dakota	61.8	51.5	60.2	55.7	55.0	41.4
Tennessee	63.2	53.6	58.0	55.9	57.8	45.7
Texas	76.9	65.4	73.7	69.9	73.2	56.1
Utah	83.0	59.1	82.2	79.2	84.0	63.8
Vermont	78.3	65.6	69.6	67.6	61.3	55.6
Virginia	85.0	71.6	81.6	77.5	78.6	62.4
Washington	85.6	75.6	73.1	74.8	70.3	66.0
West Virginia	85.0	76.5	86.1	77.3	80.3	72.4
Wisconsin	84.1	71.2	80.7	79.7	78.9	72.3
Wyoming	75.5	60.9	72.9	73.1	71.0	56.7
State Median	77.3	64.0	71.4	67.6	70.3	58.0
State Range	32.3-95.5	25.8-82.3	28.6-86.5	29.2-88.9	26.5-92.6	19.8-80.2

TABLE 8b. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 6, 7, or 8 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	How to prevent HIV, other STDs, and pregnancy	How to access valid and reliable information, products, and services related to HIV, other STDs, and pregnancy	The influences of media, family, and social and cultural norms on sexual behavior	Communication and negotiation skills‡	Goal-setting and decision-making skills‡	Compassion for persons living with HIV or AIDS§
LOCAL SURVEYS						
Albuquerque	90.0	95.0	80.0	90.0	89.5	68.5
Baltimore	47.2	40.9	41.1	36.2	43.3	41.2
Broward County	96.4	89.7	77.4	83.9	80.2	78.1
Charlotte-Mecklenberg County	93.4	82.8	89.7	93.4	86.0	69.1
Chicago	49.4	43.2	47.0	41.9	44.0	44.5
Detroit	47.0	42.3	40.2	40.1	42.2	39.8
District of Columbia	88.2	76.5	82.4	76.5	82.4	70.6
Fresno	70.0	54.5	70.0	50.0	50.0	40.0
Houston	83.8	72.7	78.2	76.3	77.4	71.8
Los Angeles	95.7	94.4	97.2	94.4	94.3	94.0
Memphis	60.4	62.8	59.1	58.5	62.3	60.4
Miami-Dade County	87.7	84.6	82.3	79.2	82.0	79.3
Milwaukee	78.2	73.8	79.1	78.2	73.4	68.9
New York City	84.2	76.0	76.6	77.6	75.7	77.3
Newark	71.9	65.5	74.0	62.8	69.9	69.7
Orange County	96.4	96.4	96.4	96.4	96.4	92.9
Philadelphia	64.1	56.5	59.4	56.4	61.9	50.5
San Diego	100.0	100.0	100.0	100.0	100.0	100.0
San Francisco	90.7	90.7	77.1	77.1	81.6	72.6
Local Median	84.2	76.0	77.4	77.1	77.4	69.7
Local Range	47.0-100.0	40.9-100.0	40.2-100.0	36.2-100.0	42.2-100.0	39.8-100.0
TERRITORIAL SURVEYS						
American Samoa	100.0	95.0	90.0	90.0	80.0	45.0
Guam	75.0	62.5	75.0	62.5	75.0	75.0
Northern Mariana Islands	100.0	75.0	100.0	100.0	100.0	100.0
Palau	100.0	70.0	40.0	30.0	100.0	50.0
Territorial Median	100.0	72.5	82.5	76.3	90.0	62.5
Territorial Range	75.0-100.0	62.5-95.0	40.0-100.0	30.0-100.0	75.0-100.0	45.0-100.0
TRIBAL SURVEYS						
Cherokee Nation	70.6	59.4	57.6	62.3	66.5	51.2
Nez Perce	66.7	66.7	66.7	66.7	66.7	33.3

*Human immunodeficiency virus.

†Sexually transmitted disease.

‡Related to eliminating or reducing risk for HIV, other STDs, and pregnancy.

§Acquired immunodeficiency syndrome.

TABLE 8c. Percentage of Secondary Schools in Which Teachers Taught Topics Related to Condom Use, and the Percentage of Secondary Schools in Which Teachers Taught all 17 HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 6, 7, or 8 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Efficacy of condoms	Importance of using condoms consistently and correctly	How to obtain condoms	How to correctly use a condom	All 4 condom use topics	All 17 HIV, STD, and pregnancy prevention topics
STATE SURVEYS						
Alabama	49.4	39.2	32.0	30.9	28.5	27.7
Alaska	27.1	26.4	17.8	10.9	10.1	8.0
Arizona	20.0	13.3	7.3	4.8	5.1	3.7
Arkansas	60.5	46.0	36.6	26.3	26.7	22.8
California	60.4	53.3	41.8	33.5	32.9	26.6
Connecticut	53.7	42.0	30.5	24.1	22.2	18.1
Delaware	51.6	34.4	21.9	19.4	16.1	13.3
Florida	48.9	38.1	30.4	28.5	27.2	25.0
Georgia	48.9	26.5	9.2	6.9	5.8	4.7
Hawaii	84.5	72.3	64.6	63.1	57.3	47.6
Idaho	40.7	24.2	12.6	8.9	7.8	6.6
Indiana	57.4	31.3	16.6	10.7	9.4	7.2
Iowa	59.1	51.6	33.7	31.1	26.8	19.7
Kansas	45.0	37.1	19.6	13.8	14.0	9.8
Kentucky	35.6	24.5	19.3	15.7	17.9	14.1
Louisiana	26.4	20.4	13.1	10.1	10.1	9.5
Maine	62.7	61.6	51.6	41.2	38.7	24.1
Maryland	59.8	49.6	34.9	25.0	23.3	19.1
Massachusetts	56.6	49.2	37.9	28.5	26.6	19.0
Michigan	38.5	21.3	12.4	8.4	9.5	6.6
Minnesota	49.4	41.6	29.3	20.8	18.3	11.0
Mississippi	34.1	30.3	26.0	22.8	21.8	20.6
Missouri	40.4	32.1	19.8	16.0	15.3	13.6
Montana	43.8	35.8	17.8	11.3	12.3	7.3
Nebraska	49.8	37.4	27.7	25.6	24.0	19.3
Nevada	75.8	65.1	28.5	24.2	25.7	17.6
New Hampshire	52.1	44.1	34.0	18.9	21.8	13.3
New Jersey	64.4	56.0	37.4	28.4	28.8	24.2
New York	67.5	62.0	42.4	34.0	30.9	28.5
North Carolina	50.7	34.0	24.6	19.2	18.1	17.9
North Dakota	32.8	25.7	15.1	10.4	9.5	7.9
Ohio	50.7	36.1	25.3	20.1	19.6	16.9
Oklahoma	43.7	38.4	29.8	27.3	26.4	24.9
Oregon	54.6	48.0	37.4	30.4	29.4	20.1
Pennsylvania	47.9	38.2	24.4	19.2	16.7	11.6
Rhode Island	52.0	52.8	24.6	20.2	16.2	15.8
South Carolina	57.7	45.7	29.4	21.9	21.6	19.0
South Dakota	28.4	23.1	16.9	10.6	9.7	10.9
Tennessee	39.1	28.1	21.8	19.6	18.4	16.2
Texas	41.6	28.3	17.1	16.5	16.6	14.2
Utah	19.5	8.1	1.9	0.0	0.0	0.0
Vermont	61.2	59.6	44.8	39.6	39.4	23.6
Virginia	53.1	35.5	22.0	17.4	16.8	13.7
Washington	59.8	55.1	33.6	22.3	17.5	12.9
West Virginia	62.3	58.6	36.0	29.1	26.7	25.7
Wisconsin	58.3	48.2	32.7	29.4	24.9	22.0
Wyoming	36.0	29.7	19.8	9.6	8.6	5.7
State Median	50.7	38.1	26.0	20.2	18.4	16.2
State Range	19.5-84.5	8.1-72.3	1.9-64.6	0.0-63.1	0.0-57.3	0.0-47.6

TABLE 8c. Percentage of Secondary Schools in Which Teachers Taught Topics Related to Condom Use, and the Percentage of Secondary Schools in Which Teachers Taught all 17 HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 6, 7, or 8 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Efficacy of condoms	Importance of using condoms consistently and correctly	How to obtain condoms	How to correctly use a condom	All 4 condom use topics	All 17 HIV, STD, and pregnancy prevention topics
LOCAL SURVEYS						
Albuquerque	85.0	75.0	60.0	52.4	42.8	36.9
Baltimore	37.1	41.4	30.2	24.7	21.0	17.3
Broward County	63.4	59.7	25.5	25.5	19.7	16.2
Charlotte-Mecklenberg County	72.3	63.4	48.3	37.2	37.2	35.3
Chicago	39.3	37.2	24.3	27.3	21.2	16.2
Detroit	29.8	23.0	20.4	16.3	18.6	17.0
District of Columbia	81.3	82.4	76.5	70.6	68.8	64.7
Fresno	36.4	36.4	18.2	0.0	0.0	0.0
Houston	56.6	46.0	36.9	31.6	27.5	27.5
Los Angeles	88.8	87.1	79.9	59.9	56.5	53.6
Memphis	44.3	42.0	27.5	22.1	21.1	21.1
Miami-Dade County	72.8	67.3	45.9	47.0	40.7	39.9
Milwaukee	66.2	66.2	48.8	44.6	40.9	32.9
New York City	66.6	63.1	45.0	38.9	37.4	35.7
Newark	62.1	62.1	42.0	40.5	32.7	31.5
Orange County	96.3	96.3	80.0	72.0	69.3	68.0
Philadelphia	47.3	39.1	26.1	21.1	19.6	16.2
San Diego	100.0	100.0	100.0	100.0	100.0	100.0
San Francisco	71.3	69.9	64.9	64.9	59.9	50.0
Local Median	66.2	63.1	45.0	38.9	37.2	32.9
Local Range	29.8-100.0	23.0-100.0	18.2-100.0	0.0-100.0	0.0-100.0	0.0-100.0
TERRITORIAL SURVEYS						
American Samoa	12.5	16.7	16.7	11.8	11.8	11.8
Guam	50.0	50.0	37.5	25.0	25.0	25.0
Northern Mariana Islands	100.0	75.0	100.0	50.0	50.0	50.0
Palau	50.0	30.0	30.0	30.0	30.0	20.0
Territorial Median	50.0	40.0	33.8	27.5	27.5	22.5
Territorial Range	12.5-100.0	16.7-75.0	16.7-100.0	11.8-50.0	11.8-50.0	11.8-50.0
TRIBAL SURVEYS						
Cherokee Nation	44.3	44.1	30.4	29.1	32.1	21.8
Nez Perce	66.7	66.7	66.7	66.7	66.7	33.3

*Human immunodeficiency virus.

†Sexually transmitted disease.

TABLE 9a. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 9, 10, 11, or 12 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	The differences between HIV and AIDS [‡]	How HIV and other STDs are transmitted	How HIV and other STDs are diagnosed and treated	Health consequences of HIV, other STDs, and pregnancy	The relationship among HIV, other STDs, and pregnancy	The relationship between alcohol and other drug use and risk for HIV, other STDs, and pregnancy	The benefits of being sexually abstinent
STATE SURVEYS							
Alabama	97.2	87.9	84.9	83.6	87.9	86.9	88.9
Alaska	62.0	65.8	59.2	65.2	56.6	62.0	65.1
Arizona	73.0	77.6	67.7	71.5	66.1	70.3	69.3
Arkansas	93.3	97.5	92.8	96.2	95.0	98.7	96.8
California	94.7	93.9	93.2	93.3	89.7	91.9	91.3
Connecticut	95.5	98.3	96.4	97.3	94.2	95.0	95.0
Delaware	100.0	100.0	93.3	96.7	96.7	100.0	100.0
Florida	88.2	88.4	86.3	87.8	87.0	85.8	88.5
Georgia	98.1	99.0	99.0	99.0	98.1	99.1	99.1
Hawaii	93.0	95.3	93.0	90.7	86.0	91.0	95.3
Idaho	96.7	96.4	92.2	97.2	94.5	96.5	97.8
Indiana	98.1	99.4	98.8	97.5	95.6	97.5	99.3
Iowa	85.2	84.7	81.1	84.6	81.9	82.3	82.2
Kansas	95.0	97.2	89.1	95.6	92.8	94.9	95.6
Kentucky	99.2	99.2	97.3	97.4	96.5	96.5	98.3
Louisiana	84.0	87.8	76.3	81.6	80.1	83.6	82.5
Maine	92.7	97.8	92.5	94.7	92.0	95.8	96.7
Maryland	100.0	100.0	99.0	100.0	98.9	100.0	100.0
Massachusetts	87.9	91.4	86.9	89.6	88.1	90.5	90.7
Michigan	94.6	94.6	92.3	91.6	90.9	90.9	93.1
Minnesota	94.6	96.3	93.3	93.4	89.7	94.0	97.1
Mississippi	98.1	98.1	97.2	97.1	95.3	98.1	98.1
Missouri	94.4	97.5	93.1	97.5	95.6	96.9	97.5
Montana	91.2	91.3	88.2	91.4	86.4	91.2	91.4
Nebraska	83.4	87.6	79.5	86.1	82.7	84.7	86.1
Nevada	96.8	100.0	96.6	100.0	100.0	100.0	98.4
New Hampshire	97.3	97.3	95.9	97.3	95.9	97.3	97.3
New Jersey	100.0	100.0	100.0	100.0	100.0	100.0	100.0
New York	99.2	99.5	98.7	99.2	99.2	99.0	98.9
North Carolina	86.8	92.0	83.0	90.3	86.0	88.3	91.9
North Dakota	88.6	88.7	81.9	87.8	83.2	86.8	86.6
Ohio	95.9	97.0	95.3	95.9	95.3	95.2	95.2
Oklahoma	71.8	70.3	66.8	69.4	65.8	67.4	68.2
Oregon	97.0	97.0	96.2	97.0	95.4	96.1	95.5
Pennsylvania	95.2	97.0	96.6	97.1	97.0	95.8	97.0
Rhode Island	93.2	93.3	93.2	93.0	88.9	93.2	93.2
South Carolina	93.2	93.9	92.5	91.8	91.1	92.3	93.9
South Dakota	67.5	73.9	65.5	69.7	65.1	71.8	66.2
Tennessee	95.2	95.2	93.3	94.1	94.2	93.4	96.2
Texas	93.2	95.2	90.5	92.4	91.7	92.6	94.0
Utah	93.6	92.3	89.9	90.7	89.6	92.0	94.6
Vermont	92.0	94.0	89.7	91.5	91.3	93.9	94.0
Virginia	94.2	93.5	87.8	90.4	86.1	89.6	92.3
Washington	98.1	98.1	94.8	97.1	96.4	93.2	95.6
West Virginia	96.7	98.1	96.3	96.3	96.3	100.0	98.1
Wisconsin	93.4	94.5	92.6	93.5	89.7	90.8	95.1
Wyoming	85.5	88.5	84.0	83.9	82.2	85.7	92.1
State Median	94.4	95.2	92.6	93.4	91.3	93.2	95.1
State Range	62.0-100.0	65.8-100.0	59.2-100.0	65.2-100.0	56.6-100.0	62.0-100.0	65.1-100.0

TABLE 9a. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 9, 10, 11, or 12 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	The differences between HIV and AIDS‡	How HIV and other STDs are transmitted	How HIV and other STDs are diagnosed and treated	Health consequences of HIV, other STDs, and pregnancy	The relationship among HIV, other STDs, and pregnancy	The relationship between alcohol and other drug use and risk for HIV, other STDs, and pregnancy	The benefits of being sexually abstinent
LOCAL SURVEYS							
Albuquerque	91.7	100.0	100.0	100.0	100.0	100.0	100.0
Baltimore	95.8	95.8	95.8	95.7	95.7	95.7	91.7
Broward County	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Charlotte-Mecklenberg County	89.2	92.8	89.5	86.1	86.1	96.4	92.8
Chicago	100.0	100.0	95.0	100.0	95.0	100.0	100.0
Detroit	100.0	100.0	91.3	95.7	95.7	91.3	100.0
District of Columbia	92.9	92.9	92.9	92.9	92.9	92.9	92.9
Fresno	100.0	100.0	100.0	100.0	100.0	87.5	100.0
Houston	93.5	96.7	96.7	96.7	96.7	93.3	93.5
Los Angeles	98.1	98.1	98.1	98.1	96.2	98.1	100.0
Memphis	96.0	96.0	96.0	95.8	95.8	87.5	96.0
Miami-Dade County	86.5	86.5	86.5	86.1	86.1	86.1	86.5
Milwaukee	93.3	93.3	93.3	93.3	93.3	86.7	93.3
New York City	97.7	98.4	97.7	97.7	97.5	96.9	97.6
Newark	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Orange County	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Philadelphia	95.6	100.0	100.0	100.0	100.0	97.6	100.0
San Diego	100.0	100.0	100.0	100.0	100.0	100.0	100.0
San Francisco	93.8	93.8	87.5	93.8	87.5	86.7	86.7
Local Median	96.0	98.4	96.7	97.7	96.2	96.4	100.0
Local Range	86.5-100.0	86.5-100.0	86.5-100.0	86.1-100.0	86.1-100.0	86.1-100.0	86.5-100.0
TERRITORIAL SURVEYS							
American Samoa	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Guam	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Northern Mariana Islands	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Palau	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Range	100.0-100.0	100.0-100.0	100.0-100.0	100.0-100.0	100.0-100.0	100.0-100.0	100.0-100.0
TRIBAL SURVEYS							
Cherokee Nation	77.3	79.7	78.6	78.8	73.3	77.0	71.8
Nez Perce	100.0	100.0	100.0	100.0	100.0	100.0	100.0

*Human immunodeficiency virus.

†Sexually transmitted disease.

‡Acquired immunodeficiency syndrome.

TABLE 9b. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 9, 10, 11, or 12 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	How to prevent HIV, other STDs, and pregnancy	How to access valid and reliable health information, products, and services related to HIV, other STDs, and pregnancy	The influences of media, family, and social and cultural norms on sexual behavior	Communication and negotiation skills [‡]	Goal-setting and decision-making skills [‡]	Compassion for persons living with HIV or AIDS [§]
STATE SURVEYS						
Alabama	82.7	85.2	83.6	84.1	86.4	77.1
Alaska	65.9	61.7	56.7	58.2	56.9	50.5
Arizona	71.6	62.6	62.2	63.2	59.5	49.5
Arkansas	96.9	91.0	93.3	91.2	90.0	82.0
California	93.3	89.7	87.7	88.5	89.1	88.8
Connecticut	95.2	93.1	90.4	91.6	89.0	84.2
Delaware	100.0	96.7	96.7	100.0	96.7	83.3
Florida	87.1	83.9	81.1	85.8	84.8	76.9
Georgia	99.0	96.4	96.3	96.4	96.4	88.8
Hawaii	95.3	93.0	95.3	90.7	93.0	80.9
Idaho	97.1	92.8	94.5	93.5	93.3	83.3
Indiana	99.4	90.0	96.2	93.4	92.6	78.0
Iowa	83.8	74.6	76.8	73.6	72.1	62.3
Kansas	94.2	84.0	89.9	91.3	87.0	70.4
Kentucky	97.5	95.7	97.5	96.6	94.9	84.8
Louisiana	82.1	75.5	76.0	74.0	73.7	66.4
Maine	96.9	94.7	89.8	92.1	86.4	74.4
Maryland	100.0	97.6	94.8	100.0	98.9	85.0
Massachusetts	92.0	87.5	84.0	86.2	82.8	75.4
Michigan	92.3	82.3	87.8	86.4	86.3	73.5
Minnesota	96.0	89.8	92.1	88.3	85.1	64.4
Mississippi	98.1	92.5	97.1	90.5	93.3	86.8
Missouri	97.5	92.5	92.4	89.2	90.4	73.8
Montana	91.3	82.1	85.4	82.8	79.7	70.3
Nebraska	85.4	77.3	79.5	70.6	74.8	66.7
Nevada	100.0	96.1	97.3	98.4	98.4	95.7
New Hampshire	97.3	93.1	94.5	90.5	87.4	83.6
New Jersey	100.0	99.0	98.1	97.1	98.1	91.6
New York	99.5	98.3	97.9	96.8	96.9	93.4
North Carolina	90.6	80.5	85.1	81.4	80.9	72.4
North Dakota	86.1	80.6	85.9	79.9	80.7	73.7
Ohio	94.7	91.2	90.9	92.8	90.6	75.5
Oklahoma	68.3	64.9	62.4	64.6	64.9	59.2
Oregon	97.0	94.1	91.6	94.7	93.8	75.4
Pennsylvania	97.0	91.0	93.7	91.6	92.4	81.1
Rhode Island	93.2	92.9	87.4	92.7	88.4	80.9
South Carolina	93.2	88.9	90.3	88.3	89.0	78.7
South Dakota	69.5	61.3	63.8	60.3	60.6	44.8
Tennessee	94.3	92.4	90.5	89.4	89.3	82.1
Texas	92.3	80.0	86.5	83.0	87.7	69.3
Utah	88.1	81.0	88.9	92.3	84.5	74.7
Vermont	94.0	94.0	91.5	85.7	87.2	70.6
Virginia	90.9	89.0	85.7	86.7	87.2	77.2
Washington	97.1	90.9	93.1	91.1	89.6	76.6
West Virginia	98.1	94.4	98.1	94.4	93.0	89.9
Wisconsin	93.9	87.3	90.5	87.7	87.3	74.5
Wyoming	85.5	74.5	79.4	79.0	80.6	68.6
State Median	94.2	90.0	90.4	89.4	87.7	76.6
State Range	65.9-100.0	61.3-99.0	56.7-98.1	58.2-100.0	56.9-98.9	44.8-95.7

TABLE 9b. Percentage of Secondary Schools in Which Teachers Taught Specific HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 9, 10, 11, or 12 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	How to prevent HIV, other STDs, and pregnancy	How to access valid and reliable health information, products, and services related to HIV, other STDs, and pregnancy	The influences of media, family, and social and cultural norms on sexual behavior	Communication and negotiation skills‡	Goal-setting and decision-making skills‡	Compassion for persons living with HIV or AIDS§
LOCAL SURVEYS						
Albuquerque	100.0	100.0	100.0	100.0	100.0	100.0
Baltimore	95.7	86.4	91.7	91.3	95.7	87.5
Broward County	100.0	100.0	91.7	91.7	95.8	87.5
Charlotte-Mecklenberg County	92.8	79.1	92.5	92.5	96.3	70.1
Chicago	100.0	100.0	95.0	89.9	95.0	95.0
Detroit	100.0	100.0	95.7	95.7	95.7	91.3
District of Columbia	92.9	92.9	92.9	85.7	85.7	85.7
Fresno	100.0	87.5	87.5	100.0	87.5	87.5
Houston	90.2	93.5	93.5	89.9	93.5	83.6
Los Angeles	98.1	98.1	96.2	98.1	98.1	96.2
Memphis	88.0	96.0	84.0	95.8	95.8	92.0
Miami-Dade County	86.1	86.1	83.3	86.1	86.1	81.1
Milwaukee	93.3	86.7	86.7	86.7	86.7	86.7
New York City	98.4	97.6	97.6	97.5	94.3	93.5
Newark	100.0	92.4	92.4	92.4	92.4	77.2
Orange County	100.0	100.0	100.0	100.0	100.0	100.0
Philadelphia	97.8	95.4	90.4	95.4	91.0	95.2
San Diego	100.0	100.0	100.0	100.0	100.0	100.0
San Francisco	93.3	86.7	86.7	86.7	86.7	80.0
Local Median	98.1	95.4	92.5	92.5	95.0	87.5
Local Range	86.1-100.0	79.1-100.0	83.3-100.0	85.7-100.0	85.7-100.0	70.1-100.0
TERRITORIAL SURVEYS						
American Samoa	100.0	100.0	100.0	100.0	100.0	83.3
Guam	100.0	100.0	100.0	100.0	100.0	100.0
Northern Mariana Islands	100.0	80.0	100.0	100.0	100.0	100.0
Palau	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Range	100.0-100.0	80.0-100.0	100.0-100.0	100.0-100.0	100.0-100.0	83.3-100.0
TRIBAL SURVEYS						
Cherokee Nation	71.4	70.3	64.8	70.3	70.7	59.9
Nez Perce	100.0	100.0	100.0	100.0	100.0	75.0

*Human immunodeficiency virus.

†Sexually transmitted disease.

‡Related to eliminating or reducing risk for HIV, other STDs, and pregnancy.

§Acquired immunodeficiency syndrome.

TABLE 9c. Percentage of Secondary Schools in Which Teachers Taught Topics Related to Condom Use, and the Percentage of Secondary Schools in Which Teachers Taught all 17 HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 9, 10, 11, or 12 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Efficacy of condoms	Importance of using condoms consistently and correctly	How to obtain condoms	How to correctly use a condom	All 4 condom use topics	All 17 HIV, STD, and pregnancy prevention topics
STATE SURVEYS						
Alabama	71.8	59.5	38.2	36.1	30.6	31.4
Alaska	49.0	47.9	39.5	30.0	30.4	23.2
Arizona	51.0	45.2	26.6	24.8	21.8	14.9
Arkansas	85.2	74.0	61.9	46.7	46.1	41.1
California	89.2	85.8	74.2	72.6	67.5	65.8
Connecticut	91.3	90.3	84.4	77.4	75.6	59.6
Delaware	96.7	96.6	93.1	85.7	85.7	67.9
Florida	64.7	60.1	50.1	41.8	40.4	38.3
Georgia	55.5	44.7	30.2	25.4	23.6	23.8
Hawaii	90.7	86.0	71.3	74.3	68.9	64.5
Idaho	77.4	64.0	47.4	37.0	35.6	31.7
Indiana	77.3	62.3	39.8	35.0	31.9	29.4
Iowa	75.0	67.4	58.1	53.2	51.7	46.3
Kansas	71.4	62.3	53.2	41.9	38.4	34.1
Kentucky	85.8	81.4	69.8	62.6	58.5	54.8
Louisiana	57.0	52.5	41.6	31.8	32.2	31.8
Maine	92.5	91.6	87.5	79.1	77.7	53.5
Maryland	94.6	91.0	77.9	71.6	67.3	56.8
Massachusetts	85.2	84.8	77.5	73.1	69.4	54.1
Michigan	73.3	69.5	53.5	49.2	41.9	34.5
Minnesota	87.9	78.5	61.8	58.4	52.2	33.2
Mississippi	80.5	65.1	55.9	43.1	41.7	38.0
Missouri	73.5	64.1	47.3	38.3	39.7	31.9
Montana	73.3	64.1	56.3	44.0	36.3	25.7
Nebraska	64.6	55.7	41.0	36.0	32.2	27.1
Nevada	86.9	85.6	70.7	61.6	61.7	59.3
New Hampshire	94.5	94.5	88.9	84.5	85.6	67.6
New Jersey	98.1	96.2	94.4	89.9	87.9	83.2
New York	96.2	94.1	90.6	84.1	83.2	76.0
North Carolina	61.2	44.7	32.0	23.4	22.6	20.4
North Dakota	58.2	45.3	30.0	21.1	20.1	17.6
Ohio	84.7	71.8	46.4	51.7	39.3	36.7
Oklahoma	55.7	52.0	39.7	35.9	29.2	30.8
Oregon	88.2	85.9	75.5	68.1	66.5	54.1
Pennsylvania	87.5	78.3	62.2	54.9	53.3	45.3
Rhode Island	88.2	86.5	73.0	68.4	62.9	49.2
South Carolina	82.9	80.0	65.5	61.1	56.3	51.0
South Dakota	37.4	33.1	25.5	21.3	15.9	17.2
Tennessee	73.3	62.1	57.1	43.3	41.4	41.3
Texas	64.3	50.9	33.3	32.9	30.1	27.0
Utah	38.5	26.8	12.4	10.1	9.0	8.9
Vermont	94.0	94.0	90.0	88.0	88.4	57.2
Virginia	72.5	64.9	52.0	45.6	44.7	41.8
Washington	89.9	86.3	73.9	70.5	66.2	52.5
West Virginia	83.8	79.2	60.3	54.3	47.8	47.8
Wisconsin	86.8	81.6	63.9	64.9	56.6	44.1
Wyoming	53.6	50.6	39.9	29.3	28.6	26.6
State Median	80.5	69.5	57.1	49.2	44.7	41.1
State Range	37.4-98.1	26.8-96.6	12.4-94.4	10.1-89.9	9.0-88.4	8.9-83.2

TABLE 9c. Percentage of Secondary Schools in Which Teachers Taught Topics Related to Condom Use, and the Percentage of Secondary Schools in Which Teachers Taught all 17 HIV,* STD,† or Pregnancy Prevention Topics in a Required Course in Any of Grades 9, 10, 11, or 12 During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Efficacy of condoms	Importance of using condoms consistently and correctly	How to obtain condoms	How to correctly use a condom	All 4 condom use topics	All 17 HIV, STD, and pregnancy prevention topics
LOCAL SURVEYS						
Albuquerque	100.0	100.0	100.0	100.0	100.0	90.9
Baltimore	87.5	91.7	75.0	70.8	70.8	65.2
Broward County	87.5	83.3	77.3	68.2	68.2	63.6
Charlotte-Mecklenberg County	58.9	47.9	39.4	26.1	22.4	23.3
Chicago	100.0	100.0	84.9	89.9	84.9	79.9
Detroit	90.5	85.7	85.7	90.5	81.0	71.4
District of Columbia	92.9	92.9	92.9	92.9	92.9	85.7
Fresno	100.0	100.0	100.0	87.5	85.7	71.4
Houston	80.4	80.4	77.3	77.3	78.1	76.6
Los Angeles	100.0	98.1	96.2	96.2	90.6	90.6
Memphis	81.8	70.8	66.7	45.8	47.8	47.8
Miami-Dade County	78.4	77.8	73.0	67.6	63.0	63.9
Milwaukee	93.3	93.3	86.7	85.7	78.6	64.3
New York City	96.9	96.7	96.8	86.7	84.5	79.4
Newark	92.4	92.4	83.5	75.9	75.9	60.7
Orange County	100.0	100.0	100.0	100.0	84.5	100.0
Philadelphia	97.8	95.5	84.3	68.7	64.2	62.4
San Diego	100.0	100.0	100.0	100.0	100.0	100.0
San Francisco	86.7	86.7	93.3	86.7	86.7	80.0
Local Median	92.9	92.9	85.7	86.7	81.0	71.4
Local Range	58.9-100.0	47.9-100.0	39.4-100.0	26.1-100.0	22.4-100.0	23.3-100.0
TERRITORIAL SURVEYS						
American Samoa	66.7	66.7	66.7	50.0	50.0	50.0
Guam	100.0	75.0	100.0	75.0	75.0	75.0
Northern Mariana Islands	100.0	100.0	100.0	100.0	100.0	80.0
Palau	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	100.0	87.5	100.0	87.5	87.5	77.5
Territorial Range	66.7-100.0	66.7-100.0	66.7-100.0	50.0-100.0	50.0-100.0	50.0-100.0
TRIBAL SURVEYS						
Cherokee Nation	60.0	56.0	49.5	45.1	39.7	42.2
Nez Perce	100.0	100.0	100.0	100.0	100.0	75.0

*Human immunodeficiency virus.

†Sexually transmitted disease.

‡Acquired immunodeficiency syndrome.

TABLE 10a. Percentage of Secondary Schools That Taught Specific Nutrition and Dietary Behavior Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Benefits of healthy eating	Food guidance using MyPyramid	Using food labels	Balancing food intake and physical activity	Eating more fruits, vegetables, and whole grain products	Choosing foods that are low in fat, saturated fat, and cholesterol	Using sugars in moderation	Using salt and sodium in moderation
STATE SURVEYS								
Alabama	95.4	80.4	77.5	79.9	84.2	84.2	78.8	79.4
Alaska	83.1	75.1	72.2	80.1	80.6	75.5	80.1	71.8
Arizona	76.0	70.0	65.8	74.1	75.1	69.6	69.8	62.7
Arkansas	98.4	94.5	95.5	98.3	97.5	97.0	97.0	94.7
California	79.2	71.0	71.4	77.0	75.8	73.6	72.7	68.0
Connecticut	90.3	79.2	83.8	89.3	90.0	87.5	86.4	80.1
Delaware	98.4	91.9	88.7	98.4	91.9	93.5	91.9	91.9
Florida	84.6	79.5	78.4	82.6	82.0	80.2	78.4	75.1
Georgia	90.9	87.1	86.4	89.7	90.9	89.6	88.9	88.0
Hawaii	97.7	91.5	94.2	90.1	96.3	91.6	92.5	86.3
Idaho	96.6	92.4	93.2	95.6	95.3	92.7	93.7	89.4
Indiana	95.8	92.8	90.7	93.0	94.4	94.7	92.5	92.4
Iowa	90.9	88.0	86.7	88.8	87.9	86.9	85.0	83.2
Kansas	91.6	89.5	86.0	90.0	90.0	88.9	86.5	84.6
Kentucky	95.1	93.0	92.3	94.6	93.9	93.2	93.4	90.2
Louisiana	93.5	87.5	87.0	91.0	90.7	86.4	85.6	82.8
Maine	96.3	89.4	92.2	94.6	94.9	91.5	90.8	86.5
Maryland	94.9	91.4	89.6	92.7	93.6	92.2	89.8	88.2
Massachusetts	88.6	81.2	81.0	86.6	86.5	84.0	83.1	78.1
Michigan	89.7	83.3	84.3	87.1	86.8	84.0	84.3	81.0
Minnesota	93.0	85.4	85.6	90.9	91.3	90.2	88.1	84.8
Mississippi	97.2	94.2	94.6	96.3	96.2	94.7	93.6	90.5
Missouri	98.4	97.1	97.0	98.0	97.4	97.1	95.8	94.5
Montana	97.5	94.0	96.2	96.7	97.6	96.1	94.3	91.3
Nebraska	95.1	89.2	84.4	93.5	91.3	90.7	86.8	85.5
Nevada	94.2	87.5	89.5	92.1	92.8	91.2	89.5	89.8
New Hampshire	96.9	90.8	90.2	96.4	95.2	92.4	91.9	88.0
New Jersey	97.6	93.6	93.5	95.9	96.2	94.4	93.4	90.4
New York	97.5	93.6	95.0	96.0	96.6	96.4	95.5	92.9
North Carolina	94.0	89.7	86.9	91.7	93.2	90.0	88.6	87.0
North Dakota	97.5	96.9	95.5	97.6	96.3	95.7	94.7	93.4
Ohio	92.1	87.0	85.3	91.0	91.1	90.3	88.5	87.7
Oklahoma	68.8	60.7	54.7	64.8	64.8	59.4	59.4	57.5
Oregon	93.6	87.3	89.2	91.2	93.1	93.2	91.5	88.7
Pennsylvania	95.8	90.8	90.5	94.9	93.6	91.3	90.7	89.3
Rhode Island	94.5	87.2	87.4	91.3	90.4	90.4	87.2	82.6
South Carolina	92.8	85.3	84.2	91.3	89.5	88.3	86.3	85.1
South Dakota	87.8	82.2	80.3	86.8	85.2	81.8	80.7	77.7
Tennessee	87.6	84.7	81.4	87.7	85.5	84.5	81.7	79.8
Texas	94.1	89.2	88.4	92.9	92.1	91.0	88.9	87.6
Utah	96.6	95.1	94.8	96.8	95.6	95.1	95.0	89.3
Vermont	91.3	85.9	87.8	89.6	90.4	89.5	87.8	84.8
Virginia	96.8	94.5	93.7	96.2	95.4	93.7	94.0	91.8
Washington	93.9	88.8	91.2	92.5	92.5	87.9	88.8	83.6
West Virginia	100.0	98.1	98.5	100.0	100.0	99.3	99.4	99.4
Wisconsin	97.8	91.9	93.4	97.2	96.5	94.8	94.1	91.5
Wyoming	97.3	88.9	92.3	97.3	94.7	93.0	93.8	87.9
State Median	94.5	89.2	88.7	92.1	92.5	91.0	88.9	87.6
State Range	68.8-100.0	60.7-98.1	54.7-98.5	64.8-100.0	64.8-100.0	59.4-99.3	59.4-99.4	57.5-99.4

TABLE 10a. Percentage of Secondary Schools That Taught Specific Nutrition and Dietary Behavior Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Benefits of healthy eating	Food guidance using MyPyramid	Using food labels	Balancing food intake and physical activity	Eating more fruits, vegetables, and whole grain products	Choosing foods that are low in fat, saturated fat, and cholesterol	Using sugars in moderation	Using salt and sodium in moderation
LOCAL SURVEYS								
Albuquerque	100.0	87.5	90.6	93.7	96.9	96.9	93.7	93.7
Baltimore	85.0	78.7	75.6	75.3	81.3	76.3	71.7	69.9
Broward County	76.9	70.9	73.6	75.9	75.6	74.6	72.9	71.5
Charlotte-Mecklenberg County	98.3	92.9	85.3	98.3	98.3	94.5	94.5	94.5
Chicago	81.0	74.6	70.4	76.9	78.8	69.6	73.0	62.9
Detroit	81.8	71.2	69.0	76.0	80.4	69.1	69.6	67.7
District of Columbia	90.7	84.4	75.8	78.7	87.8	81.9	75.8	81.9
Fresno	72.3	60.4	49.7	72.3	64.8	54.7	60.6	43.0
Houston	96.3	91.3	92.7	97.6	97.6	93.9	93.8	91.4
Los Angeles	100.0	99.2	100.0	99.2	100.0	99.2	98.3	97.5
Memphis	83.9	80.4	75.7	82.0	82.0	76.9	73.1	69.2
Miami-Dade County	91.0	86.3	80.6	89.2	88.2	83.2	80.4	79.4
Milwaukee	91.7	76.9	77.5	88.2	89.9	88.3	86.3	79.4
New York City	94.3	90.0	90.7	91.4	93.2	91.4	90.5	85.9
Newark	90.5	88.1	83.8	86.4	85.4	83.5	85.4	83.5
Orange County	88.2	85.9	83.5	90.6	88.2	85.8	88.2	81.1
Philadelphia	85.6	80.3	78.9	83.7	83.8	81.0	78.8	78.4
San Diego	33.7	30.1	27.8	29.0	31.3	29.9	25.1	21.0
San Francisco	92.0	89.1	94.5	92.0	92.0	92.0	89.4	81.6
Local Median	90.5	84.4	78.9	86.4	87.8	83.2	80.4	79.4
Local Range	33.7-100.0	30.1-99.2	27.8-100.0	29.0-99.2	31.3-100.0	29.9-99.2	25.1-98.3	21.0-97.5
TERRITORIAL SURVEYS								
American Samoa	100.0	100.0	100.0	96.0	96.0	92.0	84.0	79.2
Guam	83.3	83.3	75.0	83.3	83.3	83.3	83.3	75.0
Northern Mariana Islands	87.5	87.5	87.5	87.5	87.5	87.5	87.5	87.5
Palau	100.0	100.0	72.7	100.0	100.0	81.8	63.6	63.6
Territorial Median	93.8	93.8	81.3	91.8	91.8	85.4	83.7	77.1
Territorial Range	83.3-100.0	83.3-100.0	72.7-100.0	83.3-100.0	83.3-100.0	81.8-92.0	63.6-87.5	63.6-87.5
TRIBAL SURVEYS								
Cherokee Nation	67.8	60.9	53.8	65.2	62.7	59.4	55.4	53.6
Nez Perce	83.3	83.3	83.3	83.3	83.3	83.3	83.3	83.3

TABLE 10b. Percentage of Secondary Schools That Taught Specific Nutrition and Dietary Behavior Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Eating more calcium-rich foods	Food safety	Preparing healthy meals and snacks	Risks of unhealthy weight control practices	Accepting body size differences	Signs, symptoms, and treatment for eating disorders	All 14 nutrition and dietary behavior topics
STATE SURVEYS							
Alabama	76.0	77.0	81.8	81.8	79.1	77.3	68.0
Alaska	71.5	65.0	73.0	71.6	69.5	62.0	44.7
Arizona	62.0	58.2	65.2	67.7	60.2	54.2	40.5
Arkansas	94.1	93.9	91.4	95.4	93.5	94.1	82.5
California	65.7	59.3	67.4	71.1	66.5	59.8	43.7
Connecticut	78.4	69.6	75.7	82.1	78.8	75.6	47.9
Delaware	85.5	77.4	87.1	95.2	93.5	88.3	60.0
Florida	74.2	71.5	76.6	79.7	74.7	71.6	58.3
Georgia	82.9	80.9	82.4	88.0	83.3	82.0	68.7
Hawaii	82.8	72.9	86.9	89.3	87.8	82.3	57.9
Idaho	90.9	84.8	88.1	92.8	91.4	93.9	74.8
Indiana	89.2	88.9	89.1	93.1	88.5	90.4	75.4
Iowa	84.5	76.7	80.8	84.5	83.6	81.4	65.5
Kansas	84.1	76.5	83.1	86.1	81.3	78.3	59.5
Kentucky	87.1	84.3	87.8	91.7	87.4	87.5	71.5
Louisiana	80.9	81.6	85.3	88.7	82.2	79.2	65.5
Maine	86.1	77.5	82.7	89.4	83.5	86.5	56.1
Maryland	84.5	78.6	86.9	89.1	86.1	84.9	64.6
Massachusetts	78.5	62.9	74.6	83.4	78.7	74.3	48.4
Michigan	79.2	67.3	76.7	82.5	79.9	78.1	57.6
Minnesota	80.8	68.8	77.8	86.7	82.3	84.4	53.3
Mississippi	91.0	89.5	92.7	94.6	87.3	85.9	74.7
Missouri	92.3	91.4	93.7	96.1	90.0	90.8	78.4
Montana	92.8	82.8	86.3	93.2	86.5	87.0	71.3
Nebraska	85.1	81.4	84.4	88.7	85.5	85.9	66.8
Nevada	82.9	77.6	81.5	90.0	85.9	86.3	67.2
New Hampshire	85.8	83.0	85.9	89.3	87.9	79.9	63.7
New Jersey	88.1	84.1	90.7	92.4	92.5	89.3	74.2
New York	90.2	82.9	91.8	95.8	92.3	91.8	74.3
North Carolina	83.5	79.2	85.0	91.0	87.1	84.7	64.7
North Dakota	93.7	92.3	91.3	95.0	90.5	90.3	80.5
Ohio	82.3	80.3	85.6	86.2	82.7	81.3	62.1
Oklahoma	55.5	54.6	57.9	60.2	57.9	53.3	44.7
Oregon	83.2	74.5	80.4	85.9	84.6	82.7	54.9
Pennsylvania	88.4	82.4	90.2	90.3	86.8	84.3	68.5
Rhode Island	78.3	67.2	81.6	91.5	82.7	82.7	51.7
South Carolina	83.0	77.7	84.0	89.5	87.6	81.8	66.7
South Dakota	80.5	75.4	76.2	80.5	79.7	78.5	58.5
Tennessee	77.7	74.9	81.3	82.2	79.4	75.1	62.3
Texas	85.7	81.3	85.7	91.3	85.9	87.4	69.7
Utah	87.9	78.1	88.2	95.1	94.0	94.6	68.1
Vermont	83.2	69.5	78.0	78.8	79.1	67.1	44.0
Virginia	87.7	85.3	91.3	93.9	87.4	87.6	72.5
Washington	80.6	69.8	80.4	87.0	84.8	77.9	59.8
West Virginia	97.3	94.4	95.9	97.4	95.1	94.0	88.4
Wisconsin	89.1	81.3	87.2	93.8	92.1	91.2	67.8
Wyoming	84.6	83.9	87.9	93.9	89.0	82.7	69.7
State Median	84.1	78.1	85.0	89.3	85.9	82.7	65.5
State Range	55.5-97.3	54.6-94.4	57.9-95.9	60.2-97.4	57.9-95.1	53.3-94.6	40.5-88.4

TABLE 10b. Percentage of Secondary Schools That Taught Specific Nutrition and Dietary Behavior Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Eating more calcium-rich foods	Food safety	Preparing healthy meals and snacks	Risks of unhealthy weight control practices	Accepting body size differences	Signs, symptoms, and treatment for eating disorders	All 14 nutrition and dietary behavior topics
LOCAL SURVEYS							
Albuquerque	93.7	87.4	84.3	93.7	93.7	90.6	65.5
Baltimore	67.2	70.2	72.4	70.7	70.0	61.3	47.6
Broward County	70.2	68.8	74.2	75.6	71.1	66.2	59.9
Charlotte-Mecklenberg County	91.2	76.6	86.1	90.8	91.2	88.8	63.9
Chicago	62.8	64.8	72.8	69.9	64.7	51.4	41.0
Detroit	67.0	59.5	68.5	70.2	68.3	56.6	50.6
District of Columbia	66.6	69.7	68.7	78.1	58.0	53.3	43.9
Fresno	48.9	31.6	44.4	56.6	53.1	41.4	25.9
Houston	91.3	83.8	91.3	96.3	91.3	86.3	74.8
Los Angeles	97.6	89.3	97.5	98.3	94.3	95.0	84.2
Memphis	65.9	73.8	82.0	80.4	65.0	70.0	55.0
Miami-Dade County	75.4	79.6	80.5	82.3	82.0	74.1	61.0
Milwaukee	81.4	72.5	77.8	83.1	80.0	75.9	58.6
New York City	82.3	79.0	87.5	90.8	86.1	83.1	68.3
Newark	83.5	81.9	78.9	81.5	79.3	67.3	61.9
Orange County	76.4	71.8	83.5	85.8	83.5	81.1	62.4
Philadelphia	78.2	76.4	79.5	79.7	74.7	69.3	56.6
San Diego	19.9	13.8	19.5	27.5	22.9	20.1	13.0
San Francisco	84.2	78.9	94.5	86.7	86.7	81.6	65.8
Local Median	76.4	73.8	79.5	81.5	79.3	70.0	59.9
Local Range	19.9-97.6	13.8-89.3	19.5-97.5	27.5-98.3	22.9-94.3	20.1-95.0	13.0-84.2
TERRITORIAL SURVEYS							
American Samoa	91.7	96.0	88.0	92.0	84.0	80.0	64.0
Guam	83.3	75.0	66.7	83.3	75.0	75.0	58.3
Northern Mariana Islands	87.5	75.0	87.5	100.0	100.0	62.5	50.0
Palau	100.0	90.9	81.8	81.8	72.7	72.7	45.5
Territorial Median	89.6	83.0	84.7	87.7	79.5	73.9	54.2
Territorial Range	83.3-100.0	75.0-96.0	66.7-88.0	81.8-100.0	72.7-100.0	62.5-80.0	45.5-64.0
TRIBAL SURVEYS							
Cherokee Nation	54.6	53.8	55.1	62.4	59.4	48.2	40.3
Nez Perce	83.3	66.7	66.7	83.3	83.3	100.0	66.7

TABLE 11a. Percentage of Secondary Schools That Taught Specific Physical Activity Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Physical, psychological, or social benefits	Health-related fitness	Phases of a workout	How much physical activity is enough	Developing an individualized physical activity plan	Monitoring progress toward reaching goals
STATE SURVEYS						
Alabama	86.3	82.5	86.2	80.5	76.3	75.5
Alaska	82.1	82.1	81.5	74.7	64.0	61.7
Arizona	82.2	81.9	82.6	76.0	63.5	64.3
Arkansas	97.0	95.5	96.4	95.4	87.1	86.4
California	95.0	92.4	89.4	86.8	76.0	76.4
Connecticut	93.2	91.8	89.0	85.4	71.1	68.1
Delaware	98.4	96.8	91.9	88.7	79.0	77.4
Florida	91.8	92.4	91.1	85.8	78.9	79.5
Georgia	91.7	92.1	90.5	90.0	78.7	79.5
Hawaii	98.8	95.0	85.7	87.1	79.1	80.5
Idaho	97.1	95.5	93.1	90.0	79.8	77.7
Indiana	97.0	95.7	93.5	91.4	80.1	76.1
Iowa	95.0	92.0	89.3	84.2	72.1	75.6
Kansas	97.1	96.8	94.3	91.3	66.2	68.2
Kentucky	95.5	95.1	92.6	92.8	80.9	79.6
Louisiana	95.4	96.0	95.1	89.6	74.3	74.2
Maine	95.6	95.9	92.7	90.6	79.7	76.8
Maryland	93.6	91.0	87.5	89.2	78.0	76.2
Massachusetts	93.2	90.4	86.7	83.9	65.9	66.7
Michigan	92.4	91.9	87.8	83.2	71.1	70.5
Minnesota	95.0	92.4	88.6	86.0	74.4	72.6
Mississippi	94.6	93.7	92.7	90.6	82.8	81.4
Missouri	99.7	98.0	97.0	95.7	85.2	80.6
Montana	98.0	98.2	97.6	92.2	73.3	75.5
Nebraska	96.6	95.1	93.6	91.2	73.2	73.2
Nevada	93.5	90.8	85.5	83.9	70.0	71.8
New Hampshire	96.5	95.0	89.9	89.2	71.0	71.6
New Jersey	98.0	97.6	94.8	92.9	77.6	77.5
New York	97.6	95.7	90.3	93.1	81.9	80.9
North Carolina	94.4	93.6	93.4	91.5	77.6	77.8
North Dakota	99.5	95.8	94.8	92.6	74.6	72.2
Ohio	92.8	92.2	86.4	85.2	74.6	73.5
Oklahoma	69.5	66.4	66.1	60.7	49.7	51.1
Oregon	94.9	91.3	85.5	87.1	77.1	73.0
Pennsylvania	96.2	96.8	94.1	92.4	75.8	76.3
Rhode Island	96.7	98.9	96.9	92.3	79.9	79.2
South Carolina	96.8	95.6	94.5	93.5	85.6	85.4
South Dakota	91.1	89.3	82.9	86.7	68.3	70.4
Tennessee	93.4	93.8	93.4	89.2	71.0	73.0
Texas	96.4	95.3	92.2	91.4	78.4	81.5
Utah	99.5	97.9	94.9	95.1	85.3	84.6
Vermont	97.4	96.6	93.7	91.9	73.6	72.6
Virginia	97.3	95.9	96.2	93.7	86.3	85.5
Washington	96.5	95.1	90.0	91.7	85.0	85.2
West Virginia	96.6	98.7	96.7	96.0	82.1	85.2
Wisconsin	97.2	95.6	91.8	91.9	76.7	74.8
Wyoming	96.5	97.5	93.9	93.1	80.9	82.4
State Median	96.2	95.1	92.2	90.6	77.1	76.2
State Range	69.5-99.7	66.4-98.9	66.1-97.6	60.7-96.0	49.7-87.1	51.1-86.4

TABLE 11a. Percentage of Secondary Schools That Taught Specific Physical Activity Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Physical, psychological, or social benefits	Health-related fitness	Phases of a workout	How much physical activity is enough	Developing an individualized physical activity plan	Monitoring progress toward reaching goals
LOCAL SURVEYS						
Albuquerque	100.0	96.9	84.4	90.6	87.4	87.4
Baltimore	85.0	83.1	82.9	78.1	56.9	62.0
Broward County	88.8	90.2	90.3	85.9	71.8	71.8
Charlotte-Mecklenberg County	98.3	94.8	94.8	91.0	82.7	81.2
Chicago	87.9	89.1	86.9	80.5	61.2	65.4
Detroit	86.7	89.1	85.3	78.1	70.7	64.6
District of Columbia	90.4	90.4	90.4	80.1	70.3	77.6
Fresno	89.3	79.9	81.1	77.8	48.2	60.2
Houston	100.0	98.7	98.8	93.8	91.2	88.7
Los Angeles	99.2	94.2	90.0	94.3	87.6	84.2
Memphis	87.6	85.7	86.1	82.3	74.8	76.6
Miami-Dade County	96.1	96.1	95.2	95.2	88.3	88.3
Milwaukee	92.5	90.8	84.6	81.8	61.2	62.8
New York City	95.5	94.4	92.1	91.7	80.2	82.8
Newark	92.3	95.1	93.2	90.2	77.2	85.4
Orange County	92.7	97.6	97.6	97.6	92.7	90.3
Philadelphia	87.9	90.7	86.7	81.6	63.6	71.0
San Diego	100.0	100.0	98.3	100.0	100.0	100.0
San Francisco	94.4	94.4	88.8	88.5	89.0	86.2
Local Median	92.5	94.2	90.0	88.5	77.2	81.2
Local Range	85.0-100.0	79.9-100.0	81.1-98.8	77.8-100.0	48.2-100.0	60.2-100.0
TERRITORIAL SURVEYS						
American Samoa	92.0	88.5	100.0	84.6	84.6	88.5
Guam	91.7	100.0	100.0	90.9	81.8	90.9
Northern Mariana Islands	100.0	100.0	100.0	100.0	100.0	100.0
Palau	100.0	100.0	100.0	100.0	81.8	72.7
Territorial Median	96.0	100.0	100.0	95.5	83.2	89.7
Territorial Range	91.7-100.0	88.5-100.0	100.0-100.0	84.6-100.0	81.8-100.0	72.7-100.0
TRIBAL SURVEYS						
Cherokee Nation	71.4	68.4	67.8	61.6	53.5	56.2
Nez Perce	83.3	83.3	83.3	66.7	66.7	66.7

TABLE 11b. Percentage of Secondary Schools That Taught Specific Physical Activity Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Overcoming barriers to physical activity	Decreasing sedentary activities	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of using performance-enhancing drugs	All 12 physical activity topics
STATE SURVEYS							
Alabama	78.0	78.7	78.4	84.8	76.5	77.7	62.9
Alaska	63.4	70.6	74.1	77.0	71.3	61.3	40.4
Arizona	69.9	78.1	69.4	79.9	77.1	61.2	41.7
Arkansas	89.4	95.3	91.9	97.4	94.8	95.3	75.2
California	78.8	85.8	79.7	86.6	77.8	74.9	55.3
Connecticut	74.4	85.9	79.1	85.5	71.5	83.9	50.7
Delaware	83.6	96.8	90.3	88.7	80.6	87.1	63.9
Florida	80.3	86.4	83.4	87.4	86.7	80.8	63.7
Georgia	80.4	88.0	82.4	88.1	86.6	86.7	62.7
Hawaii	78.1	87.0	92.2	88.3	75.8	82.6	56.7
Idaho	81.4	89.6	85.8	93.8	86.1	87.4	61.4
Indiana	81.5	93.8	83.7	92.8	84.0	92.1	60.2
Iowa	75.5	83.5	76.6	85.3	74.6	80.3	48.4
Kansas	77.5	86.2	79.6	88.7	78.6	84.0	47.7
Kentucky	86.6	91.4	85.9	91.1	85.3	85.4	64.9
Louisiana	83.1	86.5	85.1	93.1	87.5	84.9	60.3
Maine	78.7	89.3	80.0	90.8	80.1	83.0	50.1
Maryland	85.6	90.9	83.9	86.0	79.7	86.0	58.6
Massachusetts	76.8	86.0	79.3	84.8	74.7	78.9	46.4
Michigan	81.8	85.9	78.4	84.7	71.5	80.4	53.0
Minnesota	71.1	84.9	74.5	84.3	71.1	81.9	47.8
Mississippi	84.3	91.1	84.6	93.6	90.1	89.1	67.2
Missouri	87.7	94.1	86.4	95.9	90.0	91.2	64.2
Montana	80.7	94.9	88.4	93.3	81.6	87.9	56.6
Nebraska	84.0	94.0	84.7	90.7	83.0	87.3	58.5
Nevada	75.6	88.5	78.3	85.0	79.8	86.0	54.5
New Hampshire	79.7	93.2	84.2	90.6	79.9	84.1	51.0
New Jersey	87.9	94.0	88.9	94.2	86.6	93.9	61.1
New York	87.6	95.0	88.7	91.0	82.7	92.2	62.8
North Carolina	87.0	91.8	86.9	90.7	84.2	82.6	62.1
North Dakota	82.3	93.9	84.9	95.4	91.5	91.0	55.4
Ohio	79.7	86.0	75.8	85.3	71.6	80.9	51.1
Oklahoma	54.4	62.5	57.1	67.0	58.2	58.5	39.0
Oregon	78.2	88.6	77.4	85.0	74.6	82.4	48.5
Pennsylvania	79.9	94.1	84.9	90.3	79.5	81.7	52.3
Rhode Island	84.5	88.9	82.5	90.0	74.8	77.8	52.5
South Carolina	88.5	91.8	92.1	93.6	85.5	86.3	68.1
South Dakota	70.7	83.0	76.4	81.6	73.3	80.7	51.9
Tennessee	82.2	90.9	84.8	91.4	82.0	75.7	55.6
Texas	84.3	92.0	84.6	92.5	89.4	93.0	65.2
Utah	82.7	94.6	85.6	88.8	82.7	92.3	60.9
Vermont	80.6	87.3	80.9	86.4	77.4	85.0	49.8
Virginia	89.8	95.9	90.1	94.5	92.9	88.0	70.4
Washington	79.7	87.0	78.9	89.5	74.4	83.2	54.6
West Virginia	92.1	99.3	93.4	94.0	93.1	95.8	72.8
Wisconsin	80.9	93.6	85.8	90.1	77.4	90.6	53.9
Wyoming	83.5	91.3	81.6	93.1	86.9	85.4	59.1
State Median	80.9	89.6	83.9	90.0	80.1	84.9	56.6
State Range	54.4-92.1	62.5-99.3	57.1-93.4	67.0-97.4	58.2-94.8	58.5-95.8	39.0-75.2

TABLE 11b. Percentage of Secondary Schools That Taught Specific Physical Activity Topics in a Required Course During the 2009–2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Overcoming barriers to physical activity	Decreasing sedentary activities	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of using performance-enhancing drugs	All 12 physical activity topics
LOCAL SURVEYS							
Albuquerque	81.2	93.7	90.6	96.9	81.2	93.5	62.4
Baltimore	75.3	79.4	77.2	76.3	63.2	69.4	41.2
Broward County	83.3	81.6	85.1	90.2	87.0	77.5	59.8
Charlotte-Mecklenberg County	81.8	93.0	89.2	86.9	79.7	88.6	70.0
Chicago	77.0	79.7	79.5	88.8	66.7	66.6	40.4
Detroit	73.4	79.5	79.2	82.5	59.1	62.1	50.0
District of Columbia	63.2	77.3	80.0	87.2	61.4	72.5	43.4
Fresno	60.9	87.7	68.1	78.6	65.0	80.1	24.6
Houston	92.4	95.0	96.3	97.5	92.5	88.6	79.7
Los Angeles	89.2	98.4	89.3	93.3	90.1	97.5	72.6
Memphis	78.1	85.5	75.1	85.7	75.1	77.4	58.8
Miami-Dade County	90.4	93.1	88.4	95.2	95.1	86.4	74.0
Milwaukee	77.4	83.1	86.1	89.0	66.3	70.2	44.5
New York City	83.6	92.0	85.5	92.1	77.2	84.6	60.7
Newark	87.4	92.5	85.7	90.2	81.1	77.5	68.3
Orange County	85.8	90.6	88.2	92.7	92.8	83.5	69.4
Philadelphia	73.3	85.1	80.8	84.7	65.9	66.9	45.4
San Diego	98.3	100.0	100.0	98.3	98.3	98.3	98.3
San Francisco	89.0	89.0	86.2	88.8	78.1	81.4	67.2
Local Median	81.8	89.0	85.7	89.0	78.1	80.1	60.7
Local Range	60.9-98.3	77.3-100.0	68.1-100.0	76.3-98.3	59.1-98.3	62.1-98.3	24.6-98.3
TERRITORIAL SURVEYS							
American Samoa	88.5	84.6	96.0	96.0	84.6	80.8	57.7
Guam	75.0	83.3	90.9	100.0	90.9	75.0	66.7
Northern Mariana Islands	100.0	100.0	100.0	100.0	87.5	75.0	62.5
Palau	81.8	90.9	100.0	100.0	81.8	100.0	54.5
Territorial Median	85.2	87.8	98.0	100.0	86.1	77.9	60.1
Territorial Range	75.0-100.0	83.3-100.0	90.9-100.0	96.0-100.0	81.8-90.9	75.0-100.0	54.5-66.7
TRIBAL SURVEYS							
Cherokee Nation	59.9	65.0	63.5	67.1	59.6	56.0	43.8
Nez Perce	66.7	66.7	50.0	83.3	83.3	83.3	50.0

TABLE 12. Percentage of Secondary Schools in Which Health Education Staff Worked on Health Education Activities with Other School Staff During the 2009-2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Physical education staff	School health services staff	School mental health or social services staff	Nutrition or food service staff	School health council, committee, or team
STATE SURVEYS					
Alabama	81.3	74.9	51.9	36.9	30.2
Alaska	56.1	48.1	53.4	29.7	27.2
Arizona	62.9	49.8	48.8	35.0	29.5
Arkansas	91.3	83.4	64.5	50.7	58.1
California	52.0	46.6	48.9	24.8	28.2
Connecticut	79.2	60.8	70.4	29.9	39.7
Delaware	95.2	90.5	72.6	58.0	50.0
Florida	74.9	57.1	50.7	37.8	36.5
Georgia	87.5	67.7	53.5	39.0	33.1
Hawaii	72.1	48.1	57.2	25.1	36.2
Idaho	85.3	61.2	70.9	45.0	42.2
Indiana	84.8	79.4	59.1	38.3	40.9
Iowa	74.7	75.1	52.0	42.4	37.2
Kansas	86.5	70.2	53.2	49.2	52.6
Kentucky	88.7	77.8	63.9	50.9	47.4
Louisiana	87.2	74.8	60.2	41.9	42.3
Maine	83.5	74.1	73.5	46.3	58.5
Maryland	81.1	65.1	62.5	25.4	34.4
Massachusetts	82.6	78.0	70.7	42.5	43.8
Michigan	71.1	41.7	56.6	37.8	48.4
Minnesota	81.7	66.9	67.3	34.2	37.7
Mississippi	77.4	73.7	62.7	60.2	60.6
Missouri	85.9	77.8	64.6	47.6	49.0
Montana	82.7	62.8	68.7	36.3	35.8
Nebraska	78.3	68.8	50.5	37.2	31.2
Nevada	84.8	65.3	56.2	27.5	42.9
New Hampshire	83.6	73.6	69.0	44.4	54.9
New Jersey	91.5	83.2	75.7	36.8	42.8
New York	86.4	67.9	74.2	45.2	54.8
North Carolina	88.8	77.8	64.2	37.9	41.7
North Dakota	78.3	45.8	62.8	53.9	41.9
Ohio	82.4	73.2	66.8	41.0	43.5
Oklahoma	67.2	51.7	54.3	44.7	52.2
Oregon	76.4	49.5	58.8	30.2	31.2
Pennsylvania	92.5	80.0	62.2	37.9	47.8
Rhode Island	95.6	78.3	70.9	43.5	48.9
South Carolina	89.0	75.5	59.6	38.9	43.0
South Dakota	68.6	48.0	40.2	36.3	34.6
Tennessee	87.9	81.1	74.0	62.7	73.9
Texas	83.2	74.8	51.9	43.9	50.9
Utah	85.2	56.5	58.7	26.4	37.8
Vermont	71.4	81.8	81.0	51.4	60.5
Virginia	93.4	82.0	67.5	40.5	43.0
Washington	76.1	69.5	67.7	28.6	33.7
West Virginia	89.9	84.6	64.7	52.3	50.5
Wisconsin	83.9	74.2	68.6	41.3	46.5
Wyoming	86.7	78.0	64.0	44.4	50.4
State Median	83.5	73.6	62.8	40.5	42.9
State Range	52.0-95.6	41.7-90.5	40.2-81.0	24.8-62.7	27.2-73.9

TABLE 12. Percentage of Secondary Schools in Which Health Education Staff Worked on Health Education Activities with Other School Staff During the 2009-2010 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Physical education staff	School health services staff	School mental health or social services staff	Nutrition or food service staff	School health council, committee, or team
LOCAL SURVEYS					
Albuquerque	65.7	65.5	71.8	24.9	59.3
Baltimore	73.2	57.6	58.5	32.6	31.0
Broward County	76.6	45.0	63.7	34.2	38.2
Charlotte-Mecklenberg County	93.1	86.6	75.6	38.2	49.9
Chicago	71.0	51.1	63.7	36.0	31.1
Detroit	72.8	50.2	68.5	50.0	44.2
District of Columbia	90.5	87.6	68.9	37.6	48.5
Fresno	50.0	45.9	48.4	9.7	9.7
Houston	91.5	86.5	54.5	42.8	46.7
Los Angeles	57.7	68.3	62.7	29.3	31.0
Memphis	79.4	72.9	74.7	59.2	56.2
Miami-Dade County	78.4	57.7	65.6	48.2	56.5
Milwaukee	64.7	74.6	76.6	46.1	45.7
New York City	84.4	58.8	68.2	40.7	42.4
Newark	87.8	86.0	75.6	50.3	54.6
Orange County	74.2	67.2	62.5	32.6	55.9
Philadelphia	84.9	83.5	58.3	57.2	36.8
San Diego	26.5	43.1	36.2	23.6	11.8
San Francisco	84.2	81.5	94.7	57.8	92.0
Local Median	76.6	67.2	65.6	38.2	45.7
Local Range	26.5-93.1	43.1-87.6	36.2-94.7	9.7-59.2	9.7-92.0
TERRITORIAL SURVEYS					
American Samoa	96.2	61.5	69.2	80.8	76.9
Guam	69.2	61.5	46.2	23.1	30.8
Northern Mariana Islands	87.5	62.5	62.5	62.5	62.5
Palau	100.0	100.0	81.8	81.8	100.0
Territorial Median	91.9	62.0	65.9	71.7	69.7
Territorial Range	69.2-100.0	61.5-100.0	46.2-81.8	23.1-81.8	30.8-100.0
TRIBAL SURVEYS					
Cherokee Nation	75.1	66.4	64.7	56.0	70.4
Nez Perce	85.7	71.4	28.6	57.1	42.9

TABLE 13. Percentage of Secondary Schools That Provided Parents and Families with Health Information on Specific Topics Designed to Increase Parent and Family Knowledge, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	HIV,* STD,† or teen pregnancy prevention	Asthma	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	All 3 topics
STATE SURVEYS						
Alabama	38.7	29.3	45.5	41.7	43.4	30.8
Alaska	13.5	11.0	34.2	33.7	36.0	19.5
Arizona	14.3	20.1	31.7	40.6	42.1	24.5
Arkansas	34.0	37.3	52.8	60.6	58.1	46.3
California	36.0	20.2	36.4	42.8	42.3	28.6
Connecticut	16.2	16.4	26.6	45.9	48.9	18.6
Delaware	34.4	26.7	41.0	52.5	51.6	36.7
Florida	29.9	21.9	35.8	46.9	44.0	29.3
Georgia	31.8	18.2	29.5	34.1	33.7	23.2
Hawaii	35.1	16.9	35.3	39.6	39.8	27.6
Idaho	28.6	9.1	33.8	43.3	43.2	23.7
Indiana	19.7	11.8	25.8	31.5	31.6	18.5
Iowa	19.8	13.0	29.1	39.7	42.0	22.9
Kansas	19.1	10.9	32.7	46.2	46.3	27.0
Kentucky	34.8	21.7	40.9	49.6	49.7	33.4
Louisiana	18.6	20.6	39.8	43.7	45.2	29.1
Maine	22.8	16.6	32.9	44.5	48.0	25.1
Maryland	39.2	24.4	43.2	48.1	45.6	33.3
Massachusetts	22.0	16.6	27.9	39.8	44.1	21.4
Michigan	33.5	16.7	33.1	41.3	41.4	26.6
Minnesota	21.4	12.2	23.5	33.8	34.6	16.3
Mississippi	27.7	29.3	51.3	54.2	58.9	42.6
Missouri	25.7	24.5	32.9	46.8	46.7	29.8
Montana	22.8	17.7	37.0	42.2	43.0	28.0
Nebraska	19.6	30.9	32.3	44.6	42.4	27.9
Nevada	93.3	16.3	33.1	40.7	39.8	25.8
New Hampshire	17.8	13.6	24.3	40.6	44.6	20.8
New Jersey	29.7	34.5	37.9	47.9	49.6	31.4
New York	38.6	30.5	41.6	55.0	54.8	36.8
North Carolina	29.2	22.7	37.0	49.0	42.7	28.2
North Dakota	21.8	15.9	41.4	47.3	48.6	31.9
Ohio	25.0	20.0	33.1	43.4	42.8	26.2
Oklahoma	43.2	30.5	42.8	44.5	47.9	33.9
Oregon	22.8	11.9	23.9	33.5	35.8	16.3
Pennsylvania	22.5	19.3	26.4	44.2	43.7	20.3
Rhode Island	21.8	17.1	24.9	41.5	38.7	20.6
South Carolina	30.4	21.3	31.5	55.1	44.9	28.5
South Dakota	11.0	6.6	33.3	37.5	38.6	26.1
Tennessee	29.4	32.4	46.0	56.2	56.1	39.7
Texas	33.0	22.3	34.1	46.9	46.9	29.6
Utah	34.8	13.1	31.4	34.6	38.4	25.0
Vermont	18.7	22.6	44.5	45.1	43.6	24.5
Virginia	30.0	25.1	32.0	55.7	48.4	28.9
Washington	50.3	21.0	35.5	41.4	39.4	26.0
West Virginia	28.1	32.2	51.1	45.2	53.0	35.7
Wisconsin	30.4	18.1	33.5	51.0	49.9	26.4
Wyoming	18.7	15.2	40.2	49.2	49.2	32.1
State Median	28.1	20.0	33.8	44.5	44.0	27.6
State Range	11.0-93.3	6.6-37.3	23.5-52.8	31.5-60.6	31.6-58.9	16.3-46.3

TABLE 13. Percentage of Secondary Schools That Provided Parents and Families with Health Information on Specific Topics Designed to Increase Parent and Family Knowledge, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	HIV* STD,† or teen pregnancy prevention	Asthma	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	All 3 topics
LOCAL SURVEYS						
Albuquerque	18.6	34.4	24.9	37.6	43.7	18.7
Baltimore	34.0	43.1	39.4	55.7	58.2	35.4
Broward County	36.9	38.3	37.9	50.8	43.5	29.2
Charlotte-Mecklenberg County	41.3	44.3	32.8	56.8	45.9	31.0
Chicago	28.4	47.2	27.2	53.6	55.7	24.0
Detroit	30.9	43.4	44.0	59.8	53.7	41.6
District of Columbia	57.8	56.0	39.6	57.6	60.3	33.5
Fresno	34.7	42.7	33.1	31.0	35.1	23.2
Houston	50.1	45.4	52.6	61.4	61.5	50.1
Los Angeles	47.2	36.2	60.0	50.1	64.5	43.8
Memphis	38.8	44.5	39.7	56.9	57.2	36.1
Miami-Dade County	48.2	34.9	48.0	65.5	64.3	44.9
Milwaukee	23.8	53.8	27.4	57.1	58.2	27.4
New York City	52.1	43.6	39.4	60.5	53.0	36.5
Newark	39.8	46.7	40.2	64.8	62.1	33.9
Orange County	57.1	23.4	42.7	51.2	53.3	33.1
Philadelphia	31.9	46.5	30.1	53.3	64.4	28.7
San Diego	94.8	24.6	35.0	34.7	34.7	25.0
San Francisco	58.3	63.2	68.7	63.0	68.3	55.4
Local Median	39.8	43.6	39.4	56.8	57.2	33.5
Local Range	18.6-94.8	23.4-63.2	24.9-68.7	31.0-65.5	34.7-68.3	18.7-55.4
TERRITORIAL SURVEYS						
American Samoa	42.3	38.5	61.5	69.2	68.0	57.7
Guam	23.1	7.7	46.2	30.8	38.5	30.8
Northern Mariana Islands	50.0	37.5	62.5	50.0	50.0	50.0
Palau	100.0	0.0	63.6	54.5	100.0	54.5
Territorial Median	46.2	22.6	62.0	52.3	59.0	52.3
Territorial Range	23.1-100.0	0.0-38.5	46.2-63.6	30.8-69.2	38.5-100.0	30.8-57.7
TRIBAL SURVEYS						
Cherokee Nation	43.7	33.9	40.8	47.0	54.7	33.9
Nez Perce	14.3	14.3	28.6	28.6	28.6	28.6

*Human immunodeficiency virus.
 †Sexually transmitted disease.

TABLE 14. Percentage of Secondary Schools in Which the Major Emphasis of the Lead Health Education Teacher's Professional Preparation Was in Each Specific Discipline, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Health and physical education combined	Health education only	Physical education only	Other education degree	Kinesiology, exercise science, or exercise physiology; home economics or family and consumer science; biology or other science	Nursing or counseling	Public health, nutrition, or another discipline
STATE SURVEYS*							
Alabama	37.9	21.0	25.4	3.1	7.8	2.9	1.8
Alaska	15.9	1.4	5.4	31.4	16.5	8.5	20.8
Arizona	22.9	2.3	17.2	15.2	10.5	19.8	12.1
Arkansas	71.8	5.4	10.5	3.2	6.1	1.1	1.9
California	15.3	12.1	14.4	12.7	31.8	3.6	10.1
Connecticut	42.2	16.5	21.9	1.6	9.3	4.8	3.7
Delaware	68.2	7.9	12.7	1.6	0.0	7.9	1.6
Florida	35.2	6.2	18.9	6.4	19.1	6.1	8.1
Georgia	66.2	8.6	8.1	9.0	3.2	3.3	1.7
Hawaii	31.2	11.3	19.2	15.1	15.6	4.5	3.1
Idaho	55.2	7.6	14.2	10.7	7.1	2.8	2.5
Indiana	74.2	6.8	10.5	1.9	4.8	0.5	1.3
Iowa	37.2	7.4	12.5	7.0	29.9	3.7	2.3
Kansas	59.6	1.2	22.7	3.8	5.4	4.9	2.4
Kentucky	65.7	3.5	15.8	6.3	3.5	1.8	3.4
Louisiana	77.0	2.5	10.3	4.7	3.2	0.9	1.4
Maine	40.0	21.5	8.4	7.6	12.6	5.8	4.1
Maryland	39.8	24.2	17.5	3.8	7.0	3.4	4.4
Massachusetts	39.1	24.3	12.8	3.1	6.2	8.3	6.2
Michigan	48.7	10.9	14.2	5.9	14.4	2.3	3.5
Minnesota	66.9	11.2	8.2	4.8	2.6	2.6	3.7
Mississippi	44.5	4.9	13.1	9.3	19.5	4.5	4.2
Missouri	54.0	5.1	17.6	4.2	12.5	2.9	3.8
Montana	69.6	2.7	11.0	11.8	2.7	0.5	1.7
Nebraska	45.5	3.1	28.0	2.0	17.3	3.0	1.0
Nevada	45.7	18.5	12.1	9.0	7.4	2.0	5.3
New Hampshire	28.4	26.6	16.5	5.0	9.9	10.7	3.0
New Jersey	79.1	5.6	3.5	2.4	1.0	6.6	1.7
New York	39.1	31.2	16.5	2.6	5.8	1.7	3.1
North Carolina	57.9	5.6	26.0	2.7	2.6	4.0	1.2
North Dakota	46.5	3.2	16.3	10.4	20.5	1.8	1.2
Ohio	65.9	9.9	7.4	5.5	6.9	2.8	1.6
Oklahoma	48.8	1.4	8.7	13.9	11.1	11.9	4.2
Oregon	39.6	14.6	8.4	12.1	12.9	2.6	9.9
Pennsylvania	86.0	2.2	2.9	1.3	3.8	2.6	1.2
Rhode Island	76.8	8.3	5.8	1.1	1.1	5.8	1.1
South Carolina	48.5	3.2	37.9	2.7	4.4	1.2	2.0
South Dakota	55.5	3.4	9.9	11.5	9.2	1.2	9.3
Tennessee	58.1	4.5	15.2	5.5	4.9	7.6	4.1
Texas	50.5	9.5	8.6	4.9	18.5	5.4	2.6
Utah	43.4	24.2	15.8	7.6	7.3	0.0	1.6
Vermont	29.5	14.8	9.5	5.7	15.2	22.3	3.0
Virginia	87.6	0.6	6.5	1.6	1.5	1.9	0.3
Washington	33.5	4.8	15.6	11.0	25.4	5.0	4.7
West Virginia	79.9	9.5	4.0	2.0	1.3	0.0	3.3
Wisconsin	61.9	10.1	11.7	4.8	6.0	2.6	2.9
Wyoming	55.9	0.0	18.7	7.5	13.8	4.2	0.0
State Median	48.8	7.4	12.8	5.5	7.3	3.3	3.0
State Range	15.3-87.6	0.0-31.2	2.9-37.9	1.1-31.4	0.0-31.8	0.0-22.3	0.0-20.8

TABLE 14. Percentage of Secondary Schools in Which the Major Emphasis of the Lead Health Education Teacher’s Professional Preparation Was in Each Specific Discipline, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Health and physical education combined	Health education only	Physical education only	Other education degree	Kinesiology, exercise science, or exercise physiology; home economics or family and consumer science; biology or other science	Nursing or counseling	Public health, nutrition, or another discipline
LOCAL SURVEYS*							
Albuquerque	35.8	53.5	0.0	3.6	7.1	0.0	0.0
Baltimore	24.2	12.0	26.5	8.7	12.1	11.0	5.4
Broward County	30.9	12.5	15.9	4.9	23.2	7.2	5.3
Charlotte-Mecklenberg County	52.0	10.8	30.0	3.6	0.0	0.0	3.6
Chicago	29.8	5.9	27.5	5.9	11.8	13.0	6.0
Detroit	49.9	4.0	26.3	7.0	2.9	4.3	5.6
District of Columbia	72.9	3.8	3.8	3.9	0.0	11.8	3.8
Fresno	11.5	0.0	0.0	5.8	82.7	0.0	0.0
Houston	62.1	2.8	29.3	1.4	2.9	0.0	1.5
Los Angeles	6.2	40.0	2.7	5.3	35.5	1.7	8.7
Memphis	68.5	3.9	14.2	6.1	0.0	3.7	3.7
Miami-Dade County	25.9	5.1	22.3	6.4	19.4	12.5	8.4
Milwaukee	34.6	5.1	10.0	31.8	1.9	9.7	6.9
New York City	31.8	12.2	28.3	6.9	11.2	2.7	6.9
Newark	63.4	0.0	20.4	2.8	9.5	2.0	2.0
Orange County	35.3	21.5	11.9	4.8	24.0	0.0	2.4
Philadelphia	79.3	0.0	6.2	2.3	3.1	9.2	0.0
San Diego	0.0	3.3	0.0	14.8	61.0	7.3	13.7
San Francisco	2.7	5.7	2.7	30.2	27.6	5.6	25.5
Local Median	34.6	5.1	14.2	5.8	11.2	4.3	5.3
Local Range	0.0-79.3	0.0-53.5	0.0-30.0	1.4-31.8	0.0-82.7	0.0-13.0	0.0-25.5
TERRITORIAL SURVEYS *							
American Samoa	73.1	3.8	3.8	7.7	3.8	0.0	7.7
Guam	50.0	8.3	16.7	0.0	8.3	16.7	0.0
Northern Mariana Islands	14.3	0.0	0.0	28.6	28.6	14.3	14.3
Palau	0.0	9.1	9.1	81.8	0.0	0.0	0.0
Territorial Median	32.2	6.1	6.5	18.2	6.1	7.2	3.9
Territorial Range	0.0-73.1	0.0-9.1	0.0-16.7	0.0-81.8	0.0-28.6	0.0-16.7	0.0-14.3
TRIBAL SURVEYS *							
Cherokee Nation	44.6	3.1	11.5	8.4	9.5	21.7	1.2
Nez Perce	83.3	0.0	0.0	16.7	0.0	0.0	0.0

*Percentages for each row might not add up to 100.0 because of rounding.

TABLE 15. Percentage of Secondary Schools in Which All Staff Who Teach Health Education Topics Were Certified* in Health Education, the Lead Health Education Teacher Was Certified to Teach Health Education in Middle School or High School, and the Lead Health Education Teacher Had Experience Teaching Health Education Courses or Topics for a Specific Number of Years, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010

Site	All staff who teach health education topics are certified in health education	Lead health education teacher is certified to teach health education	Number of years lead health education teacher has taught health education courses or topics				
			1 year	2–5 years	6–9 years	10–14 years	≥15 years
STATE SURVEYS							
Alabama	77.0	76.7	5.0	25.0	24.0	15.6	30.3
Alaska	25.6	29.2	21.0	33.7	18.6	11.8	14.9
Arizona	42.8	36.3	20.1	28.0	17.0	9.9	25.0
Arkansas	98.8	94.7	7.9	28.9	16.6	12.3	34.2
California	61.8	58.4	7.2	29.9	21.9	14.4	26.6
Colorado	66.5	NA	NA	NA	NA	NA	NA
Connecticut	82.8	83.5	5.6	21.7	20.9	22.6	29.2
Delaware	92.0	88.7	3.2	27.4	19.4	21.0	29.0
Florida	70.0	59.1	11.2	31.0	15.5	13.1	29.1
Georgia	97.7	95.2	3.3	19.6	16.5	22.7	37.9
Hawaii	76.4	65.2	3.5	29.5	25.0	18.3	23.7
Idaho	87.8	87.5	6.7	22.1	17.7	15.3	38.2
Indiana	97.0	95.2	3.0	17.2	9.3	13.8	56.7
Iowa	86.9	81.2	4.9	25.1	15.1	17.0	37.9
Kansas	84.9	79.3	4.5	26.7	14.9	13.6	40.3
Kentucky	89.1	87.3	5.9	32.0	20.9	12.9	28.2
Louisiana	85.7	90.4	4.9	17.3	21.8	12.9	43.1
Maine	80.2	78.6	3.8	23.1	25.8	10.5	36.9
Maryland	84.2	77.9	6.6	22.1	16.6	22.2	32.5
Massachusetts	79.8	76.7	9.0	21.8	14.5	17.4	37.4
Michigan	89.2	81.6	11.0	26.3	17.1	21.0	24.6
Minnesota	85.7	82.8	6.0	22.4	15.2	17.8	38.6
Mississippi	90.0	76.5	12.2	35.5	15.0	12.5	24.9
Missouri	88.1	88.3	8.8	34.2	16.1	17.0	23.9
Montana	94.8	94.0	4.9	20.5	17.5	16.7	40.5
Nebraska	81.0	75.2	4.8	23.6	12.8	19.2	39.7
Nevada	94.0	88.3	6.8	25.9	23.9	14.4	28.9
New Hampshire	75.0	64.1	1.7	21.9	16.4	25.6	34.5
New Jersey	94.2	95.1	2.8	14.9	18.5	21.0	42.9
New Mexico	79.2	NA	NA	NA	NA	NA	NA
New York	79.2	73.1	7.6	28.9	14.6	16.6	32.2
North Carolina	84.9	79.7	4.5	20.8	17.7	16.5	40.5
North Dakota	93.4	95.3	5.3	24.1	14.4	14.9	41.3
Ohio	84.1	83.3	6.1	19.4	18.8	19.6	36.1
Oklahoma	88.4	80.6	14.8	28.0	18.8	14.6	23.8
Oregon	69.9	74.3	3.8	33.7	13.8	15.5	33.1
Pennsylvania	95.0	96.5	3.6	19.4	20.8	15.0	41.1
Rhode Island	96.6	93.7	3.3	12.9	14.3	21.9	47.6
South Carolina	84.9	68.0	8.1	27.5	14.0	14.1	36.2
South Dakota	93.4	88.2	9.9	23.1	15.8	16.5	34.8
Tennessee	84.4	78.6	9.2	22.1	20.2	11.6	36.9
Texas	88.2	83.6	10.6	27.8	18.0	13.9	29.7
Utah	93.9	95.4	3.1	20.1	19.0	18.9	38.9
Vermont	69.3	60.6	5.9	24.1	16.7	20.5	32.9
Virginia	95.7	96.5	2.0	10.4	15.4	16.2	56.0
Washington	74.0	77.6	3.9	26.3	23.0	17.9	28.9
West Virginia	97.2	98.7	5.6	26.6	17.3	9.4	41.2
Wisconsin	91.8	88.8	6.4	19.3	14.1	13.3	46.9
Wyoming	96.1	82.2	9.3	16.6	5.5	17.3	51.2
State Median	85.7	82.2	5.9	24.1	17.0	16.2	36.1
State Range	25.6-98.8	29.2-98.7	1.7-21.0	10.4-35.5	5.5-25.8	9.4-25.6	14.9-56.7

TABLE 15. Percentage of Secondary Schools in Which All Staff Who Teach Health Education Topics Were Certified* in Health Education, the Lead Health Education Teacher Was Certified to Teach Health Education in Middle School or High School, and the Lead Health Education Teacher Had Experience Teaching Health Education Courses or Topics for a Specific Number of Years, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	All staff who teach health education topics are certified in health education	Lead health education teacher is certified to teach health education	Number of years lead health education teacher has taught health education courses or topics				
			1 year	2–5 years	6–9 years	10–14 years	≥15 years
LOCAL SURVEYS							
Albuquerque	100.0	96.9	6.2	22.0	12.5	15.6	43.7
Baltimore	58.6	48.1	9.8	28.0	11.3	18.4	32.6
Broward County	77.8	58.2	6.3	20.3	25.9	10.5	37.0
Charlotte-Mecklenberg County	94.0	64.4	3.5	14.4	23.3	21.8	37.0
Chicago	68.0	57.8	10.8	37.8	9.6	15.0	26.9
Detroit	81.5	74.5	9.8	15.0	20.5	25.6	29.1
District of Columbia	77.8	81.9	8.9	12.0	15.1	15.1	48.7
Fresno	71.4	44.6	12.5	24.9	12.5	17.0	33.2
Houston	62.7	64.0	12.3	22.4	12.5	16.0	36.8
Los Angeles	82.5	85.5	0.8	36.0	31.9	12.3	19.0
Memphis	93.4	81.7	5.1	21.7	18.7	7.3	47.2
Miami-Dade County	57.4	51.7	10.0	23.6	16.3	13.7	36.4
Milwaukee	62.5	43.7	26.6	20.8	10.0	12.6	30.0
New York City	49.5	41.7	13.2	47.1	14.0	10.9	14.8
Newark	100.0	82.3	13.2	34.1	23.4	6.8	22.5
Orange County	68.8	69.4	6.9	30.3	23.2	14.2	25.5
Philadelphia	91.1	93.9	9.9	22.4	13.7	10.7	43.4
San Diego	0.0	48.5	17.0	39.9	24.6	13.4	5.1
San Francisco	50.5	47.4	27.5	26.8	21.9	13.3	10.5
Local Median	71.4	64.0	9.9	23.6	16.3	13.7	32.9
Local Range	0.0-100.0	41.7-96.9	0.0-27.5	12.0-47.1	9.6-31.9	6.8-25.6	5.1-48.7
TERRITORIAL SURVEYS							
American Samoa	3.8	34.6	0.0	23.1	38.5	15.4	23.1
Guam	92.3	84.6	15.4	23.1	23.1	7.7	30.8
Marshall Islands	21.1	NA	NA	NA	NA	NA	NA
Northern Mariana Islands	42.9	28.6	42.9	28.6	28.6	0.0	0.0
Palau	0.0	9.1	18.2	18.2	36.4	18.2	9.1
Territorial Median	21.1	31.6	16.8	23.1	32.5	11.6	16.1
Territorial Range	0.0-92.3	9.1-84.6	0.0-42.9	18.2-28.6	23.1-38.5	0.0-18.2	0.0-30.8
TRIBAL SURVEYS							
Cherokee Nation	79.3	78.0	9.7	28.6	25.6	11.2	24.8
Nez Perce	100.0	100.0	0.0	0.0	0.0	14.3	85.7

NA=Data not available.

*Certification, licensure, or endorsement by the state.

TABLE 16a. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Alcohol- or other drug-use prevention	Asthma	Emotional and mental health	Foodborne illness prevention	HIV† prevention	Human sexuality	Injury prevention and safety
STATE SURVEYS							
Alabama	49.6	19.5	21.5	17.0	44.3	22.9	44.2
Alaska	30.7	11.2	27.7	24.6	17.8	9.9	32.3
Arizona	40.3	19.2	30.8	21.6	22.5	15.7	41.8
Arkansas	44.8	35.4	37.8	22.0	26.5	22.1	65.9
California	33.6	14.8	26.2	16.0	40.0	29.5	32.4
Connecticut	31.4	10.6	27.3	9.9	26.4	27.1	38.3
Delaware	45.9	18.3	37.7	13.1	53.2	54.8	31.7
Florida	39.0	23.5	33.6	25.8	46.4	37.1	47.9
Georgia	38.1	20.0	24.7	14.5	47.4	36.8	38.5
Hawaii	51.3	14.9	46.4	25.7	66.3	61.5	39.1
Idaho	59.2	11.4	46.8	14.8	45.9	43.5	39.5
Indiana	21.4	10.2	20.4	11.6	28.4	17.8	29.3
Iowa	26.5	9.1	24.3	21.0	33.6	28.0	27.5
Kansas	32.0	11.0	27.3	21.8	30.0	30.1	40.2
Kentucky	34.2	18.8	30.9	19.6	31.8	23.4	52.5
Louisiana	55.2	21.8	41.3	20.3	27.2	21.9	56.3
Maine	42.4	17.3	43.9	13.8	41.6	46.4	41.5
Maryland	54.3	19.0	52.3	18.0	66.5	58.8	43.7
Massachusetts	44.9	9.6	44.9	11.8	37.4	41.5	33.8
Michigan	38.5	14.8	33.3	18.9	54.4	47.2	27.6
Minnesota	43.4	16.2	67.5	18.8	41.1	42.7	45.1
Mississippi	43.1	31.1	43.0	32.1	29.3	24.3	51.8
Missouri	37.9	26.1	36.5	23.0	28.4	23.8	49.5
Montana	40.8	19.8	32.2	19.9	39.4	29.8	51.2
Nebraska	28.5	37.4	24.2	18.0	20.9	19.5	36.1
Nevada	60.7	16.8	46.9	23.6	49.0	49.9	40.3
New Hampshire	58.7	20.9	59.9	18.4	46.0	46.2	47.6
New Jersey	48.1	34.6	48.2	21.2	42.1	50.2	54.0
New York	60.0	24.8	52.1	23.5	55.6	48.0	49.6
North Carolina	43.1	36.2	35.5	23.4	45.6	41.0	58.8
North Dakota	39.6	11.1	33.2	25.8	31.9	25.7	38.5
Ohio	37.6	20.7	40.6	23.8	27.8	26.6	46.0
Oklahoma	58.3	42.3	50.6	44.4	65.5	31.0	70.4
Oregon	28.1	12.8	28.1	20.1	27.4	28.0	37.7
Pennsylvania	43.3	21.2	34.3	15.3	37.7	31.8	43.1
Rhode Island	27.5	13.6	39.9	18.7	24.6	24.7	43.0
South Carolina	38.1	19.6	32.2	22.0	49.5	48.7	51.4
South Dakota	31.1	8.5	25.8	13.3	15.3	15.6	26.6
Tennessee	48.9	42.7	52.8	29.7	44.2	28.2	65.6
Texas	47.1	24.2	36.2	21.2	39.5	36.1	59.3
Utah	59.0	15.6	47.4	17.7	52.2	64.4	38.4
Vermont	53.4	23.7	57.7	18.1	31.9	33.0	38.4
Virginia	42.6	27.7	29.7	20.7	34.0	30.7	44.3
Washington	34.7	20.1	29.9	28.6	52.1	41.0	41.9
West Virginia	36.0	20.0	26.5	20.6	28.8	23.7	39.0
Wisconsin	51.7	14.9	46.9	15.2	30.0	34.4	41.1
Wyoming	42.4	13.4	31.7	24.0	25.4	27.1	49.7
State Median	42.4	19.2	35.5	20.3	37.7	30.7	41.9
State Range	21.4-60.7	8.5-42.7	20.4-67.5	9.9-44.4	15.3-66.5	9.9-64.4	26.6-70.4

TABLE 16a. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Alcohol- or other drug-use prevention	Asthma	Emotional and mental health	Foodborne illness prevention	HIV† prevention	Human sexuality	Injury prevention and safety
LOCAL SURVEYS							
Albuquerque	46.8	21.9	34.2	28.0	58.0	68.7	34.3
Baltimore	38.4	32.7	42.3	19.6	45.5	36.4	43.3
Broward County	32.1	36.7	36.7	26.4	70.0	69.5	47.2
Charlotte-Mecklenberg County	43.8	45.2	31.9	23.8	69.5	62.3	51.3
Chicago	41.7	56.3	40.6	25.5	43.7	32.1	47.8
Detroit	51.4	53.5	43.1	32.1	61.4	50.8	47.0
District of Columbia	61.5	35.3	46.9	32.3	82.5	79.5	54.4
Fresno	26.0	19.8	18.3	7.7	63.3	52.7	32.0
Houston	63.9	63.0	44.1	33.5	74.0	49.3	81.1
Los Angeles	62.1	29.6	49.6	37.3	80.0	70.2	42.7
Memphis	67.1	56.8	48.3	40.4	74.7	42.4	70.2
Miami-Dade County	51.5	24.2	40.1	30.1	62.4	44.3	56.1
Milwaukee	40.4	53.6	40.7	25.9	48.1	48.1	43.2
New York City	49.8	35.5	51.2	35.0	63.9	47.8	54.1
Newark	57.6	61.3	52.1	29.9	80.5	71.0	62.0
Orange County	32.9	11.6	32.4	23.7	88.2	78.9	46.2
Philadelphia	47.6	54.3	31.2	24.7	61.2	42.7	56.2
San Diego	35.1	18.0	31.8	23.2	78.4	70.1	38.0
San Francisco	92.3	41.7	81.4	39.4	81.8	73.9	47.0
Local Median	47.6	36.7	40.7	28.0	69.5	52.7	47.2
Local Range	26.0-92.3	11.6-63.0	18.3-81.4	7.7-40.4	43.7-88.2	32.1-79.5	32.0-81.1
TERRITORIAL SURVEYS							
American Samoa	100.0	57.7	69.2	76.9	92.3	88.5	92.3
Guam	30.8	7.7	23.1	23.1	46.2	38.5	15.4
Northern Mariana Islands	87.5	37.5	62.5	37.5	100.0	75.0	87.5
Palau	63.6	0.0	100.0	27.3	100.0	63.6	100.0
Territorial Median	75.6	22.6	65.9	32.4	96.2	69.3	89.9
Territorial Range	30.8-100.0	0.0-57.7	23.1-100.0	23.1-76.9	46.2-100.0	38.5-88.5	15.4-100.0
TRIBAL SURVEYS							
Cherokee Nation	52.3	44.2	45.1	37.5	64.1	29.4	59.1
Nez Perce	57.1	14.3	14.3	0.0	42.9	57.1	28.6

*Such as workshops, conferences, continuing education, or any other kind of in-service.
 †Human immunodeficiency virus.

TABLE 16b. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD [†] prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
STATE SURVEYS							
Alabama	31.9	49.8	24.6	34.1	25.8	34.4	46.1
Alaska	23.6	29.6	10.2	13.2	24.2	21.8	36.5
Arizona	34.0	39.7	17.0	18.7	21.1	31.7	48.5
Arkansas	42.9	68.4	24.2	25.0	33.5	43.2	58.9
California	24.1	30.0	27.6	35.9	17.7	29.3	41.0
Connecticut	26.7	51.8	19.5	23.2	21.2	22.6	50.2
Delaware	38.3	60.7	44.3	51.6	18.0	35.6	50.0
Florida	40.3	53.8	33.5	40.0	30.9	36.3	69.1
Georgia	30.3	48.4	34.2	43.7	21.1	28.5	47.5
Hawaii	43.7	70.3	58.8	62.7	33.7	45.2	56.3
Idaho	39.6	55.6	28.4	39.8	38.8	40.2	58.4
Indiana	21.0	38.2	16.5	20.3	12.8	19.5	37.7
Iowa	31.6	33.3	26.2	30.6	17.1	17.3	42.9
Kansas	40.5	55.5	23.4	27.4	19.7	30.0	61.9
Kentucky	39.9	50.9	23.2	26.4	25.4	29.5	55.3
Louisiana	43.5	63.3	21.5	26.1	50.0	43.3	66.0
Maine	51.3	58.2	35.2	41.3	36.6	29.8	54.0
Maryland	43.9	57.7	45.9	55.5	48.9	45.0	61.9
Massachusetts	41.4	48.2	33.9	37.5	32.2	24.9	63.8
Michigan	43.4	48.9	39.4	48.1	24.8	29.7	42.5
Minnesota	38.5	51.0	32.3	36.0	40.5	31.7	57.5
Mississippi	47.5	53.6	27.8	29.2	43.1	41.7	54.7
Missouri	43.7	54.8	21.1	26.8	30.3	30.7	58.9
Montana	32.6	56.4	28.5	33.9	38.9	33.2	52.6
Nebraska	32.2	39.1	21.3	21.3	24.4	19.4	44.7
Nevada	56.4	56.5	39.6	48.2	49.3	49.8	64.4
New Hampshire	59.0	64.9	36.8	43.1	43.0	38.7	60.1
New Jersey	40.9	69.0	35.2	41.5	64.9	34.1	76.6
New York	53.5	60.8	42.4	47.6	38.2	44.6	63.5
North Carolina	42.9	72.5	36.7	40.7	25.1	43.7	63.7
North Dakota	40.7	54.6	17.8	27.0	31.0	33.2	51.9
Ohio	42.0	51.5	23.4	25.8	29.7	26.9	53.3
Oklahoma	45.2	56.8	32.7	43.4	39.4	51.3	76.9
Oregon	25.1	31.8	19.5	23.9	27.2	17.5	43.7
Pennsylvania	40.1	64.4	25.3	28.8	22.6	27.9	52.7
Rhode Island	38.4	44.6	20.2	22.1	19.0	19.0	68.6
South Carolina	38.6	69.3	40.8	46.4	22.5	33.4	60.3
South Dakota	24.6	44.9	10.5	16.2	23.5	25.7	50.3
Tennessee	49.9	72.4	26.3	31.5	73.1	40.9	80.4
Texas	42.5	65.2	43.4	40.9	27.4	35.5	57.8
Utah	42.7	53.4	40.2	53.0	30.4	44.0	49.9
Vermont	46.0	58.5	21.2	28.3	33.9	31.2	52.0
Virginia	38.6	75.9	26.0	29.9	32.5	32.4	64.0
Washington	37.2	47.0	34.8	44.5	34.7	28.4	47.3
West Virginia	40.4	56.1	21.3	24.7	18.5	40.7	51.9
Wisconsin	45.4	55.2	27.7	31.1	47.9	34.5	50.7
Wyoming	44.3	54.5	21.7	20.6	30.8	38.2	57.1
State Median	40.5	54.8	27.6	31.5	30.4	33.2	54.7
State Range	21.0-59.0	29.6-75.9	10.2-58.8	13.2-62.7	12.8-73.1	17.3-51.3	36.5-80.4

TABLE 16b. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific Health Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD† prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
LOCAL SURVEYS							
Albuquerque	37.5	49.9	53.1	59.4	46.8	40.6	50.0
Baltimore	45.9	58.3	33.5	39.0	23.0	40.6	57.4
Broward County	27.1	39.3	44.9	61.8	29.9	30.2	82.7
Charlotte-Mecklenberg County	38.6	84.1	53.5	61.9	31.7	45.3	68.9
Chicago	56.2	74.5	38.8	39.4	22.7	32.9	62.2
Detroit	58.9	76.4	41.3	54.2	33.5	46.1	59.7
District of Columbia	58.6	88.3	58.7	70.7	35.2	48.2	47.0
Fresno	18.3	23.7	39.6	67.2	34.3	13.0	47.3
Houston	68.9	91.2	61.7	64.1	52.3	59.5	66.3
Los Angeles	40.5	32.5	59.4	76.9	51.3	63.3	67.1
Memphis	56.8	84.9	46.1	69.2	64.3	50.5	75.1
Miami-Dade County	52.2	66.9	36.6	53.0	37.0	39.7	65.3
Milwaukee	38.1	54.6	45.4	48.1	54.9	33.8	56.1
New York City	52.3	64.8	47.9	53.8	38.8	45.6	59.1
Newark	55.2	88.5	70.2	74.5	71.4	36.1	77.5
Orange County	25.7	45.1	67.3	69.7	41.8	28.6	67.4
Philadelphia	60.3	82.8	27.4	39.5	29.0	30.1	60.9
San Diego	18.3	24.6	61.8	74.9	25.3	33.1	48.0
San Francisco	76.5	68.3	68.5	76.5	60.3	81.3	89.4
Local Median	52.2	66.9	47.9	61.9	37.0	40.6	62.2
Local Range	18.3-76.5	23.7-91.2	27.4-70.2	39.0-76.9	22.7-71.4	13.0-81.3	47.0-89.4
TERRITORIAL SURVEYS							
American Samoa	92.3	100.0	92.3	92.3	88.0	100.0	84.6
Guam	23.1	23.1	15.4	38.5	30.8	38.5	23.1
Northern Mariana Islands	87.5	87.5	62.5	87.5	75.0	75.0	75.0
Palau	100.0	100.0	63.6	100.0	72.7	72.7	72.7
Territorial Median	89.9	93.8	63.1	89.9	73.9	73.9	73.9
Territorial Range	23.1-100.0	23.1-100.0	15.4-92.3	38.5-100.0	30.8-88.0	38.5-100.0	23.1-84.6
TRIBAL SURVEYS							
Cherokee Nation	47.0	51.5	28.1	37.7	35.8	50.3	72.0
Nez Perce	57.1	85.7	28.6	28.6	14.3	14.3	71.4

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Sexually transmitted disease.

TABLE 17a. Percentage of Secondary Schools in Which the Lead Health Education Teacher Wanted to Receive Professional Development* on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Alcohol- or other drug-use prevention	Asthma	Emotional and mental health	Foodborne illness prevention	HIV† prevention	Human sexuality	Injury prevention and safety
STATE SURVEYS							
Alabama	86.3	69.0	73.1	63.1	71.5	59.9	68.4
Alaska	68.0	44.2	70.2	43.0	49.6	50.5	50.6
Arizona	68.6	58.2	71.2	48.2	58.2	54.0	63.8
Arkansas	81.4	69.3	72.3	57.6	66.9	61.6	78.6
California	66.7	55.2	71.3	51.9	56.7	57.6	59.8
Connecticut	80.4	52.6	79.8	42.7	72.7	77.2	53.9
Delaware	81.4	56.9	83.6	45.8	73.3	81.7	59.3
Florida	72.2	63.9	72.5	59.1	66.9	64.3	71.7
Georgia	73.1	53.9	66.1	55.3	71.6	62.5	68.3
Hawaii	80.1	74.6	87.3	71.9	68.7	73.2	73.9
Idaho	75.0	55.3	74.5	52.5	63.6	64.7	62.9
Indiana	71.4	41.7	61.7	35.3	58.0	59.1	48.8
Iowa	68.1	43.4	67.0	46.1	57.3	65.1	50.8
Kansas	73.5	50.3	67.9	46.8	66.8	68.3	59.2
Kentucky	73.2	54.0	73.4	47.4	64.2	59.7	64.6
Louisiana	75.8	69.7	69.7	59.4	64.8	54.1	76.8
Maine	70.6	41.6	71.5	46.2	58.0	67.2	57.0
Maryland	78.5	65.5	76.0	53.8	75.8	75.2	64.8
Massachusetts	84.7	55.5	83.6	52.8	75.0	79.6	63.7
Michigan	74.5	58.5	73.7	48.7	63.8	65.8	62.9
Minnesota	74.7	46.2	75.6	42.1	69.1	79.3	56.6
Mississippi	78.2	67.9	76.7	64.9	74.4	67.5	76.2
Missouri	74.8	52.3	67.6	48.4	61.0	58.9	63.5
Montana	70.2	58.6	65.6	46.8	66.3	65.5	62.8
Nebraska	64.1	42.5	59.9	37.0	48.4	52.6	44.5
Nevada	76.5	57.7	73.1	54.8	73.8	75.5	64.6
New Hampshire	76.1	48.6	79.2	50.2	64.7	75.1	65.5
New Jersey	81.4	63.3	81.8	60.8	78.1	81.1	68.8
New York	81.6	65.7	82.2	63.1	76.2	79.2	66.6
North Carolina	76.4	64.5	70.2	57.6	69.8	67.0	67.4
North Dakota	59.9	53.2	70.8	43.0	53.0	53.6	54.9
Ohio	78.6	51.1	76.1	50.2	67.6	66.5	59.6
Oklahoma	69.5	55.5	63.5	43.8	54.3	47.2	62.6
Oregon	67.7	47.8	68.9	40.8	62.2	66.7	47.0
Pennsylvania	79.7	58.4	76.2	56.4	79.3	80.7	61.9
Rhode Island	84.4	61.2	77.8	54.2	70.9	75.1	61.9
South Carolina	72.9	63.8	70.3	51.8	67.1	68.3	61.8
South Dakota	62.8	45.6	58.3	37.3	52.6	48.8	54.9
Tennessee	74.7	63.6	72.0	54.3	61.3	57.5	66.3
Texas	79.7	62.1	71.6	58.4	71.5	69.5	68.9
Utah	78.7	63.1	82.8	54.5	78.7	80.5	62.4
Vermont	61.2	37.7	67.6	35.4	51.5	61.1	44.2
Virginia	73.9	62.8	70.5	51.5	56.3	56.6	69.4
Washington	68.4	51.6	70.3	44.5	58.1	60.5	51.5
West Virginia	79.4	62.4	68.0	56.3	72.4	67.6	65.2
Wisconsin	72.0	50.6	71.7	42.6	65.8	72.9	51.3
Wyoming	66.8	60.9	70.6	52.9	62.5	59.9	59.5
State Median	74.7	56.9	71.6	51.5	66.3	65.8	62.8
State Range	59.9-86.3	37.7-74.6	58.3-87.3	35.3-71.9	48.4-79.3	47.2-81.7	44.2-78.6

TABLE 17a. Percentage of Secondary Schools in Which the Lead Health Education Teacher Wanted to Receive Professional Development* on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Alcohol- or other drug-use prevention	Asthma	Emotional and mental health	Foodborne illness prevention	HIV† prevention	Human sexuality	Injury prevention and safety
LOCAL SURVEYS							
Albuquerque	84.4	59.3	78.1	65.6	71.9	81.2	67.7
Baltimore	73.1	73.7	75.5	68.9	74.1	65.0	67.6
Broward County	68.3	68.2	69.2	62.5	59.4	64.6	60.5
Charlotte-Mecklenberg County	67.2	69.6	66.2	62.7	64.8	65.1	55.4
Chicago	80.3	76.1	82.7	65.6	70.1	70.5	78.3
Detroit	76.2	83.2	83.1	70.1	73.7	74.1	77.0
District of Columbia	79.0	82.4	88.4	70.8	66.6	72.8	76.5
Fresno	54.7	44.6	73.4	42.8	54.7	54.7	46.8
Houston	93.9	90.2	90.1	86.3	91.4	91.5	88.9
Los Angeles	83.4	71.3	88.0	74.7	79.8	80.0	75.9
Memphis	75.2	76.1	84.1	70.8	69.3	78.4	78.1
Miami-Dade County	69.2	67.4	70.8	63.2	68.5	68.3	73.5
Milwaukee	70.5	54.1	74.2	59.0	57.0	57.9	58.7
New York City	74.9	72.0	76.5	66.7	66.7	73.5	69.4
Newark	79.0	72.2	85.3	65.7	83.5	82.8	73.9
Orange County	78.4	62.0	73.5	54.6	69.0	71.3	61.7
Philadelphia	77.0	74.0	80.4	73.5	76.8	81.0	75.1
San Diego	53.1	44.2	58.3	36.9	40.1	38.2	44.5
San Francisco	76.1	57.9	89.1	59.4	78.9	76.0	62.9
Local Median	76.1	71.3	78.1	65.6	69.3	72.8	69.4
Local Range	53.1-93.9	44.2-90.2	58.3-90.1	36.9-86.3	40.1-91.4	38.2-91.5	44.5-88.9
TERRITORIAL SURVEYS							
American Samoa	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Guam	76.9	100.0	84.6	92.3	76.9	84.6	76.9
Northern Mariana Islands	75.0	75.0	75.0	75.0	75.0	75.0	75.0
Palau	27.3	81.8	36.4	63.6	63.6	72.7	27.3
Territorial Median	76.0	90.9	79.8	83.7	76.0	79.8	76.0
Territorial Range	27.3-100.0	75.0-100.0	36.4-100.0	63.6-100.0	63.6-100.0	72.7-100.0	27.3-100.0
TRIBAL SURVEYS							
Cherokee Nation	66.2	61.0	69.2	52.3	62.4	54.2	67.7
Nez Perce	85.7	28.6	71.4	42.9	71.4	71.4	57.1

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Human immunodeficiency virus.

TABLE 17b. Percentage of Secondary Schools in Which the Lead Health Education Teacher Wanted to Receive Professional Development* on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD [†] prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
STATE SURVEYS							
Alabama	72.7	73.6	62.2	67.2	79.3	74.7	73.6
Alaska	62.1	58.3	50.1	53.3	68.2	58.8	72.6
Arizona	71.8	66.7	56.1	55.0	69.5	62.2	76.0
Arkansas	78.3	84.3	66.5	71.2	75.8	77.3	79.1
California	71.2	62.7	54.4	56.8	70.8	60.9	76.8
Connecticut	72.4	62.8	70.8	74.5	79.9	69.1	79.9
Delaware	79.7	68.4	76.7	73.3	78.3	67.8	73.8
Florida	75.9	73.2	61.9	65.8	73.8	66.1	78.0
Georgia	74.7	75.6	68.1	72.9	68.8	70.5	72.1
Hawaii	85.1	75.5	72.2	71.0	84.1	74.7	87.8
Idaho	72.1	66.9	59.4	65.7	76.4	69.0	80.2
Indiana	63.9	65.0	59.1	62.8	70.0	60.3	75.7
Iowa	67.2	62.9	60.8	64.3	74.3	58.9	70.7
Kansas	72.9	72.8	66.7	68.0	73.5	69.0	73.3
Kentucky	74.6	73.2	60.7	63.6	71.5	69.7	71.7
Louisiana	73.9	74.7	59.1	64.2	72.5	70.6	78.3
Maine	71.0	61.6	64.2	63.6	68.0	58.2	72.8
Maryland	77.0	65.0	70.9	74.8	77.7	62.8	83.2
Massachusetts	78.4	69.6	74.5	78.6	84.5	68.5	90.3
Michigan	72.5	69.7	60.0	62.2	73.8	65.2	77.6
Minnesota	70.9	67.4	68.5	71.4	77.8	64.2	72.7
Mississippi	80.2	80.3	75.6	73.7	79.2	73.9	83.0
Missouri	73.2	73.3	63.1	64.5	73.3	69.6	77.3
Montana	76.5	74.2	58.1	62.6	70.3	61.3	75.0
Nebraska	63.1	59.8	53.0	56.8	66.2	53.8	67.9
Nevada	76.5	68.8	75.7	75.3	75.4	70.0	77.5
New Hampshire	78.4	68.4	65.6	71.5	80.6	64.0	76.9
New Jersey	80.1	76.0	75.6	78.5	84.2	74.3	83.2
New York	80.0	74.5	75.1	76.3	84.8	72.5	81.2
North Carolina	78.0	77.9	68.7	69.4	77.9	68.3	80.0
North Dakota	58.8	55.8	49.6	53.3	71.5	53.1	65.8
Ohio	72.0	68.6	69.1	73.2	78.6	67.7	77.5
Oklahoma	64.5	67.0	51.9	54.8	65.7	63.5	72.7
Oregon	62.6	58.2	62.2	64.8	69.6	60.4	72.2
Pennsylvania	76.0	74.0	77.4	79.4	80.9	71.8	76.3
Rhode Island	72.6	71.9	73.3	70.1	76.1	64.6	78.6
South Carolina	73.6	69.1	67.1	67.0	74.8	68.0	77.2
South Dakota	64.3	66.8	53.0	57.7	68.7	58.7	69.5
Tennessee	74.9	70.7	60.5	62.5	71.9	65.1	78.9
Texas	78.5	73.0	69.9	69.4	79.5	72.7	83.9
Utah	76.1	66.2	74.3	78.9	82.7	72.1	79.5
Vermont	61.0	46.7	54.1	55.0	65.8	47.6	60.8
Virginia	80.0	76.5	54.2	56.9	68.9	65.6	75.0
Washington	67.8	62.6	57.1	59.4	66.7	61.0	74.0
West Virginia	73.1	67.4	67.6	71.3	79.6	66.8	75.9
Wisconsin	72.5	63.9	70.4	73.5	74.0	62.6	72.5
Wyoming	71.3	62.9	60.6	62.5	69.8	65.6	68.1
State Median	73.1	68.6	64.2	67.0	74.0	66.1	76.3
State Range	58.8-85.1	46.7-84.3	49.6-77.4	53.3-79.4	65.7-84.8	47.6-77.3	60.8-90.3

TABLE 17b. Percentage of Secondary Schools in Which the Lead Health Education Teacher Wanted to Receive Professional Development* on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Nutrition and dietary behavior	Physical activity and fitness	Pregnancy prevention	STD† prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
LOCAL SURVEYS							
Albuquerque	87.5	71.8	75.0	74.2	84.3	78.1	87.5
Baltimore	75.7	72.7	68.7	72.4	75.2	68.0	84.2
Broward County	76.0	63.2	58.4	66.0	71.5	60.2	64.2
Charlotte-Mecklenberg County	74.1	66.2	65.1	57.7	72.6	59.2	76.4
Chicago	80.4	81.4	67.0	71.2	80.5	76.0	85.6
Detroit	80.9	85.7	76.1	75.2	79.7	68.7	82.0
District of Columbia	85.5	78.8	69.8	66.6	82.4	73.7	85.4
Fresno	54.8	37.8	54.7	54.7	66.4	52.5	60.6
Houston	87.5	88.9	93.8	92.6	95.0	91.4	89.9
Los Angeles	84.1	76.9	79.8	75.7	92.0	72.0	90.5
Memphis	85.6	79.7	74.3	75.4	79.1	62.0	82.3
Miami-Dade County	76.6	72.8	63.7	68.4	72.1	64.7	76.6
Milwaukee	68.2	66.8	58.7	53.4	68.2	57.1	74.2
New York City	76.3	78.8	68.0	69.5	77.2	73.0	79.6
Newark	83.4	74.6	78.6	79.4	83.1	81.2	89.9
Orange County	76.0	69.0	63.3	66.6	76.1	68.9	85.6
Philadelphia	75.3	78.8	81.0	82.9	86.8	73.3	86.6
San Diego	56.8	49.8	42.1	42.0	61.7	51.5	59.8
San Francisco	76.0	60.0	73.4	73.4	81.3	70.9	86.5
Local Median	76.3	72.8	68.7	71.2	79.1	68.9	84.2
Local Range	54.8-87.5	37.8-88.9	42.1-93.8	42.0-92.6	61.7-95.0	51.5-91.4	59.8-90.5
TERRITORIAL SURVEYS							
American Samoa	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Guam	69.2	76.9	84.6	76.9	84.6	69.2	92.3
Northern Mariana Islands	75.0	75.0	75.0	75.0	75.0	75.0	62.5
Palau	27.3	18.2	72.7	63.6	27.3	18.2	54.5
Territorial Median	72.1	76.0	79.8	76.0	79.8	72.1	77.4
Territorial Range	27.3-100.0	18.2-100.0	72.7-100.0	63.6-100.0	27.3-100.0	18.2-100.0	54.5-100.0
TRIBAL SURVEYS							
Cherokee Nation	70.9	73.0	49.3	54.9	61.0	66.2	74.4
Nez Perce	71.4	57.1	42.9	71.4	57.1	71.4	85.7

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Sexually transmitted disease.

TABLE 18. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Critical HIV[†] and STD[‡] Prevention Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Describing how widespread HIV and other STD infections are and the consequences of these infections	Understanding the modes of transmission and effective prevention strategies	Identifying populations of youth who are at high risk of being infected	Implementing health education strategies using prevention messages that are likely to be effective in reaching youth	All 4 topics
STATE SURVEYS					
Alabama	43.1	25.8	27.9	28.1	17.4
Alaska	18.5	19.5	13.7	13.9	7.3
Arizona	14.9	17.7	14.9	21.5	9.8
Arkansas	24.1	25.6	22.1	34.5	18.7
California	38.6	39.4	35.8	34.3	28.6
Connecticut	21.7	23.0	16.0	23.8	12.5
Delaware	50.8	53.2	50.8	59.0	44.3
Florida	43.3	43.6	37.6	37.1	27.9
Georgia	41.2	38.8	34.2	33.4	25.9
Hawaii	64.2	62.7	57.1	51.6	48.4
Idaho	45.7	46.2	38.9	44.2	29.3
Indiana	27.8	28.1	22.6	21.5	14.4
Iowa	29.3	31.0	27.5	27.1	18.1
Kansas	23.5	24.6	20.7	25.0	15.2
Kentucky	24.9	27.4	22.0	26.7	14.2
Louisiana	21.9	25.5	20.3	29.0	14.6
Maine	40.6	43.8	36.4	44.4	27.4
Maryland	64.7	65.1	60.0	55.3	47.5
Massachusetts	34.6	35.2	32.6	38.1	24.2
Michigan	50.2	53.3	46.5	43.0	36.3
Minnesota	31.7	33.7	24.6	32.3	16.3
Mississippi	25.8	24.2	25.7	33.0	20.7
Missouri	25.2	25.8	20.1	31.4	16.1
Montana	33.5	34.9	31.3	36.9	25.6
Nebraska	17.0	19.0	15.6	25.2	12.3
Nevada	58.6	59.0	51.4	49.5	40.9
New Hampshire	40.3	43.5	35.3	47.7	28.5
New Jersey	37.5	39.2	33.5	44.7	28.2
New York	52.7	52.9	45.3	51.3	38.0
North Carolina	40.2	41.3	35.0	41.0	26.5
North Dakota	28.2	28.0	26.5	36.4	20.8
Ohio	26.7	27.9	24.7	29.6	19.3
Oklahoma	56.2	60.2	45.1	43.0	33.4
Oregon	22.9	23.9	17.5	26.8	14.1
Pennsylvania	31.6	31.6	30.0	34.1	22.8
Rhode Island	24.5	25.6	26.1	28.7	16.9
South Carolina	43.6	46.5	40.8	42.0	32.1
South Dakota	9.7	9.3	9.5	17.0	7.0
Tennessee	40.4	40.0	31.5	37.4	24.8
Texas	32.9	35.8	31.0	37.4	24.4
Utah	49.5	47.4	38.5	43.8	30.3
Vermont	33.6	36.7	31.6	41.8	22.1
Virginia	27.1	28.8	25.8	33.8	19.3
Washington	47.0	49.7	42.2	37.3	31.6
West Virginia	25.3	26.3	23.8	27.7	19.8
Wisconsin	28.5	31.6	24.9	33.3	17.7
Wyoming	16.7	18.5	15.9	27.1	12.1
State Median	32.9	33.7	30.0	34.3	22.1
State Range	9.7-64.7	9.3-65.1	9.5-60.0	13.9-59.0	7.0-48.4

TABLE 18. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Critical HIV[†] and STD[‡] Prevention Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Describing how widespread HIV and other STD infections are and the consequences of these infections	Understanding the modes of transmission and effective prevention strategies	Identifying populations of youth who are at high risk of being infected	Implementing health education strategies using prevention messages that are likely to be effective in reaching youth	All 4 topics
LOCAL SURVEYS					
Albuquerque	56.2	46.8	34.3	50.0	28.0
Baltimore	48.6	47.4	42.2	43.1	35.6
Broward County	71.5	72.9	71.0	60.2	57.1
Charlotte-Mecklenberg County	57.9	59.7	54.0	55.9	44.6
Chicago	42.5	42.5	41.4	44.5	35.2
Detroit	57.9	57.8	53.1	46.1	38.8
District of Columbia	78.0	81.1	77.9	78.1	71.7
Fresno	61.8	65.7	55.0	45.9	40.5
Houston	73.9	73.9	67.4	67.4	58.6
Los Angeles	79.3	80.9	77.7	72.7	68.7
Memphis	74.7	71.4	74.6	64.1	53.8
Miami-Dade County	64.1	66.0	58.7	51.6	47.0
Milwaukee	45.2	48.4	44.6	46.8	29.8
New York City	61.3	60.9	56.8	56.6	49.9
Newark	78.6	78.6	78.6	71.5	71.5
Orange County	79.0	79.0	76.5	69.5	67.3
Philadelphia	54.4	51.5	50.8	47.8	41.4
San Diego	94.9	94.9	94.9	94.9	94.9
San Francisco	76.5	76.5	73.6	73.6	68.3
Local Median	64.1	66.0	58.7	56.6	49.9
Local Range	42.5-94.9	42.5-94.9	34.3-94.9	43.1-94.9	28.0-94.9
TERRITORIAL SURVEYS					
American Samoa	100.0	100.0	92.3	92.3	84.6
Guam	38.5	38.5	38.5	38.5	30.8
Northern Mariana Islands	100.0	100.0	100.0	87.5	87.5
Palau	90.9	90.9	90.9	81.8	72.7
Territorial Median	95.5	95.5	91.6	84.7	78.7
Territorial Range	38.5-100.0	38.5-100.0	38.5-100.0	38.5-92.3	30.8-87.5
TRIBAL SURVEYS					
Cherokee Nation	60.6	60.3	54.2	52.1	43.2
Nez Perce	28.6	28.6	28.6	57.1	14.3

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Human immunodeficiency virus.

‡Sexually transmitted disease.

TABLE 19a. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific HIV† Prevention Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Teaching HIV prevention education to students with physical, medical, or cognitive disabilities	Teaching HIV prevention education to students of various cultural backgrounds	Using interactive teaching methods for HIV prevention education†	Teaching essential skills for health behavior change related to HIV prevention and guiding student practice of these skills	Teaching about health-promoting social norms and beliefs related to HIV prevention	Strategies for involving parents, families, and others in student learning of HIV prevention education	Assessing students' performance in HIV prevention education
STATE SURVEYS							
Alabama	22.2	20.4	20.0	22.9	26.8	19.4	21.5
Alaska	6.8	9.2	6.2	8.8	9.6	6.3	7.3
Arizona	9.5	12.0	12.0	11.6	13.3	7.5	8.5
Arkansas	14.1	18.8	19.1	21.2	20.0	17.5	18.7
California	24.0	28.3	28.7	32.3	32.4	19.1	20.4
Connecticut	9.8	10.9	17.1	13.5	14.9	7.0	13.3
Delaware	36.1	32.2	46.8	47.5	42.6	27.8	33.8
Florida	20.5	27.1	33.0	35.2	31.5	24.5	29.2
Georgia	20.7	26.4	33.7	31.9	29.0	21.6	25.1
Hawaii	41.4	46.0	53.8	52.7	50.0	35.7	49.3
Idaho	20.0	22.8	36.7	32.4	33.2	21.1	25.9
Indiana	12.1	15.1	19.5	21.0	20.0	12.7	12.5
Iowa	10.4	15.9	23.2	20.9	20.2	10.9	15.8
Kansas	11.9	12.3	16.9	18.8	17.1	12.0	15.1
Kentucky	14.3	15.7	21.8	20.8	19.0	12.6	14.6
Louisiana	15.9	17.2	17.1	19.7	19.2	13.5	16.7
Maine	19.5	17.4	31.7	35.3	34.1	18.1	23.9
Maryland	34.8	39.5	43.8	47.1	44.4	28.4	42.7
Massachusetts	13.8	20.3	26.4	28.8	26.9	13.1	19.3
Michigan	26.6	30.1	41.1	41.7	40.6	30.1	37.6
Minnesota	12.4	14.0	22.8	21.4	23.1	10.9	15.1
Mississippi	17.9	20.9	21.4	23.9	22.5	19.5	18.3
Missouri	14.0	15.6	19.8	21.9	18.9	11.5	16.6
Montana	15.7	15.3	30.3	30.3	25.3	17.3	26.2
Nebraska	8.2	11.1	15.8	14.7	13.4	8.6	8.5
Nevada	34.7	39.1	44.2	49.6	46.3	32.1	39.6
New Hampshire	16.9	10.4	31.4	35.6	32.2	15.9	26.2
New Jersey	25.1	24.7	34.1	34.3	33.3	20.2	25.2
New York	31.5	35.4	44.4	45.5	42.7	30.8	34.4
North Carolina	20.1	26.4	30.8	33.1	29.7	21.2	23.0
North Dakota	16.5	16.5	24.1	25.7	25.2	15.8	19.2
Ohio	15.8	16.1	19.2	21.4	20.7	16.6	17.6
Oklahoma	28.0	34.1	35.0	39.3	40.3	32.2	29.4
Oregon	11.1	14.9	19.9	19.2	19.4	12.3	14.9
Pennsylvania	18.9	21.3	21.5	25.8	24.2	16.7	20.7
Rhode Island	8.4	19.3	14.7	18.0	18.2	9.5	11.6
South Carolina	25.1	30.5	33.3	37.4	35.9	27.0	31.2
South Dakota	5.9	6.7	6.2	8.3	8.7	5.1	6.1
Tennessee	20.6	22.6	27.7	29.9	30.6	24.7	23.9
Texas	18.9	25.9	24.8	25.7	28.1	19.8	21.4
Utah	22.4	24.4	33.4	37.5	33.6	22.1	21.9
Vermont	12.6	8.0	29.6	29.2	24.9	16.7	21.5
Virginia	17.2	18.8	24.1	23.1	22.1	13.8	16.5
Washington	25.5	29.7	35.2	35.3	34.4	25.2	31.1
West Virginia	17.2	17.8	19.9	22.3	20.4	13.8	19.6
Wisconsin	12.3	15.1	20.0	23.0	22.9	13.2	16.2
Wyoming	6.2	5.1	12.1	15.7	15.6	6.2	15.7
State Median	17.2	18.8	24.1	25.7	25.2	16.7	20.4
State Range	5.9-41.4	5.1-46.0	6.2-53.8	8.3-52.7	8.7-50.0	5.1-35.7	6.1-49.3

TABLE 19a. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific HIV† Prevention Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Teaching HIV prevention education to students with physical, medical, or cognitive disabilities	Teaching HIV prevention education to students of various cultural backgrounds	Using interactive teaching methods for HIV prevention education‡	Teaching essential skills for health behavior change related to HIV prevention and guiding student practice of these skills	Teaching about health-promoting social norms and beliefs related to HIV prevention	Strategies for involving parents, families, and others in student learning of HIV prevention education	Assessing students' performance in HIV prevention education
LOCAL SURVEYS							
Albuquerque	31.2	31.2	43.7	40.5	34.3	24.9	32.2
Baltimore	33.1	34.8	32.0	37.1	38.9	25.4	26.8
Broward County	40.7	58.5	61.9	63.6	62.8	44.5	58.8
Charlotte-Mecklenberg County	30.1	39.3	50.8	48.2	49.3	28.1	37.4
Chicago	30.1	32.2	32.1	33.1	33.5	24.8	30.2
Detroit	36.4	41.3	42.5	50.7	43.7	34.4	37.5
District of Columbia	56.0	65.4	77.9	74.8	62.3	57.8	68.7
Fresno	16.9	55.0	40.5	60.4	36.7	13.0	36.7
Houston	58.3	62.1	72.7	67.4	61.4	45.0	53.9
Los Angeles	52.9	66.4	71.2	71.2	71.8	46.1	58.9
Memphis	46.5	53.4	58.6	62.1	63.7	53.7	50.4
Miami-Dade County	38.5	46.8	48.7	53.2	48.2	42.2	47.3
Milwaukee	23.2	39.9	41.5	41.5	43.4	24.7	34.5
New York City	47.6	54.9	54.9	54.9	55.5	48.4	50.3
Newark	49.0	62.5	74.1	74.1	71.5	54.2	62.8
Orange County	50.9	62.6	72.0	76.7	69.6	55.6	62.6
Philadelphia	26.4	35.3	39.1	41.6	42.5	27.1	35.3
San Diego	94.9	94.9	94.9	94.9	94.9	93.2	1.6
San Francisco	60.5	70.9	65.8	73.8	68.5	52.5	60.5
Local Median	40.7	54.9	54.9	60.4	55.5	44.5	47.3
Local Range	16.9-94.9	31.2-94.9	32.0-94.9	33.1-94.9	33.5-94.9	13.0-93.2	1.6-68.7
TERRITORIAL SURVEYS							
American Samoa	65.4	76.9	84.6	80.8	80.8	76.9	73.1
Guam	23.1	38.5	30.8	30.8	30.8	30.8	23.1
Northern Mariana Islands	75.0	87.5	100.0	87.5	100.0	75.0	75.0
Palau	54.5	90.9	90.9	90.9	90.9	63.6	72.7
Territorial Median	60.0	82.2	87.8	84.2	85.9	69.3	72.9
Territorial Range	23.1-75.0	38.5-90.9	30.8-100.0	30.8-90.9	30.8-100.0	30.8-76.9	23.1-75.0
TRIBAL SURVEYS							
Cherokee Nation	32.3	42.5	45.6	45.1	44.8	34.8	35.2
Nez Perce	0.0	0.0	28.6	0.0	0.0	14.3	14.3

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Human immunodeficiency virus.

‡Such as role plays or cooperative group activities.

TABLE 19b. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific HIV† Prevention Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Implementing standards-based HIV prevention education curricula and student assessment	Using technology to improve HIV prevention education instruction	Teaching HIV prevention education to students with limited English proficiency	Addressing community concerns and challenges related to HIV prevention education	At least 6 of these 11 topics
STATE SURVEYS					
Alabama	18.6	21.9	14.1	15.6	17.3
Alaska	6.8	6.3	5.7	6.8	6.8
Arizona	9.5	8.5	8.2	6.5	9.2
Arkansas	20.7	22.2	11.7	15.2	18.7
California	28.9	18.9	19.4	19.4	25.6
Connecticut	13.2	10.7	6.5	6.2	8.0
Delaware	36.7	36.0	18.0	20.0	31.1
Florida	31.1	29.2	20.2	22.6	27.8
Georgia	26.6	20.3	15.6	19.2	24.3
Hawaii	56.3	46.6	36.2	35.3	49.4
Idaho	28.9	24.4	14.3	18.2	24.0
Indiana	16.3	12.6	7.6	10.4	13.1
Iowa	19.1	18.7	8.9	11.7	15.6
Kansas	13.8	15.0	9.0	7.6	12.4
Kentucky	15.5	15.2	9.7	10.0	14.1
Louisiana	17.1	15.3	10.9	13.8	16.7
Maine	25.6	23.2	8.8	13.1	20.5
Maryland	44.8	39.5	23.9	31.8	39.1
Massachusetts	23.0	18.6	12.4	12.2	17.9
Michigan	39.6	30.3	13.9	26.6	33.3
Minnesota	17.5	18.5	6.3	7.8	12.1
Mississippi	20.0	19.4	13.4	15.6	19.6
Missouri	15.9	18.1	11.2	12.9	15.9
Montana	20.2	23.9	7.9	12.6	20.5
Nebraska	11.0	13.3	6.2	8.7	8.1
Nevada	46.0	43.6	33.4	36.8	36.6
New Hampshire	24.3	23.4	5.4	7.1	15.9
New Jersey	29.0	28.6	16.4	17.8	24.5
New York	39.8	35.2	24.4	27.8	34.7
North Carolina	28.2	25.1	15.9	18.3	23.3
North Dakota	18.9	17.9	7.4	12.3	16.0
Ohio	17.1	17.2	10.6	13.0	14.8
Oklahoma	28.8	30.8	20.3	26.8	30.3
Oregon	20.0	13.6	9.9	11.9	14.1
Pennsylvania	23.8	22.8	14.0	15.9	19.3
Rhode Island	16.2	14.6	8.3	11.6	12.6
South Carolina	34.5	33.8	20.0	22.5	29.8
South Dakota	5.1	6.3	4.7	6.3	5.1
Tennessee	27.0	24.3	16.5	22.1	23.3
Texas	23.6	21.0	17.4	16.6	22.2
Utah	25.7	21.4	17.7	15.0	22.1
Vermont	27.8	18.1	3.6	11.6	18.7
Virginia	20.0	21.2	11.9	13.3	17.0
Washington	35.6	24.5	18.0	19.4	27.8
West Virginia	20.2	17.0	11.0	13.3	17.7
Wisconsin	19.2	17.0	8.1	8.7	15.3
Wyoming	15.8	7.8	3.5	5.2	10.4
State Median	20.7	20.3	11.7	13.3	18.7
State Range	5.1-56.3	6.3-46.6	3.5-36.2	5.2-36.8	5.1-49.4

TABLE 19b. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Specific HIV† Prevention Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Implementing standards-based HIV prevention education curricula and student assessment	Using technology to improve HIV prevention education instruction	Teaching HIV prevention education to students with limited English proficiency	Addressing community concerns and challenges related to HIV prevention education	At least 6 of these 11 topics
LOCAL SURVEYS					
Albuquerque	46.8	18.6	31.1	21.8	31.2
Baltimore	26.2	29.7	20.3	24.6	32.3
Broward County	58.8	60.9	50.7	40.7	58.8
Charlotte-Mecklenberg County	45.2	38.9	26.5	30.3	38.7
Chicago	28.4	26.2	22.7	23.8	28.0
Detroit	39.9	36.4	30.4	32.1	40.1
District of Columbia	70.9	59.3	40.4	59.3	65.5
Fresno	42.0	36.7	22.2	7.7	26.0
Houston	62.0	61.0	41.1	46.0	63.4
Los Angeles	68.6	55.3	53.5	51.4	66.5
Memphis	57.5	56.0	36.9	48.7	55.2
Miami-Dade County	45.6	47.9	35.8	40.3	46.3
Milwaukee	38.0	28.5	18.1	25.0	31.6
New York City	51.1	41.9	43.0	42.8	50.1
Newark	62.8	53.2	42.9	55.7	61.8
Orange County	72.0	67.3	50.8	57.8	62.5
Philadelphia	43.2	32.0	20.7	28.8	31.4
San Diego	94.9	1.7	0.0	94.9	94.9
San Francisco	73.9	60.5	52.5	57.9	63.0
Local Median	51.1	41.9	35.8	40.7	50.1
Local Range	26.2-94.9	1.7-67.3	0.0-53.5	7.7-94.9	26.0-94.9
TERRITORIAL SURVEYS					
American Samoa	69.2	57.7	73.1	65.4	88.5
Guam	23.1	30.8	23.1	30.8	23.1
Northern Mariana Islands	87.5	87.5	75.0	75.0	87.5
Palau	63.6	60.0	81.8	72.7	81.8
Territorial Median	66.4	58.9	74.1	69.1	84.7
Territorial Range	23.1-87.5	30.8-87.5	23.1-81.8	30.8-75.0	23.1-88.5
TRIBAL SURVEYS					
Cherokee Nation	35.2	35.9	27.7	25.5	34.9
Nez Perce	14.3	14.3	0.0	0.0	0.0

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Human immunodeficiency virus.

TABLE 20. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Teaching Methods, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Teaching students with physical, medical, or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Teaching students of different sexual orientations or gender identities	Using interactive teaching methods ^f	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
STATE SURVEYS									
Alabama	49.7	45.1	34.7	20.1	49.8	35.7	45.4	66.9	27.8
Alaska	46.8	55.2	39.4	11.5	58.2	38.6	42.2	59.0	14.7
Arizona	47.3	53.8	68.1	10.3	56.6	42.6	44.8	58.2	26.7
Arkansas	56.2	56.6	33.2	13.6	67.4	71.4	54.8	83.9	43.9
California	41.0	55.3	64.7	17.1	53.6	33.0	38.3	51.4	23.4
Connecticut	43.6	31.0	14.7	10.2	48.6	23.9	40.4	51.4	33.4
Delaware	26.2	36.1	13.1	13.1	70.5	37.7	60.6	54.1	44.2
Florida	49.0	53.2	51.7	17.2	63.4	41.4	51.8	67.7	35.5
Georgia	38.4	40.3	29.1	9.2	51.6	34.9	36.6	53.0	36.5
Hawaii	36.0	49.0	39.0	25.6	62.4	31.0	42.4	49.1	54.4
Idaho	33.1	40.7	30.4	9.8	57.0	40.2	52.1	54.2	35.4
Indiana	34.1	28.3	21.9	6.6	42.5	28.3	35.4	50.8	21.4
Iowa	41.1	28.9	18.5	11.8	53.5	30.0	44.9	56.5	23.3
Kansas	36.2	31.7	23.2	9.9	57.1	36.5	44.0	59.0	27.4
Kentucky	42.1	37.7	22.3	8.5	56.3	40.0	47.0	65.5	39.1
Louisiana	39.1	34.5	18.1	9.6	56.4	46.4	48.9	67.5	42.6
Maine	43.1	14.0	11.2	16.1	51.7	26.5	42.0	45.4	36.7
Maryland	61.3	51.7	33.1	17.5	66.1	38.4	51.0	67.6	50.9
Massachusetts	37.0	30.5	30.6	17.9	47.5	25.5	37.8	46.6	33.2
Michigan	30.5	24.0	10.0	9.7	50.2	29.9	38.9	47.8	32.4
Minnesota	57.2	47.7	30.6	14.0	52.2	30.7	51.4	59.7	34.4
Mississippi	38.3	39.8	22.9	11.8	58.8	48.0	50.3	72.2	36.4
Missouri	48.0	38.4	19.5	8.8	61.1	40.3	50.0	64.9	37.8
Montana	23.0	33.0	8.2	8.1	49.3	32.8	38.3	50.8	28.2
Nebraska	39.4	29.3	14.9	10.4	44.9	28.8	43.7	53.4	25.6
Nevada	40.1	52.2	45.0	8.6	57.3	36.2	44.9	60.3	30.8
New Hampshire	53.0	15.5	9.3	20.2	53.7	32.9	48.4	58.5	47.0
New Jersey	52.6	31.8	21.2	21.3	63.2	32.1	48.7	68.4	41.8
New York	41.1	32.5	24.6	22.4	57.1	34.6	47.9	59.9	41.6
North Carolina	43.9	49.4	35.2	12.5	64.0	40.1	44.1	68.7	42.0
North Dakota	41.1	31.3	14.1	7.5	50.8	38.0	48.6	61.3	29.4
Ohio	38.6	33.2	17.6	11.8	44.1	31.2	45.4	52.7	29.3
Oklahoma	57.2	54.9	28.9	15.4	57.4	53.3	56.5	70.0	35.0
Oregon	33.3	42.4	42.1	9.5	48.8	26.3	41.9	52.3	23.9
Pennsylvania	49.5	36.9	25.3	14.3	56.1	27.5	41.4	58.3	36.3
Rhode Island	31.0	19.1	16.9	12.6	35.2	27.8	37.0	38.9	31.0
South Carolina	38.3	43.0	30.8	16.7	59.3	38.6	46.3	66.6	43.8
South Dakota	34.6	25.3	8.5	5.2	37.1	29.0	28.1	48.4	20.4
Tennessee	53.9	41.9	21.3	10.0	64.5	56.6	56.7	76.8	41.2
Texas	50.2	50.7	49.0	13.8	62.0	39.6	51.9	73.4	42.7
Utah	21.8	32.9	34.7	6.9	48.2	24.0	35.6	49.3	30.7
Vermont	40.0	17.4	7.7	15.9	52.8	24.6	45.7	51.9	41.1
Virginia	49.1	40.3	26.2	8.8	60.7	38.1	47.4	69.0	47.4
Washington	33.2	35.8	24.9	12.5	43.7	32.8	41.6	47.3	46.3
West Virginia	40.7	31.3	10.1	9.6	56.8	39.4	41.7	55.6	44.4
Wisconsin	40.1	33.8	25.8	12.4	51.1	28.9	42.2	52.7	37.3
Wyoming	33.6	21.4	14.7	3.4	33.6	31.9	53.9	65.2	46.2
State Median	40.7	36.1	24.6	11.8	56.1	34.6	44.9	58.3	36.3
State Range	21.8-61.3	14.0-56.6	7.7-68.1	3.4-25.6	33.6-70.5	23.9-71.4	28.1-60.6	38.9-83.9	14.7-54.4

TABLE 20. Percentage of Secondary Schools in Which the Lead Health Education Teacher Received Professional Development* During the 2 Years Before the Survey on Teaching Methods, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Teaching students with physical, medical, or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Teaching students of different sexual orientations or gender identities	Using interactive teaching methods†	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
LOCAL SURVEYS									
Albuquerque	37.5	49.9	53.1	18.7	65.5	34.3	25.8	34.3	31.2
Baltimore	61.1	39.6	19.5	26.0	56.1	49.7	56.9	71.8	30.2
Broward County	44.6	67.3	62.4	25.3	61.5	37.1	57.6	68.0	41.5
Charlotte-Mecklenberg County	44.9	50.8	36.6	18.2	67.0	35.4	41.0	52.5	41.8
Chicago	68.8	53.6	40.7	21.7	66.3	53.0	57.1	70.5	43.0
Detroit	31.7	39.5	14.4	23.7	49.7	41.9	47.0	57.7	41.5
District of Columbia	24.1	27.3	12.0	59.2	74.1	40.7	50.0	60.7	62.6
Fresno	35.8	57.2	64.8	40.3	57.2	38.2	45.9	75.4	23.4
Houston	48.7	60.1	63.7	27.2	78.6	49.9	63.2	81.5	53.6
Los Angeles	50.4	73.6	81.5	44.3	77.9	48.9	56.9	70.9	48.2
Memphis	43.7	51.3	18.2	19.5	66.3	54.0	53.2	71.0	60.0
Miami-Dade County	32.9	48.1	47.7	30.9	62.3	49.5	45.0	62.9	37.7
Milwaukee	53.3	54.9	26.6	20.3	68.2	60.2	66.9	70.3	33.2
New York City	33.0	40.9	39.4	25.8	57.3	37.8	44.2	58.6	40.1
Newark	25.2	27.1	14.6	32.3	62.8	39.1	38.3	61.3	41.3
Orange County	44.3	55.8	53.5	34.9	69.7	34.7	44.2	62.6	32.4
Philadelphia	32.8	40.6	16.7	20.5	57.7	39.0	51.4	61.9	43.8
San Diego	45.5	63.9	76.8	32.4	52.6	40.8	32.0	55.2	17.0
San Francisco	60.1	78.7	71.0	55.2	84.2	57.4	68.3	70.4	55.2
Local Median	44.3	51.3	40.7	26.0	65.5	40.8	50.0	62.9	41.5
Local Range	24.1-68.8	27.1-78.7	12.0-81.5	18.2-59.2	49.7-84.2	34.3-60.2	25.8-68.3	34.3-81.5	17.0-62.6
TERRITORIAL SURVEYS									
American Samoa	38.5	50.0	96.2	88.5	88.5	88.5	92.3	92.3	96.2
Guam	69.2	69.2	61.5	30.8	61.5	38.5	41.7	30.8	23.1
Northern Mariana Islands	62.5	75.0	62.5	37.5	87.5	75.0	75.0	75.0	62.5
Palau	36.4	81.8	81.8	45.5	90.9	90.9	90.9	90.9	72.7
Territorial Median	50.5	72.1	72.2	41.5	88.0	81.8	83.0	83.0	67.6
Territorial Range	36.4-69.2	50.0-81.8	61.5-96.2	30.8-88.5	61.5-90.9	38.5-90.9	41.7-92.3	30.8-92.3	23.1-96.2
TRIBAL SURVEYS									
Cherokee Nation	59.6	51.7	29.3	15.3	51.3	51.9	51.6	70.3	41.4
Nez Perce	0.0	42.9	0.0	14.3	50.0	42.9	57.1	28.6	71.4

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Such as role plays or cooperative group activities.

TABLE 21. Percentage of Secondary Schools in Which the Lead Health Education Teacher Wanted to Receive Professional Development* on Teaching Methods, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010

Site	Teaching students with physical, medical, or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Teaching students of different sexual orientations or gender identities	Using interactive teaching methods ^f	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
STATE SURVEYS									
Alabama	74.1	70.0	58.6	52.5	68.6	77.3	77.5	70.6	73.1
Alaska	51.6	51.8	44.1	42.8	58.1	62.4	71.8	58.7	58.3
Arizona	58.2	56.5	52.4	52.6	58.2	66.2	68.4	63.4	62.6
Arkansas	65.7	62.4	53.2	47.0	63.1	72.3	74.0	69.6	73.0
California	58.8	50.5	50.2	53.4	57.3	62.5	71.0	54.7	64.8
Connecticut	73.1	62.2	52.8	61.6	78.5	71.7	81.2	64.2	74.9
Delaware	75.0	75.0	66.7	66.7	77.4	80.3	83.9	60.0	82.0
Florida	68.1	61.5	52.6	56.2	66.8	68.6	73.3	65.5	68.3
Georgia	62.6	59.6	54.8	47.3	64.1	71.6	75.9	63.6	70.6
Hawaii	76.3	70.1	77.0	71.1	75.0	74.2	81.4	64.7	79.0
Idaho	61.6	55.1	48.6	42.2	67.8	66.7	74.2	63.6	66.3
Indiana	52.9	37.7	35.0	40.3	58.4	58.1	62.9	58.5	59.5
Iowa	51.0	45.8	38.9	47.1	54.1	63.0	66.6	54.7	69.5
Kansas	59.6	49.7	41.5	44.2	58.8	65.4	67.1	64.6	64.4
Kentucky	64.0	53.7	48.1	40.5	64.1	67.6	67.3	61.8	69.5
Louisiana	58.9	56.9	48.8	50.4	61.6	69.3	72.0	67.5	69.0
Maine	58.6	34.7	29.5	55.1	63.7	66.3	76.1	61.3	74.1
Maryland	73.9	67.0	64.6	69.2	67.2	71.8	75.8	63.1	76.1
Massachusetts	69.8	65.0	59.5	74.1	76.8	76.5	81.8	71.4	79.8
Michigan	62.1	51.6	40.6	55.0	69.6	71.0	73.3	63.8	70.3
Minnesota	56.8	56.4	46.9	60.2	67.2	67.3	71.7	61.2	74.2
Mississippi	73.4	69.8	64.3	53.4	73.7	77.4	77.2	72.7	77.2
Missouri	64.8	54.1	42.4	44.4	65.0	74.0	77.4	67.6	69.2
Montana	57.1	44.4	31.6	43.9	65.4	65.6	76.4	70.2	72.3
Nebraska	51.0	43.0	34.9	37.3	49.1	61.1	63.8	58.7	62.2
Nevada	58.6	64.4	65.5	57.7	72.6	77.7	80.6	58.2	72.7
New Hampshire	60.8	45.8	41.7	64.6	69.7	72.4	77.0	62.2	73.4
New Jersey	82.1	67.7	64.5	75.0	82.6	81.1	87.8	79.6	79.9
New York	75.2	68.5	61.4	74.1	77.8	80.0	84.0	72.0	80.0
North Carolina	72.5	66.1	61.0	56.8	70.3	75.5	75.8	64.8	70.2
North Dakota	51.0	39.9	25.9	37.3	52.7	54.2	67.9	54.2	60.6
Ohio	60.0	54.0	42.7	52.2	70.0	72.4	72.9	65.4	69.5
Oklahoma	62.6	55.3	41.7	41.4	59.7	66.6	72.0	70.8	64.1
Oregon	53.8	51.4	42.1	52.9	62.9	65.6	71.0	53.7	64.5
Pennsylvania	71.3	60.0	50.9	61.1	69.2	72.5	76.4	67.7	76.7
Rhode Island	71.9	61.0	55.7	59.5	71.4	74.5	79.9	68.9	82.3
South Carolina	65.2	61.1	56.9	53.5	63.4	73.8	72.8	63.1	70.7
South Dakota	53.1	46.6	30.0	34.5	56.9	54.7	63.5	60.3	60.7
Tennessee	70.6	62.5	58.1	51.8	67.3	76.1	77.5	68.7	69.4
Texas	68.7	63.8	63.7	55.6	65.7	70.0	76.9	69.0	70.4
Utah	66.1	66.5	61.4	55.6	71.6	71.6	83.7	71.0	74.4
Vermont	48.0	39.0	34.3	58.6	62.7	70.3	73.3	55.6	70.3
Virginia	72.5	58.5	58.3	46.6	61.3	68.0	75.8	62.6	70.5
Washington	58.6	57.5	48.5	54.1	57.8	63.9	71.2	56.5	62.5
West Virginia	68.7	50.0	46.4	48.1	67.1	78.2	70.7	66.8	70.5
Wisconsin	57.3	54.3	47.3	60.1	64.9	69.4	74.8	59.9	71.8
Wyoming	65.3	57.5	48.7	63.5	63.7	69.4	71.4	61.8	63.4
State Median	62.6	56.9	48.8	53.4	65.4	70.3	74.2	63.6	70.4
State Range	48.0-82.1	34.7-75.0	25.9-77.0	34.5-75.0	49.1-82.6	54.2-81.1	62.9-87.8	53.7-79.6	58.3-82.3

TABLE 21. Percentage of Secondary Schools in Which the Lead Health Education Teacher Wanted to Receive Professional Development* on Teaching Methods, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2010 (continued)

Site	Teaching students with physical, medical, or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Teaching students of different sexual orientations or gender identities	Using interactive teaching methods†	Encouraging family or community involvement	Teaching skills for behavior change	Classroom management techniques	Assessing or evaluating students in health education
LOCAL SURVEYS									
Albuquerque	68.8	71.9	65.6	71.8	84.3	78.1	81.2	62.5	78.1
Baltimore	67.1	68.4	65.2	63.1	64.8	72.4	74.4	65.0	77.3
Broward County	67.1	68.3	57.9	65.3	67.6	66.4	67.7	59.3	61.9
Charlotte-Mecklenberg County	71.9	75.0	73.3	56.8	71.8	70.6	64.1	56.0	68.0
Chicago	76.8	66.2	67.5	71.9	74.5	79.2	87.6	81.1	73.8
Detroit	84.4	69.1	62.2	72.3	73.4	80.0	85.7	78.0	75.8
District of Columbia	90.8	84.7	78.7	85.5	79.4	85.4	88.3	85.5	82.3
Fresno	37.8	47.1	37.2	37.2	54.8	42.9	60.3	35.6	34.2
Houston	86.5	85.1	84.0	84.0	87.5	91.5	92.5	88.9	88.8
Los Angeles	84.1	75.1	76.8	81.6	75.9	88.8	89.6	70.3	79.9
Memphis	84.1	81.2	78.8	78.8	74.2	84.1	86.2	84.1	75.3
Miami-Dade County	72.3	63.9	60.2	65.9	70.4	71.4	71.1	67.4	61.5
Milwaukee	60.3	57.9	49.2	67.4	66.2	62.2	70.0	61.5	75.7
New York City	73.9	70.2	71.1	74.4	72.9	76.7	80.6	76.6	76.8
Newark	88.6	84.1	83.8	77.1	86.0	86.0	88.4	86.7	76.8
Orange County	71.3	61.8	64.2	68.8	63.4	80.9	85.5	78.4	78.2
Philadelphia	83.8	78.8	71.9	78.0	75.8	78.5	86.0	81.0	78.9
San Diego	46.7	40.2	40.4	50.1	45.2	53.6	68.4	44.8	50.2
San Francisco	79.0	81.8	76.3	84.2	81.6	86.5	89.3	86.7	81.4
Local Median	73.9	70.2	67.5	71.9	73.4	78.5	85.5	76.6	76.8
Local Range	37.8-90.8	40.2-85.1	37.2-84.0	37.2-85.5	45.2-87.5	42.9-91.5	60.3-92.5	35.6-88.9	34.2-88.8
TERRITORIAL SURVEYS									
American Samoa	96.2	96.2	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Guam	69.2	69.2	69.2	76.9	84.6	84.6	92.3	69.2	76.9
Northern Mariana Islands	71.4	71.4	85.7	85.7	85.7	85.7	71.4	85.7	85.7
Palau	90.9	63.6	72.7	45.5	72.7	81.8	63.6	90.9	90.9
Territorial Median	81.2	70.3	79.2	81.3	85.2	85.2	81.9	88.3	88.3
Territorial Range	69.2-96.2	63.6-96.2	69.2-100.0	45.5-100.0	72.7-100.0	81.8-100.0	63.6-100.0	69.2-100.0	76.9-100.0
TRIBAL SURVEYS									
Cherokee Nation	63.0	62.7	47.4	43.0	60.8	69.8	74.2	70.4	70.4
Nez Perce	57.1	71.4	28.6	42.9	42.9	42.9	85.7	57.1	28.6

*Such as workshops, conferences, continuing education, or any other kind of in-service.

†Such as role plays or cooperative group activities.

TABLE 22. Percentage of Secondary Schools That Required Physical Education in Any of Grades 6–12, the Percentage of Schools in Which at Least 1 Physical Education Teacher or Specialist Received Professional Development on Physical Education During the 2 Years Before the Survey, the Percentage of Schools That Offered Opportunities for All Students to Participate in Intramural Activities or Physical Activity Clubs,* and the Percentage of Schools in Which Children or Adolescents Use Any of the School's Indoor Physical Activity or Athletic Facilities for Community-Sponsored Physical Activity Classes or Lessons,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Required physical education	Physical education teacher or specialist received professional development on physical education	Offered intramural activities or physical activity clubs	Children or adolescents use school's indoor physical activity or athletic facilities for community-sponsored physical activity classes or lessons
STATE SURVEYS				
Alabama	95.0	84.9	48.8	66.6
Alaska	88.7	37.5	82.4	86.8
Arizona	78.0	66.7	72.1	66.3
Arkansas	98.9	93.6	50.0	70.4
California	98.5	81.2	76.3	82.5
Colorado	87.1	87.5	72.3	86.8
Connecticut	98.5	95.6	74.2	81.3
Delaware	98.4	93.9	73.6	81.0
Florida	94.1	92.7	77.2	66.9
Georgia	88.4	91.0	58.3	65.1
Hawaii	95.8	86.3	82.5	77.2
Idaho	90.8	74.9	59.7	78.0
Indiana	98.3	75.3	56.3	76.7
Iowa	97.1	80.8	50.9	86.3
Kansas	97.0	88.5	43.2	81.8
Kentucky	89.3	77.7	60.5	74.5
Louisiana	98.3	80.6	56.0	63.6
Maine	99.7	93.1	76.8	86.7
Maryland	94.0	94.0	87.1	90.1
Massachusetts	97.5	92.1	85.7	86.1
Michigan	89.6	83.6	60.5	80.4
Minnesota	94.4	87.3	65.2	88.6
Mississippi	94.0	82.6	57.2	54.8
Missouri	97.1	84.7	62.8	79.7
Montana	97.1	85.7	61.1	83.6
Nebraska	100.0	77.0	45.9	90.4
Nevada	95.0	88.7	85.3	70.6
New Hampshire	97.2	98.3	80.0	86.2
New Jersey	100.0	96.0	78.7	90.1
New Mexico	98.0	66.2	60.2	72.4
New York	100.0	96.7	86.1	87.9
North Carolina	97.0	92.0	60.6	71.1
North Dakota	97.0	77.5	45.9	72.6
Ohio	92.0	78.4	51.5	62.6
Oklahoma	61.1	74.5	52.4	74.2
Oregon	96.2	71.9	63.8	84.5
Pennsylvania	97.0	91.2	75.9	82.8
Rhode Island	99.0	89.3	73.1	78.9
South Carolina	93.4	94.3	58.2	72.2
South Dakota	95.0	76.8	40.6	84.1
Tennessee	93.0	90.7	60.7	68.0
Texas	97.7	93.5	53.1	79.4
Utah	99.5	92.1	72.2	92.8
Vermont	100.0	95.6	82.3	96.8
Virginia	96.5	95.6	67.9	88.9
Washington	96.7	82.7	63.0	87.0
West Virginia	98.0	88.4	61.7	88.6
Wisconsin	96.9	92.7	70.5	95.3
Wyoming	98.4	83.7	58.0	84.7
State Median	97.0	87.5	62.8	81.3
State Range	61.1-100.0	37.5-98.3	40.6-87.1	54.8-96.8

TABLE 22. Percentage of Secondary Schools That Required Physical Education in Any of Grades 6–12, the Percentage of Schools in Which at Least 1 Physical Education Teacher or Specialist Received Professional Development on Physical Education During the 2 Years Before the Survey, the Percentage of Schools That Offered Opportunities for All Students to Participate in Intramural Activities or Physical Activity Clubs,* and the Percentage of Schools in Which Children or Adolescents Use Any of the School’s Indoor Physical Activity or Athletic Facilities for Community-Sponsored Physical Activity Classes or Lessons,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Required physical education	Physical education teacher or specialist received professional development on physical education	Offered intramural activities or physical activity clubs	Children or adolescents use school’s indoor physical activity or athletic facilities for community-sponsored physical activity classes or lessons
LOCAL SURVEYS				
Albuquerque	100.0	94.8	91.4	89.0
Baltimore	83.5	86.8	79.3	70.8
Broward County	94.0	93.6	81.3	69.3
Charlotte-Mecklenberg County	96.3	94.2	47.5	69.5
Chicago	95.0	90.3	92.2	69.8
Detroit	79.7	87.3	71.8	68.2
District of Columbia	89.0	97.1	77.2	76.5
Fresno	95.6	95.6	100.0	91.8
Houston	97.5	98.7	64.1	48.2
Los Angeles	98.3	97.6	89.4	89.3
Memphis	85.3	98.1	77.1	54.4
Miami-Dade County	99.0	99.1	78.8	67.0
Milwaukee	89.9	89.9	93.0	81.0
New York City	100.0	94.9	89.0	82.4
Newark	100.0	100.0	76.7	70.6
Orange County	95.3	100.0	86.8	82.3
Philadelphia	95.9	96.5	84.8	60.4
San Diego	100.0	100.0	78.8	68.2
San Francisco	97.1	93.9	91.3	75.8
Local Median	95.9	95.6	81.3	70.6
Local Range	79.7-100.0	86.8-100.0	47.5-100.0	48.2-91.8
TERRITORIAL SURVEYS				
American Samoa	96.2	100.0	92.3	69.2
Guam	100.0	69.2	100.0	100.0
Marshall Islands	98.5	65.7	74.5	73.2
Northern Mariana Islands	85.7	75.0	100.0	87.5
Palau	100.0	81.8	90.9	9.1
Territorial Median	98.5	75.0	92.3	73.2
Territorial Range	85.7-100.0	65.7-100.0	74.5-100.0	9.1-100.0
TRIBAL SURVEYS				
Cherokee Nation	59.5	81.2	51.8	80.5
Nez Perce	100.0	85.7	28.6	57.1

*Any physical activity programs that are voluntary for students, in which students are given an equal opportunity to participate regardless of physical ability.
 †Outside of school hours or when school is not in session.

TABLE 23. Percentage of Secondary Schools That Taught a Required Physical Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
STATE SURVEYS							
Alabama	91.1	92.1	91.6	86.1	39.1	37.3	37.3
Alaska	78.1	79.5	78.9	79.0	71.6	64.7	64.6
Arizona	81.6	79.1	75.6	52.4	35.3	33.0	32.2
Arkansas	95.9	93.8	90.5	92.0	68.4	66.2	65.4
California	98.5	98.8	98.8	93.9	93.9	32.2	29.2
Colorado	77.0	73.7	70.9	74.3	64.1	52.2	49.4
Connecticut	96.6	97.1	97.1	91.5	92.1	74.1	65.1
Delaware	97.0	97.3	97.3	85.7	86.2	39.1	30.4
Florida	92.9	91.2	91.0	77.8	62.8	53.7	52.9
Georgia	80.8	80.9	81.0	77.0	39.4	34.5	35.5
Hawaii	82.9	83.4	70.2	83.9	70.3	22.6	19.4
Idaho	81.0	87.9	82.8	69.8	63.2	52.5	48.6
Indiana	93.0	97.1	96.5	94.0	34.5	12.4	13.0
Iowa	98.2	98.5	98.4	94.3	93.3	91.9	91.2
Kansas	91.2	88.7	85.4	92.4	18.0	9.4	9.4
Kentucky	79.9	80.8	79.1	72.9	27.7	14.2	14.0
Louisiana	98.4	98.1	97.4	92.1	94.5	63.4	59.4
Maine	99.5	99.0	99.0	92.9	82.1	26.2	24.4
Maryland	91.4	90.0	89.7	80.6	49.0	34.2	35.3
Massachusetts	97.6	98.1	97.6	92.0	91.8	73.4	68.9
Michigan	73.7	71.0	67.1	78.3	45.7	37.8	37.1
Minnesota	94.6	95.1	93.6	88.4	73.3	21.3	18.1
Mississippi	87.9	90.0	90.0	77.7	77.6	75.0	74.8
Missouri	95.9	97.1	97.1	89.6	60.7	48.1	47.6
Montana	92.0	95.6	95.6	94.1	97.1	17.3	17.3
Nebraska	99.1	99.3	99.3	89.0	48.5	21.3	21.2
Nevada	88.2	56.4	85.3	86.6	92.7	34.4	32.1
New Hampshire	94.8	96.4	96.4	91.0	77.7	44.7	38.5
New Jersey	98.9	99.5	99.5	99.0	100.0	100.0	100.0
New Mexico	72.3	82.9	44.5	87.6	57.4	52.9	50.6
New York	99.3	99.2	99.6	98.7	99.4	98.9	99.4
North Carolina	95.4	94.7	94.7	85.7	14.7	8.8	8.8
North Dakota	92.7	95.7	95.8	92.5	73.4	42.0	38.6
Ohio	85.6	85.1	86.9	77.6	72.0	25.9	26.7
Oklahoma	66.0	47.5	46.9	15.4	13.0	12.4	12.4
Oregon	91.6	92.6	92.6	84.5	53.5	31.6	26.8
Pennsylvania	94.7	96.2	96.1	90.7	94.2	89.9	85.4
Rhode Island	97.3	97.8	97.8	94.0	95.8	95.8	95.8
South Carolina	90.2	91.1	91.1	81.8	46.4	42.9	42.5
South Dakota	84.0	86.4	85.1	66.1	21.1	15.1	20.9
Tennessee	88.8	90.1	90.2	74.3	55.7	32.8	31.8
Texas	95.6	94.9	81.3	92.3	88.6	69.8	68.4
Utah	96.6	97.2	92.1	94.0	93.5	81.1	61.4
Vermont	100.0	100.0	100.0	89.6	86.1	54.7	50.5
Virginia	93.6	94.7	88.2	91.5	96.8	7.5	6.1
Washington	95.5	96.8	96.0	82.1	77.2	50.7	44.7
West Virginia	98.5	98.7	98.7	85.3	56.2	33.0	34.2
Wisconsin	97.9	98.1	98.1	91.9	90.0	81.2	46.2
Wyoming	100.0	100.0	98.7	93.8	54.1	29.6	29.1
State Median	93.6	94.9	92.6	88.4	71.6	39.1	37.3
State Range	66.0-100.0	47.5-100.0	44.5-100.0	15.4-99.0	13.0-100.0	7.5-100.0	6.1-100.0

TABLE 23. Percentage of Secondary Schools That Taught a Required Physical Education Course in Each Grade,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Grade 6	Grade 7	Grade 8	Grade 9	Grade 10	Grade 11	Grade 12
LOCAL SURVEYS							
Albuquerque	87.5	48.2	20.3	100.0	15.4	16.7	16.7
Baltimore	70.0	69.4	72.8	61.4	73.3	58.3	62.5
Broward County	85.3	85.8	85.8	86.0	79.2	75.0	75.0
Charlotte-Mecklenberg County	96.8	96.8	96.8	84.6	29.2	25.0	26.1
Chicago	92.5	93.9	93.9	72.3	88.1	31.1	21.5
Detroit	63.9	63.9	61.9	48.5	95.2	85.7	85.7
District of Columbia	78.9	80.0	78.9	76.8	100.0	61.5	61.5
Fresno	100.0	100.0	100.0	87.3	85.7	0.0	0.0
Houston	98.0	98.0	98.0	91.1	93.7	87.5	87.5
Los Angeles	98.6	98.6	98.6	94.2	94.0	35.7	31.7
Memphis	69.4	71.4	71.4	67.7	54.5	50.0	45.5
Miami-Dade County	98.5	90.9	91.7	78.7	72.2	37.8	30.6
Milwaukee	87.7	87.7	87.7	69.0	100.0	87.5	87.5
New York City	98.7	98.6	99.3	96.1	98.3	96.6	98.3
Newark	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Orange County	96.4	96.4	96.4	86.1	50.0	35.7	33.3
Philadelphia	93.7	93.9	93.8	78.6	81.2	71.9	80.0
San Diego	100.0	100.0	100.0	100.0	97.0	36.4	36.4
San Francisco	100.0	100.0	100.0	84.0	87.5	18.8	18.8
Local Median	96.4	93.9	93.9	84.6	87.5	50.0	45.5
Local Range	63.9-100.0	48.2-100.0	20.3-100.0	48.5-100.0	15.4-100.0	0.0-100.0	0.0-100.0
TERRITORIAL SURVEYS							
American Samoa	95.0	95.0	95.0	85.7	100.0	80.0	80.0
Guam	100.0	100.0	37.5	80.0	50.0	25.0	25.0
Marshall Islands	95.1	93.2	93.3	55.3	100.0	100.0	75.0
Northern Mariana Islands	100.0	100.0	100.0	80.0	75.0	50.0	50.0
Palau	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	100.0	100.0	95.0	80.0	100.0	80.0	75.0
Territorial Range	95.0-100.0	93.2-100.0	37.5-100.0	55.3-100.0	50.0-100.0	25.0-100.0	25.0-100.0
TRIBAL SURVEYS							
Cherokee Nation	67.5	52.0	52.0	12.1	4.6	4.7	4.7
Nez Perce	50.0	75.0	75.0	60.0	75.0	75.0	75.0

*Among schools with students in that grade.

TABLE 24. Among Secondary Schools That Required Physical Education for Students in Any of Grades 6–12, the Percentage That Allowed Students to be Exempted* from Taking Required Physical Education for Specific Reasons, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Religious reasons	Long-term physical or medical disability	Cognitive disability	Enrollment in other courses	Participation in school sports	Participation in other school activities ¹	Participation in community sports activities	High physical fitness competency test score	Participation in vocational training	Participation in community service activities	Could not be exempted for any of these 7 reasons ¹
STATE SURVEYS											
Alabama	22.6	50.3	22.7	13.3	12.9	25.2	4.5	5.0	4.6	0.9	65.1
Alaska	48.7	81.9	51.1	38.1	36.9	19.2	20.9	7.6	13.2	11.2	41.4
Arizona	54.6	86.1	47.4	32.1	15.1	27.3	9.6	4.6	6.7	5.9	52.1
Arkansas	35.2	70.8	31.7	17.3	49.1	24.2	4.9	1.3	6.0	1.7	39.9
California	29.6	84.6	29.7	8.8	18.5	19.1	9.7	5.0	2.3	0.8	62.1
Colorado	58.2	88.5	42.5	37.1	44.5	33.2	17.2	3.6	6.2	5.2	30.7
Connecticut	38.3	88.1	19.5	5.8	1.6	2.5	1.6	1.5	1.1	1.3	88.9
Delaware	44.2	89.5	32.7	14.0	3.1	9.3	3.0	3.1	3.1	3.1	77.8
Florida	60.9	85.7	47.7	57.8	42.4	62.9	36.0	10.3	8.7	7.7	14.9
Georgia	48.6	75.6	37.4	42.6	6.0	50.9	1.8	2.7	7.0	2.6	36.8
Hawaii	46.2	86.6	40.2	27.1	6.8	18.3	2.6	2.6	4.1	1.5	61.0
Idaho	45.9	85.8	48.6	30.9	13.6	13.9	7.0	2.8	5.1	3.3	58.3
Indiana	43.0	80.2	35.3	20.4	10.2	15.0	0.9	1.7	0.9	0.3	65.4
Iowa	65.2	84.0	28.9	44.7	25.7	9.5	8.7	0.5	5.0	0.5	46.3
Kansas	55.1	79.6	41.8	16.9	4.8	9.7	1.7	1.5	1.4	2.3	74.7
Kentucky	38.3	72.5	25.3	20.9	1.4	26.1	1.4	1.4	2.5	1.4	64.1
Louisiana	41.7	80.7	42.5	23.7	9.7	29.7	4.2	2.7	8.2	4.6	54.0
Maine	40.6	74.8	23.6	14.3	4.4	3.1	2.4	1.2	1.2	0.8	83.2
Maryland	42.7	67.7	21.6	21.0	0.9	7.8	0.5	0.4	2.3	1.8	73.8
Massachusetts	47.7	89.7	24.3	17.0	7.0	9.1	4.1	1.3	4.5	2.5	72.2
Michigan	54.3	88.8	40.6	41.6	16.0	34.5	4.0	5.5	7.5	3.5	40.2
Minnesota	32.6	82.7	35.4	17.1	1.9	4.5	1.6	0.9	3.2	2.1	80.9
Mississippi	47.1	83.4	54.8	31.7	66.6	70.6	6.6	5.6	10.7	5.0	17.1
Missouri	41.9	81.0	39.3	23.2	4.1	9.4	2.0	2.6	5.0	3.4	71.7
Montana	30.7	86.3	34.3	17.8	2.1	2.1	1.5	0.5	1.0	0.6	79.2
Nebraska	37.0	74.4	37.4	18.2	8.6	7.8	1.6	1.6	5.1	4.0	72.5
Nevada	52.4	91.3	42.3	32.6	56.9	47.1	20.5	2.7	4.1	2.7	21.5
New Hampshire	39.5	88.4	29.8	22.6	11.9	10.1	6.5	4.0	2.9	2.3	67.9
New Jersey	37.6	86.3	23.8	2.9	3.8	3.2	3.3	0.7	1.9	1.0	88.2
New Mexico	46.6	89.5	41.2	20.7	25.6	20.2	4.9	3.9	7.0	4.8	53.4
New York	23.7	67.0	17.2	3.9	11.3	2.8	3.5	1.7	2.2	1.8	83.5
North Carolina	32.6	67.5	23.5	17.9	2.8	11.1	1.5	0.8	2.7	1.1	77.4
North Dakota	20.0	68.3	24.4	18.7	6.2	5.2	0.6	0.6	2.9	1.3	77.5
Ohio	35.3	69.1	17.2	13.5	14.2	16.5	2.9	1.7	3.5	2.8	66.3
Oklahoma	59.8	78.4	53.5	30.5	58.5	32.8	12.7	13.8	23.7	16.4	37.4
Oregon	59.5	87.2	38.9	30.2	11.0	14.9	10.1	4.3	7.1	5.9	61.3
Pennsylvania	44.1	82.5	31.5	11.2	4.6	5.4	1.8	0.6	7.1	1.4	78.4
Rhode Island	25.1	79.0	13.9	7.3	2.2	1.2	2.2	2.3	3.1	2.1	89.3
South Carolina	54.2	84.8	32.2	27.6	4.6	54.8	3.3	0.9	2.2	1.6	39.3
South Dakota	19.0	61.1	27.2	11.8	4.4	4.8	0.0	0.0	0.5	0.0	84.8
Tennessee	44.2	79.8	40.3	14.7	12.9	37.8	1.7	3.9	4.5	2.0	54.8
Texas	39.2	77.3	42.1	21.1	72.9	55.0	20.6	4.8	18.9	3.1	17.4
Utah	61.3	93.7	56.8	27.7	32.5	12.2	6.4	12.2	4.7	2.3	45.3
Vermont	31.5	78.7	21.5	11.5	17.7	5.3	7.9	0.0	1.7	1.7	73.4
Virginia	42.0	74.2	31.4	13.2	3.3	8.5	3.3	0.6	2.7	0.6	80.5
Washington	64.4	87.9	43.6	45.0	40.1	30.8	27.5	2.8	11.2	4.7	29.5
West Virginia	28.5	70.7	18.1	10.2	1.2	3.2	2.3	1.2	2.4	1.2	86.8
Wisconsin	41.0	80.2	21.3	10.8	1.8	3.8	2.0	0.5	2.3	1.5	85.1
Wyoming	24.0	85.2	34.4	19.8	5.0	4.9	0.9	0.0	3.3	0.0	76.9
State Median	42.0	81.9	34.3	19.8	9.7	12.2	3.3	1.7	4.1	2.1	65.4
State Range	19.0-65.2	50.3-93.7	13.9-56.8	2.9-57.8	0.9-72.9	1.2-70.6	0.0-36.0	0.0-13.8	0.5-23.7	0.0-16.4	14.9-89.3

TABLE 24. Among Secondary Schools That Required Physical Education for Students in Any of Grades 6–12, the Percentage That Allowed Students to be Exempted* from Taking Required Physical Education for Specific Reasons, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Religious reasons	Long-term physical or medical disability	Cognitive disability	Enrollment in other courses	Participation in school sports	Participation in other school activities [†]	Participation in community sports activities	High physical fitness competency test score	Participation in vocational training	Participation in community service activities	Could not be exempted for any of these 7 reasons [‡]
LOCAL SURVEYS											
Albuquerque	58.4	89.3	43.6	18.8	7.8	13.1	5.2	5.2	5.2	5.2	75.8
Baltimore	51.2	76.6	32.6	28.3	4.6	10.5	4.7	3.1	4.6	7.5	69.8
Broward County	59.4	82.6	56.5	49.2	52.7	67.8	31.1	9.1	5.4	3.6	15.0
Charlotte-Mecklenberg County	40.6	59.4	26.2	24.1	2.0	14.1	0.0	0.0	0.0	0.0	71.9
Chicago	59.7	78.5	19.7	2.2	2.2	12.2	2.2	2.2	2.2	2.2	86.3
Detroit	58.5	89.2	38.9	29.1	24.4	50.5	4.7	1.5	9.2	6.0	41.9
District of Columbia	46.5	61.2	12.8	12.9	6.4	12.8	9.6	9.6	6.4	6.4	71.1
Fresno	54.6	87.9	34.7	4.0	8.6	23.0	4.6	0.0	0.0	0.0	64.4
Houston	28.1	39.5	27.8	16.8	25.4	38.0	12.8	7.7	6.4	3.9	54.3
Los Angeles	26.6	73.0	19.5	6.8	27.8	25.1	4.3	6.7	0.8	2.5	60.4
Memphis	45.9	70.2	49.6	21.1	10.5	63.3	4.1	8.2	10.5	5.9	31.0
Miami-Dade County	67.2	86.6	50.8	67.9	22.9	36.2	15.9	5.9	7.8	3.0	21.0
Milwaukee	55.4	80.8	26.1	11.5	1.6	6.4	3.2	0.0	1.7	4.8	82.4
New York City	35.6	79.7	28.3	8.4	16.7	5.6	6.1	3.4	4.6	4.2	73.8
Newark	45.4	95.9	34.2	0.0	57.8	27.5	15.6	5.9	0.0	4.1	32.1
Orange County	66.7	82.5	50.8	63.1	48.9	61.1	41.0	9.7	7.2	12.0	17.0
Philadelphia	56.6	84.6	36.3	10.0	5.4	2.2	3.9	3.7	1.4	3.0	81.8
San Diego	7.5	100.0	61.2	1.5	44.8	49.3	1.5	3.0	26.9	1.5	47.8
San Francisco	40.4	74.5	27.9	6.0	8.9	8.9	6.2	6.0	0.0	3.0	75.9
Local Median	51.2	80.8	34.2	12.9	10.5	23.0	4.7	5.2	4.6	3.9	64.4
Local Range	7.5-67.2	39.5-100.0	12.8-61.2	0.0-67.9	1.6-57.8	2.2-67.8	0.0-41.0	0.0-9.7	0.0-26.9	0.0-12.0	15.0-86.3
TERRITORIAL SURVEYS											
American Samoa	0.0	12.0	8.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0
Guam	8.3	83.3	8.3	25.0	0.0	16.7	0.0	0.0	0.0	0.0	66.7
Marshall Islands	29.7	58.6	65.5	34.9	42.2	34.4	40.6	34.9	26.1	41.0	34.6
Northern Mariana Islands	0.0	66.7	33.3	16.7	0.0	33.3	0.0	0.0	0.0	0.0	66.7
Palau	0.0	100.0	100.0	10.0	0.0	0.0	0.0	10.0	0.0	0.0	80.0
Territorial Median	0.0	66.7	33.3	16.7	0.0	16.7	0.0	0.0	0.0	0.0	66.7
Territorial Range	0.0-29.7	12.0-100.0	8.0-100.0	0.0-34.9	0.0-42.2	0.0-34.4	0.0-40.6	0.0-34.9	0.0-26.1	0.0-41.0	34.6-100.0
TRIBAL SURVEYS											
Cherokee Nation	59.8	87.2	65.2	36.4	63.3	40.8	17.8	16.4	28.0	18.1	30.2
Nez Perce	0.0	42.9	14.3	14.3	0.0	0.0	0.0	0.0	0.0	0.0	85.7

*For one grading period or longer.

†Such as ROTC, band, or chorus.

‡These reasons included enrollment in other courses, participation in school sports, participation in other school activities, participation in community sports activities, high physical fitness competency test score, participation in vocational training, and participation in community service activities.

TABLE 25. Percentage of Secondary Schools That Provided Those Who Teach Physical Education with Materials for Teaching Physical Education, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Goals, objectives, and expected outcomes for physical education	Chart describing annual scope and sequence of instruction for physical education	Plans for how to assess student performance in physical education	Written physical education curriculum	All 4 types of materials
STATE SURVEYS					
Alabama	97.7	68.5	70.6	83.6	59.1
Alaska	67.8	51.7	52.2	60.6	40.9
Arizona	87.3	64.5	69.5	69.2	52.0
Arkansas	97.4	74.3	80.0	93.6	67.4
California	94.6	80.7	86.4	80.1	69.4
Colorado	93.4	72.4	82.8	80.4	63.1
Connecticut	96.6	87.0	93.2	92.8	81.4
Delaware	95.3	92.4	90.7	89.2	82.7
Florida	97.7	88.1	89.8	90.8	80.1
Georgia	95.6	88.5	88.8	94.2	80.6
Hawaii	96.8	86.4	87.7	81.7	75.0
Idaho	85.9	68.0	70.2	76.0	59.0
Indiana	91.5	71.7	74.4	90.8	62.7
Iowa	96.6	77.4	78.0	91.6	64.8
Kansas	94.6	72.4	75.8	89.0	62.9
Kentucky	97.0	85.7	86.8	89.7	76.8
Louisiana	96.5	75.7	84.3	92.0	69.9
Maine	92.7	78.9	85.4	91.5	71.6
Maryland	99.1	96.1	95.7	97.4	93.9
Massachusetts	93.7	83.0	84.4	88.1	74.6
Michigan	94.3	78.9	78.2	82.1	68.2
Minnesota	89.3	75.3	77.0	78.5	65.0
Mississippi	97.3	74.6	84.7	95.9	70.3
Missouri	96.1	80.6	88.9	92.3	75.4
Montana	95.5	81.5	76.7	90.7	67.2
Nebraska	90.1	66.3	74.8	83.3	57.7
Nevada	94.9	79.7	82.5	89.1	75.2
New Hampshire	96.8	79.8	84.6	91.2	71.7
New Jersey	99.0	89.4	92.2	98.3	85.3
New Mexico	92.6	69.6	80.8	85.2	65.8
New York	95.0	87.5	91.5	90.3	81.1
North Carolina	96.7	82.5	84.2	92.0	76.1
North Dakota	86.7	62.4	63.8	70.7	50.3
Ohio	91.6	65.6	70.2	82.3	56.3
Oklahoma	86.4	49.3	62.7	59.1	39.1
Oregon	89.0	69.7	68.8	66.8	54.5
Pennsylvania	96.8	86.6	88.8	91.3	81.0
Rhode Island	93.4	74.7	77.9	85.0	64.8
South Carolina	99.1	84.9	93.4	89.3	78.9
South Dakota	81.9	49.6	58.8	60.6	40.0
Tennessee	96.9	77.0	85.4	91.4	72.0
Texas	95.3	79.6	89.8	83.9	71.9
Utah	97.8	86.0	89.9	93.8	79.9
Vermont	93.6	76.8	82.3	85.5	72.2
Virginia	97.2	88.8	90.1	93.2	81.5
Washington	93.0	74.0	82.3	68.9	60.9
West Virginia	100.0	82.9	91.9	91.7	78.0
Wisconsin	97.7	85.7	87.5	93.4	78.4
Wyoming	96.9	83.2	93.0	79.9	72.6
State Median	95.3	78.9	84.3	89.2	71.6
State Range	67.8-100.0	49.3-96.1	52.2-95.7	59.1-98.3	39.1-93.9

TABLE 25. Percentage of Secondary Schools That Provided Those Who Teach Physical Education with Materials for Teaching Physical Education, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Goals, objectives, and expected outcomes for physical education	Chart describing annual scope and sequence of instruction for physical education	Plans for how to assess student performance in physical education	Written physical education curriculum	All 4 types of materials
LOCAL SURVEYS					
Albuquerque	97.4	84.1	86.4	97.4	81.2
Baltimore	96.4	86.4	86.3	87.6	83.8
Broward County	98.5	92.5	92.7	92.4	88.2
Charlotte-Mecklenberg County	100.0	92.4	98.0	94.2	90.4
Chicago	91.2	76.8	78.6	71.7	62.9
Detroit	92.0	85.0	89.7	89.6	81.6
District of Columbia	97.2	88.3	80.1	85.8	70.6
Fresno	100.0	90.9	100.0	100.0	90.9
Houston	97.6	77.4	94.0	91.5	72.5
Los Angeles	96.6	91.6	92.5	89.2	83.2
Memphis	100.0	96.0	98.0	96.7	90.8
Miami-Dade County	100.0	95.3	98.1	97.1	93.3
Milwaukee	98.5	91.4	87.3	88.7	81.6
New York City	95.6	87.4	94.5	83.3	77.8
Newark	100.0	94.3	96.0	100.0	90.3
Orange County	100.0	91.5	91.3	89.2	82.8
Philadelphia	95.7	89.2	92.1	90.3	84.7
San Diego	100.0	100.0	100.0	98.5	98.5
San Francisco	93.8	90.5	93.8	90.8	87.5
Local Median	97.6	90.9	92.7	90.8	83.8
Local Range	91.2-100.0	76.8-100.0	78.6-100.0	71.7-100.0	62.9-98.5
TERRITORIAL SURVEYS					
American Samoa	100.0	100.0	100.0	100.0	100.0
Guam	92.3	69.2	84.6	92.3	69.2
Marshall Islands	85.3	75.1	79.6	85.4	64.9
Northern Mariana Islands	100.0	87.5	100.0	100.0	87.5
Palau	100.0	100.0	100.0	100.0	100.0
Territorial Median	100.0	87.5	100.0	100.0	87.5
Territorial Range	85.3-100.0	69.2-100.0	79.6-100.0	85.4-100.0	64.9-100.0
TRIBAL SURVEYS					
Cherokee Nation	88.7	59.4	71.3	65.0	47.2
Nez Perce	85.7	57.1	57.1	85.7	57.1

TABLE 26. Percentage of Secondary Schools That Allowed Students to Purchase Snack Foods or Beverages from One or More Vending Machines or at the School Store, Canteen, or Snack Bar, the Percentage That Limited the Package or Serving Size of Any Individual Food or Beverage Item in These Venues, the Percentage That Allowed Students to Purchase Fruits and Non-Fried Vegetables From These Venues, and the Percentage That Always or Almost Always Offered Fruits or Non-Fried Vegetables at School Celebrations,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Allowed students to purchase snack foods or beverages	Limited the package or serving size of any individual food or beverage items	Allowed students to purchase fruits (not fruit juice)	Allowed students to purchase non-fried vegetables (not vegetable juice)	Always or almost always offered fruits or non-fried vegetables at school celebrations	Made fruits and non-fried vegetables available in both ways
STATE SURVEYS						
Alabama	63.5	57.4	9.1	6.6	32.0	4.3
Alaska	53.0	29.8	8.7	7.7	40.7	3.8
Arizona	51.6	33.4	23.3	15.7	42.2	10.1
Arkansas	51.9	38.3	16.0	8.6	21.2	4.2
California	66.4	51.5	41.7	33.0	35.5	14.9
Colorado	70.5	41.7	26.5	16.7	33.3	9.5
Connecticut	53.2	40.0	28.3	20.2	43.6	13.3
Delaware	58.2	40.6	27.0	17.5	35.4	11.0
Florida	73.0	47.5	39.0	32.4	34.4	16.7
Georgia	81.2	52.7	25.1	15.5	30.5	8.5
Hawaii	37.0	17.8	2.8	2.8	38.9	1.4
Idaho	80.7	44.8	28.0	13.6	32.9	9.2
Indiana	77.0	52.8	33.2	23.3	22.3	8.1
Iowa	76.4	51.7	31.6	14.9	22.4	7.8
Kansas	76.1	45.0	26.7	15.3	17.0	4.0
Kentucky	70.8	47.3	16.3	10.8	23.1	4.2
Louisiana	84.5	64.4	11.9	6.2	29.9	6.5
Maine	67.5	43.8	31.7	22.2	48.5	14.4
Maryland	63.5	41.2	31.3	27.0	40.6	12.9
Massachusetts	67.6	46.6	36.0	28.9	43.1	15.9
Michigan	78.0	48.9	46.3	36.3	27.6	16.8
Minnesota	75.9	49.9	39.9	22.4	28.4	11.4
Mississippi	69.5	54.8	21.2	11.1	32.4	7.4
Missouri	75.2	45.0	31.0	21.0	23.9	8.8
Montana	79.1	40.9	27.7	11.7	29.6	5.6
Nebraska	78.4	45.1	24.3	14.6	15.9	5.3
Nevada	90.2	83.5	39.1	30.8	37.2	22.5
New Hampshire	82.7	63.8	50.2	33.5	44.2	20.9
New Jersey	65.8	48.9	38.6	30.6	38.6	16.0
New Mexico	61.9	43.5	21.7	11.7	42.4	11.0
New York	78.4	48.2	38.6	30.7	41.2	16.9
North Carolina	65.0	41.6	26.8	22.5	26.6	8.3
North Dakota	64.0	32.2	17.8	10.9	28.1	4.4
Ohio	70.5	41.5	34.1	26.3	26.7	8.3
Oklahoma	75.3	42.8	25.3	14.7	22.3	5.2
Oregon	61.8	44.6	21.7	15.4	41.7	11.0
Pennsylvania	67.8	45.7	40.5	33.2	40.4	17.9
Rhode Island	66.7	43.7	30.4	25.0	46.2	14.7
South Carolina	80.3	51.7	32.4	24.9	36.8	14.7
South Dakota	76.3	36.4	17.9	5.5	12.2	5.3
Tennessee	57.2	45.3	21.8	15.8	28.4	9.9
Texas	63.9	54.0	35.5	31.0	25.1	9.8
Utah	92.8	60.3	38.9	32.4	31.4	14.8
Vermont	70.1	49.1	42.9	35.8	52.4	23.4
Virginia	72.5	45.8	30.9	25.2	37.2	13.2
Washington	80.2	50.1	36.3	25.1	29.1	10.5
West Virginia	61.9	54.0	7.8	2.3	33.4	4.0
Wisconsin	72.9	45.5	31.1	21.3	24.8	11.4
Wyoming	70.6	30.0	21.3	13.4	29.7	8.7
State Median	70.5	45.5	28.3	20.2	32.4	9.9
State Range	37.0-92.8	17.8-83.5	2.8-50.2	2.3-36.3	12.2-52.4	1.4-23.4

TABLE 26. Percentage of Secondary Schools That Allowed Students to Purchase Snack Foods or Beverages from One or More Vending Machines or at the School Store, Canteen, or Snack Bar, the Percentage That Limited the Package or Serving Size of Any Individual Food or Beverage Item in These Venues, the Percentage That Allowed Students to Purchase Fruits and Non-Fried Vegetables From These Venues, and the Percentage That Always or Almost Always Offered Fruits or Non-Fried Vegetables at School Celebrations,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Allowed students to purchase snack foods or beverages	Limited the package or serving size of any individual food or beverage items	Allowed students to purchase fruits (not fruit juice)	Allowed students to purchase non-fried vegetables (not vegetable juice)	Always or almost always offered fruits or non-fried vegetables at school celebrations	Made fruits and non-fried vegetables available in both ways
LOCAL SURVEYS						
Albuquerque	62.9	57.4	38.3	30.3	51.2	24.5
Baltimore	26.8	15.1	12.7	6.4	51.8	7.4
Broward County	89.4	46.0	41.3	25.4	45.3	20.4
Charlotte-Mecklenberg County	69.5	35.5	33.4	27.4	18.6	11.2
Chicago	25.8	15.1	8.0	5.1	37.3	3.9
Detroit	34.2	22.2	19.7	12.8	53.3	13.7
District of Columbia	42.9	26.4	19.9	14.3	30.6	11.1
Fresno	73.1	52.7	33.5	25.3	40.1	20.9
Houston	75.7	27.6	46.1	32.5	35.9	13.3
Los Angeles	88.0	66.1	45.8	30.1	44.1	24.9
Memphis	35.0	29.4	18.8	14.1	40.9	9.6
Miami-Dade County	81.0	52.0	44.4	32.7	42.6	20.7
Milwaukee	21.5	17.1	10.1	7.2	26.5	4.3
New York City	65.3	37.1	17.3	12.8	49.2	10.8
Newark	45.6	30.1	15.6	9.0	33.7	6.6
Orange County	78.7	60.8	46.7	40.1	41.5	22.3
Philadelphia	45.4	21.1	17.8	10.6	43.6	10.8
San Diego	62.7	56.7	38.8	32.8	29.9	16.4
San Francisco	50.5	47.6	29.2	20.5	91.1	23.4
Local Median	62.7	35.5	29.2	20.5	41.5	13.3
Local Range	21.5-89.4	15.1-66.1	8.0-46.7	5.1-40.1	18.6-91.1	3.9-24.9
TERRITORIAL SURVEYS						
American Samoa	0.0	0.0	0.0	0.0	76.9	0.0
Guam	76.9	38.5	23.1	15.4	46.2	15.4
Marshall Islands	40.4	11.0	34.1	24.8	24.2	11.7
Northern Mariana Islands	50.0	28.6	12.5	0.0	75.0	12.5
Palau	63.6	20.0	63.6	54.5	63.6	63.6
Territorial Median	50.0	20.0	23.1	15.4	63.6	12.5
Territorial Range	0.0-76.9	0.0-38.5	0.0-63.6	0.0-54.5	24.2-76.9	0.0-63.6
TRIBAL SURVEYS						
Cherokee Nation	76.9	44.4	28.6	20.8	17.6	5.0
Nez Perce	71.4	42.9	28.6	28.6	0.0	0.0

*When foods or beverages are offered.

TABLE 27. Percentage of Secondary Schools That Allowed Students to Purchase Less Nutritious Foods and Beverages From Vending Machines or at the School Store, Canteen, or Snack Bar and the Percentage That Did Not Sell Baked Goods,* Salty Snacks,* Candy, Soda Pop or Fruit Drinks,† or Sports Drinks in These Venues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	2% or whole milk (plain or flavored)	Foods or beverages containing caffeine	Ice cream or frozen yogurt*	Water ices or frozen slushes that do not contain juice	Cookies, crackers, cakes, pastries, or other baked goods	Salty snacks	Chocolate candy	Other kinds of candy	Soda pop or fruit drinks	Sports drinks	Did not sell any of these 6 items
STATE SURVEYS											
Alabama	12.6	21.1	2.7	7.8	6.1	10.5	3.3	9.0	21.1	38.8	55.8
Alaska	12.5	23.6	4.0	11.5	17.3	26.2	20.4	23.9	25.9	38.0	54.1
Arizona	24.3	19.3	15.7	9.5	24.9	25.8	17.3	20.6	22.8	35.3	56.2
Arkansas	20.7	30.1	7.7	8.5	22.5	22.6	15.3	17.0	30.4	39.6	52.6
California	36.0	5.1	11.4	12.7	18.7	20.1	4.6	10.1	11.0	51.9	44.1
Colorado	22.2	25.6	12.1	17.7	40.9	38.8	29.8	32.5	23.4	50.7	38.4
Connecticut	24.1	3.2	13.4	7.8	14.4	14.5	2.4	3.6	3.1	8.4	77.1
Delaware	26.9	17.3	4.7	7.9	17.3	15.7	10.9	15.7	14.1	34.4	57.6
Florida	43.4	29.2	30.7	25.3	42.4	38.6	21.4	30.0	37.4	58.5	35.3
Georgia	43.2	39.3	41.4	18.7	51.0	50.5	44.1	51.8	42.5	62.5	28.5
Hawaii	6.3	7.0	5.3	11.3	9.5	16.2	7.4	12.3	8.8	16.1	75.8
Idaho	38.7	43.4	14.2	16.0	44.4	47.1	46.3	47.8	49.7	56.3	25.8
Indiana	41.0	45.7	22.8	18.6	42.3	43.3	37.5	41.0	47.4	67.7	28.5
Iowa	35.6	35.8	14.5	12.1	27.9	24.4	17.5	22.2	37.9	65.5	30.9
Kansas	30.3	48.3	13.4	13.9	43.0	38.9	43.1	44.6	50.3	69.0	27.4
Kentucky	30.3	36.8	7.0	10.6	17.1	20.4	19.0	26.4	34.3	46.1	45.7
Louisiana	25.1	43.0	6.7	17.8	36.5	48.6	45.1	65.8	41.7	60.2	21.2
Maine	26.8	9.6	11.3	5.4	19.7	19.6	4.7	7.6	10.9	40.3	52.3
Maryland	41.0	27.0	31.9	21.6	46.8	49.7	29.7	34.5	33.3	47.8	38.8
Massachusetts	34.7	11.4	25.0	14.2	27.8	23.8	8.3	11.7	15.6	37.1	48.4
Michigan	50.2	34.1	27.7	24.7	52.6	48.1	35.4	40.7	37.5	62.5	26.7
Minnesota	37.5	45.4	20.6	19.4	47.0	44.8	39.1	45.0	41.5	66.1	25.7
Mississippi	27.1	19.6	14.9	9.5	11.6	13.5	4.3	13.2	16.6	40.6	51.1
Missouri	37.2	38.4	18.3	14.7	41.8	38.7	33.2	37.5	43.8	63.9	28.2
Montana	28.8	42.8	12.8	12.9	45.3	33.7	36.8	41.1	43.1	71.4	25.8
Nebraska	34.8	43.3	16.1	15.6	32.5	32.2	35.1	39.1	48.7	69.6	24.2
Nevada	43.1	6.5	13.8	15.8	19.8	15.3	8.3	27.7	11.5	66.5	26.6
New Hampshire	49.1	22.4	30.7	13.5	32.4	26.5	6.8	11.2	19.2	42.9	35.5
New Jersey	39.5	15.2	26.3	15.6	26.5	23.2	7.5	11.2	15.5	40.5	47.3
New Mexico	28.7	12.7	3.9	7.2	17.7	20.7	14.2	21.2	16.0	39.2	52.3
New York	39.8	18.8	29.3	16.2	35.5	31.6	10.5	15.5	24.4	45.8	38.2
North Carolina	30.8	29.0	17.2	13.9	38.5	39.5	27.3	30.5	33.2	51.3	41.2
North Dakota	18.8	29.9	9.3	14.9	21.7	20.0	21.4	24.5	33.4	53.5	41.8
Ohio	43.3	33.2	27.6	20.9	42.1	41.8	28.3	30.4	29.8	50.8	35.3
Oklahoma	31.0	49.1	14.2	12.7	38.7	39.3	43.9	48.6	47.2	61.5	27.4
Oregon	15.4	23.0	7.3	7.1	20.4	17.2	15.9	22.4	22.9	39.5	47.9
Pennsylvania	47.6	27.2	28.2	21.1	35.6	33.7	19.3	23.0	24.1	48.5	41.4
Rhode Island	26.5	1.0	5.7	6.6	8.8	8.5	0.0	2.0	4.2	16.8	72.6
South Carolina	39.0	41.6	20.4	20.7	45.9	45.8	37.7	50.3	40.5	58.2	28.8
South Dakota	37.8	36.4	9.5	16.8	26.1	24.2	15.4	17.9	40.1	66.5	31.9
Tennessee	28.4	20.2	14.3	10.9	25.2	23.1	21.7	25.2	22.0	30.6	61.4
Texas	38.3	20.1	29.2	19.9	29.2	26.8	18.9	21.9	20.9	48.4	42.8
Utah	53.1	46.6	31.4	32.5	69.7	68.5	75.2	74.0	54.3	73.0	8.3
Vermont	52.1	16.8	27.5	14.1	33.4	26.7	11.0	14.1	18.4	41.4	46.0
Virginia	38.0	27.6	24.6	14.1	38.5	34.5	24.9	33.4	37.1	46.1	37.2
Washington	36.8	33.7	12.1	17.0	41.1	40.0	26.5	35.6	37.9	64.4	26.2
West Virginia	13.5	4.3	0.6	2.3	6.0	8.3	1.7	5.2	2.9	10.3	81.6
Wisconsin	44.2	31.0	18.0	13.8	35.8	31.7	22.7	27.9	28.0	58.4	33.5
Wyoming	30.7	33.6	7.0	13.7	32.1	33.6	32.1	32.3	34.8	63.7	31.9
State Median	34.8	27.6	14.3	14.1	32.4	26.8	20.4	25.2	29.8	50.7	38.4
State Range	6.3-53.1	1.0-49.1	0.6-41.4	2.3-32.5	6.0-69.7	8.3-68.5	0.0-75.2	2.0-74.0	2.9-54.3	8.4-73.0	8.3-81.6

TABLE 27. Percentage of Secondary Schools That Allowed Students to Purchase Less Nutritious Foods and Beverages From Vending Machines or at the School Store, Canteen, or Snack Bar and the Percentage That Did Not Sell Baked Goods,* Salty Snacks,* Candy, Soda Pop or Fruit Drinks,† or Sports Drinks in These Venues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	2% or whole milk (plain or flavored)	Foods or beverages containing caffeine	Ice cream or frozen yogurt*	Water ices or frozen slushes that do not contain juice	Cookies, crackers, cakes, pastries, or other baked goods	Salty snacks	Chocolate candy	Other kinds of candy	Soda pop or fruit drinks	Sports drinks	Did not sell any of these 6 items
LOCAL SURVEYS											
Albuquerque	41.9	8.3	2.9	15.9	19.1	19.1	5.5	5.5	13.6	24.8	59.0
Baltimore	12.6	10.4	6.5	7.7	20.7	24.3	13.0	14.2	15.6	9.2	74.4
Broward County	56.4	51.0	51.4	27.5	71.5	67.9	58.0	65.8	73.8	83.9	12.1
Charlotte-Mecklenberg County	35.1	22.7	21.8	19.4	50.5	49.5	34.6	47.9	38.3	45.2	34.2
Chicago	12.8	4.0	1.0	7.1	12.0	7.0	6.9	10.9	11.0	14.9	75.2
Detroit	21.7	12.7	6.9	11.7	24.0	19.7	8.1	10.3	16.4	25.3	66.5
District of Columbia	28.6	8.5	11.4	8.6	25.7	19.9	17.1	22.8	22.8	11.4	62.8
Fresno	20.3	8.2	15.9	15.9	7.7	15.4	0.0	3.8	4.4	52.7	47.3
Houston	55.6	30.1	50.1	48.7	51.3	54.3	34.2	38.2	35.5	55.6	28.2
Los Angeles	51.9	2.6	14.5	13.7	23.5	14.6	4.0	16.6	10.3	73.4	21.7
Memphis	24.0	12.5	8.8	20.1	14.3	16.0	14.7	14.7	10.6	16.0	76.9
Miami-Dade County	47.7	12.6	33.0	19.6	40.7	29.9	14.6	24.3	17.4	67.8	24.6
Milwaukee	11.7	7.2	2.9	7.4	7.1	8.5	1.3	2.7	10.1	11.5	81.5
New York City	19.1	5.1	8.0	8.4	17.2	18.0	3.8	8.0	12.8	11.8	64.0
Newark	22.0	4.1	13.1	13.6	1.8	6.7	1.8	4.1	14.4	19.8	70.2
Orange County	59.5	6.8	24.4	17.8	39.0	31.5	9.3	11.4	13.4	65.0	30.7
Philadelphia	22.2	4.2	22.2	11.6	16.8	17.6	8.3	12.3	10.7	16.5	67.4
San Diego	32.8	10.4	20.9	22.4	22.4	19.4	9.0	9.0	7.5	55.2	43.3
San Francisco	20.8	5.8	0.0	2.9	11.8	5.8	2.9	0.0	3.2	9.2	75.8
Local Median	24.0	8.3	13.1	13.7	20.7	19.1	8.3	11.4	13.4	24.8	62.8
Local Range	11.7-59.5	2.6-51.0	0.0-51.4	2.9-48.7	1.8-71.5	5.8-67.9	0.0-58.0	0.0-65.8	3.2-73.8	9.2-83.9	12.1-81.5
TERRITORIAL SURVEYS											
American Samoa	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0
Guam	38.5	0.0	0.0	7.7	0.0	0.0	0.0	0.0	7.7	61.5	38.5
Marshall Islands	29.5	24.7	24.7	29.3	31.0	27.8	24.7	27.9	26.3	26.3	64.3
Northern Mariana Islands	37.5	0.0	12.5	25.0	25.0	12.5	12.5	0.0	0.0	50.0	50.0
Palau	45.5	63.6	54.5	45.5	63.6	63.6	63.6	63.6	54.5	63.6	36.4
Territorial Median	37.5	0.0	12.5	25.0	25.0	12.5	12.5	0.0	7.7	50.0	50.0
Territorial Range	0.0-45.5	0.0-63.6	0.0-54.5	0.0-45.5	0.0-63.6	0.0-63.6	0.0-63.6	0.0-63.6	0.0-54.5	0.0-63.6	36.4-100.0
TRIBAL SURVEYS											
Cherokee Nation	35.8	46.6	14.5	21.1	47.9	49.8	48.2	55.6	43.3	63.4	27.4
Nez Perce	14.3	42.9	0.0	14.3	42.9	42.9	42.9	42.9	42.9	57.1	28.6

*That are not low in fat.

†That are not 100% juice.

TABLE 28. Percentage of Secondary Schools That Implemented Strategies to Promote Healthy Eating, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Priced nutritious foods and beverages at a lower cost while increasing the price of less nutritious foods and beverages	Collected suggestions from students, families, and school staff on nutritious food preferences and strategies to promote healthy eating	Provided information to students or families on the nutrition and caloric content of foods available	Conducted taste tests to determine food preferences for nutritious items	Provided opportunities for students to visit the cafeteria to learn about food safety, food preparation, or other nutrition-related topics	At least 3 of these 5 strategies
STATE SURVEYS						
Alabama	8.9	38.6	43.4	18.0	17.0	17.9
Alaska	10.7	44.1	38.6	7.1	24.8	15.6
Arizona	7.7	41.9	39.4	21.9	23.8	21.0
Arkansas	4.0	44.3	43.5	13.0	15.0	13.5
California	9.3	47.3	53.2	30.7	19.6	28.0
Colorado	10.3	50.0	40.7	22.2	14.7	21.9
Connecticut	10.6	56.8	52.6	20.6	19.3	25.7
Delaware	19.1	45.3	68.8	31.6	18.5	31.8
Florida	10.0	51.8	49.4	30.6	28.3	28.5
Georgia	11.3	45.7	44.3	24.7	18.8	23.2
Hawaii	3.8	34.1	33.0	8.0	37.2	11.5
Idaho	17.5	49.5	44.2	13.1	22.6	24.1
Indiana	11.4	49.0	46.8	21.5	15.2	20.6
Iowa	12.8	50.5	45.7	23.2	18.2	22.5
Kansas	13.6	46.9	45.8	23.7	16.8	20.3
Kentucky	7.7	32.9	44.8	14.4	14.1	13.8
Louisiana	8.3	33.3	38.1	12.6	14.7	12.2
Maine	13.1	58.9	57.4	26.5	28.5	33.3
Maryland	9.3	39.4	50.1	19.1	16.3	20.7
Massachusetts	7.5	60.0	54.3	32.3	20.3	29.7
Michigan	12.2	50.4	53.9	25.7	15.0	25.3
Minnesota	19.0	60.6	54.7	28.2	24.5	33.0
Mississippi	7.4	48.2	46.0	14.3	19.0	20.9
Missouri	7.2	48.4	44.1	17.2	18.7	19.3
Montana	7.5	50.9	33.1	12.3	23.9	19.6
Nebraska	15.0	44.7	36.2	13.9	17.0	18.0
Nevada	8.7	30.2	42.9	8.0	17.4	13.7
New Hampshire	10.2	66.3	55.7	35.2	17.4	31.2
New Jersey	11.0	68.5	58.4	34.5	26.9	34.6
New Mexico	11.4	42.2	39.2	14.9	17.6	18.2
New York	11.3	66.8	59.3	33.5	29.0	38.7
North Carolina	6.6	32.9	41.4	16.4	13.0	14.2
North Dakota	10.3	42.3	40.3	16.8	24.0	22.2
Ohio	15.6	46.1	51.1	17.5	20.4	22.7
Oklahoma	20.0	53.3	42.2	15.1	17.6	21.4
Oregon	4.8	39.5	43.1	15.9	24.9	17.4
Pennsylvania	13.3	61.1	65.8	40.2	25.6	37.3
Rhode Island	12.8	52.7	52.9	32.4	30.9	33.4
South Carolina	10.6	42.9	47.4	25.7	16.5	20.7
South Dakota	7.8	39.0	30.1	9.6	14.7	8.6
Tennessee	10.4	44.9	46.6	25.9	23.9	24.1
Texas	9.0	41.6	54.1	21.1	17.8	20.5
Utah	9.5	41.5	38.5	11.1	17.2	13.9
Vermont	12.2	70.8	51.9	54.3	43.5	50.2
Virginia	4.9	44.8	58.5	28.3	17.0	26.6
Washington	8.8	33.4	36.5	18.2	15.7	16.2
West Virginia	0.6	37.6	50.7	15.4	13.1	14.2
Wisconsin	15.2	57.3	58.9	24.9	23.4	30.7
Wyoming	7.6	36.1	44.1	11.4	20.6	12.7
State Median	10.3	45.7	45.8	20.6	18.7	21.0
State Range	0.6-20.0	30.2-70.8	30.1-68.8	7.1-54.3	13.0-43.5	8.6-50.2

TABLE 28. Percentage of Secondary Schools That Implemented Strategies to Promote Healthy Eating, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Priced nutritious foods and beverages at a lower cost while increasing the price of less nutritious foods and beverages	Collected suggestions from students, families, and school staff on nutritious food preferences and strategies to promote healthy eating	Provided information to students or families on the nutrition and caloric content of foods available	Conducted taste tests to determine food preferences for nutritious items	Provided opportunities for students to visit the cafeteria to learn about food safety, food preparation, or other nutrition-related topics	At least 3 of these 5 strategies
LOCAL SURVEYS						
Albuquerque	5.5	35.2	37.9	18.8	10.7	21.7
Baltimore	6.2	32.1	40.0	16.9	17.4	17.1
Broward County	12.6	45.1	59.2	22.2	25.0	26.9
Charlotte-Mecklenberg County	9.8	30.4	45.1	21.1	19.0	15.4
Chicago	4.0	36.6	40.0	10.8	14.3	10.0
Detroit	12.4	50.1	56.0	31.8	27.2	34.0
District of Columbia	20.0	47.1	42.8	24.2	12.0	21.2
Fresno	3.8	37.4	53.3	33.0	28.6	36.8
Houston	9.9	30.8	42.1	9.8	12.2	16.0
Los Angeles	8.1	44.4	50.3	31.5	28.5	29.8
Memphis	14.2	44.9	44.6	38.4	35.0	31.4
Miami-Dade County	8.3	51.5	56.7	19.8	30.1	24.3
Milwaukee	4.6	40.1	49.6	17.7	15.2	15.2
New York City	5.1	55.5	51.3	29.5	36.4	34.4
Newark	5.8	80.5	49.8	29.4	38.1	33.3
Orange County	13.0	50.2	50.1	58.2	15.2	30.6
Philadelphia	9.5	45.9	56.3	43.4	24.2	35.4
San Diego	12.1	54.5	53.0	31.3	23.9	34.8
San Francisco	12.9	62.6	67.5	21.7	34.1	31.2
Local Median	9.5	45.1	50.1	24.2	24.2	29.8
Local Range	3.8-20.0	30.4-80.5	37.9-67.5	9.8-58.2	10.7-38.1	10.0-36.8
TERRITORIAL SURVEYS						
American Samoa	3.8	53.8	53.8	34.6	64.0	46.2
Guam	0.0	15.4	38.5	23.1	16.7	15.4
Marshall Islands	38.5	63.3	67.7	38.5	35.4	52.4
Northern Mariana Islands	14.3	71.4	85.7	42.9	57.1	71.4
Palau	9.1	63.6	63.6	45.5	54.5	45.5
Territorial Median	9.1	63.3	63.6	38.5	54.5	46.2
Territorial Range	0.0-38.5	15.4-71.4	38.5-85.7	23.1-45.5	16.7-64.0	15.4-71.4
TRIBAL SURVEYS						
Cherokee Nation	18.6	59.4	42.2	12.2	21.8	24.9
Nez Perce	14.3	42.9	28.6	28.6	14.3	14.3

TABLE 29. Percentage of Secondary Schools That Promoted Candy, Meals from Fast-Food Restaurants, or Soft Drinks Through the Distribution of Products* to Students; the Percentage That Prohibited Advertisements for Candy, Fast Food Restaurants, or Soft Drinks in Specific Locations; and the Percentage That Prohibited All Forms of Advertising and Promotion,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Promoted candy, meals from fast food restaurants, or soft drinks through the distribution of products to students	Prohibited advertisements for candy, fast food restaurants, or soft drinks				Prohibited all forms of advertising and promotion
		In the school building	On school grounds [‡]	On school buses or other vehicles used to transport students	In school publications	
STATE SURVEYS						
Alabama	9.2	81.5	54.7	76.3	70.1	45.3
Alaska	2.4	67.6	66.8	67.1	65.9	61.4
Arizona	2.3	75.0	70.0	75.9	70.3	63.0
Arkansas	2.2	57.2	45.9	62.2	49.6	39.2
California	3.3	81.7	76.9	81.8	71.7	65.1
Colorado	3.6	59.4	56.2	61.5	59.8	44.4
Connecticut	0.4	84.4	79.9	81.9	80.5	71.7
Delaware	0.0	69.1	61.4	72.1	67.1	58.4
Florida	2.7	65.9	55.0	73.1	60.1	46.7
Georgia	4.1	64.7	48.6	73.3	56.5	38.4
Hawaii	2.4	83.0	79.2	78.2	75.0	67.7
Idaho	3.4	58.0	52.7	69.7	58.6	41.5
Indiana	2.2	54.7	42.5	67.1	46.9	33.1
Iowa	1.6	47.7	40.7	59.2	49.9	34.5
Kansas	2.5	48.2	42.0	56.8	45.2	33.5
Kentucky	1.5	58.9	44.7	70.1	52.5	38.6
Louisiana	0.9	79.9	68.8	78.8	70.3	61.6
Maine	1.2	81.5	79.9	83.9	79.9	72.0
Maryland	2.6	67.3	63.6	72.5	61.2	51.2
Massachusetts	0.5	81.6	79.1	80.4	80.4	73.0
Michigan	2.0	64.4	57.5	68.3	60.4	49.4
Minnesota	2.4	67.9	59.9	74.1	68.8	53.8
Mississippi	3.2	75.9	60.0	78.7	68.0	51.3
Missouri	2.3	53.2	45.0	61.4	50.7	36.2
Montana	0.5	51.9	47.0	63.7	56.2	37.8
Nebraska	1.4	43.1	40.3	53.7	45.6	32.1
Nevada	3.8	77.1	68.0	80.8	68.4	59.8
New Hampshire	2.4	77.0	66.8	80.8	77.4	59.1
New Jersey	0.7	82.0	79.5	79.2	80.8	71.6
New Mexico	2.0	71.8	68.1	74.9	70.1	57.4
New York	0.9	80.7	78.9	79.0	79.5	71.1
North Carolina	2.3	68.4	56.8	77.6	63.4	49.1
North Dakota	1.2	51.4	42.9	50.9	46.9	34.8
Ohio	5.1	60.0	42.2	64.6	54.0	32.9
Oklahoma	3.4	54.7	44.3	63.1	52.9	37.3
Oregon	2.7	71.3	58.3	74.2	67.1	49.8
Pennsylvania	2.2	73.0	64.8	75.2	70.0	56.1
Rhode Island	2.2	89.7	79.9	87.4	82.6	72.4
South Carolina	4.2	66.2	57.1	75.5	59.3	46.6
South Dakota	1.4	47.6	42.4	49.3	49.0	34.9
Tennessee	2.8	68.6	53.4	74.2	59.5	41.5
Texas	2.8	68.8	62.1	71.9	60.2	52.6
Utah	2.8	64.9	55.4	76.8	60.2	46.9
Vermont	0.9	72.3	63.4	71.1	73.4	58.8
Virginia	1.2	68.0	54.8	78.3	60.1	46.4
Washington	4.3	67.7	62.9	74.5	62.8	52.1
West Virginia	4.0	70.6	55.3	74.9	61.6	46.8
Wisconsin	4.2	60.2	54.4	63.8	61.4	43.1
Wyoming	0.7	48.7	44.9	60.2	50.5	38.9
State Median	2.3	67.9	57.1	74.1	61.4	49.1
State Range	0.0-9.2	43.1-89.7	40.3-79.9	49.3-87.4	45.2-82.6	32.1-73.0

TABLE 29. Percentage of Secondary Schools That Promoted Candy, Meals from Fast-Food Restaurants, or Soft Drinks Through the Distribution of Products* to Students; the Percentage That Prohibited Advertisements for Candy, Fast Food Restaurants, or Soft Drinks in Specific Locations; and the Percentage That Prohibited All Forms of Advertising and Promotion,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Promoted candy, meals from fast food restaurants, or soft drinks through the distribution of products to students	Prohibited advertisements for candy, fast food restaurants, or soft drinks				
		In the school building	On school grounds‡	On school buses or other vehicles used to transport students	In school publications	Prohibited all forms of advertising and promotion
LOCAL SURVEYS						
Albuquerque	0.0	83.8	86.4	78.3	83.6	69.4
Baltimore	2.7	61.2	67.1	63.2	65.6	53.9
Broward County	1.4	74.6	66.5	73.6	61.6	53.6
Charlotte-Mecklenberg County	5.9	62.2	58.7	75.3	62.6	45.5
Chicago	5.1	77.0	71.2	65.2	75.0	57.1
Detroit	5.8	73.9	72.4	72.1	72.4	66.3
District of Columbia	0.0	76.6	67.7	64.8	67.8	55.9
Fresno	0.0	76.4	67.6	80.8	68.1	59.3
Houston	11.6	64.1	59.1	57.4	57.4	51.8
Los Angeles	2.6	90.2	87.8	86.1	82.8	75.7
Memphis	3.7	78.4	74.8	79.6	77.8	64.8
Miami-Dade County	1.8	80.3	72.7	76.1	66.8	61.2
Milwaukee	3.2	72.4	68.3	65.2	73.5	54.3
New York City	1.5	83.3	80.6	75.4	79.7	70.7
Newark	0.0	79.4	73.2	65.1	73.5	61.8
Orange County	2.2	73.8	69.4	75.5	59.0	45.5
Philadelphia	1.5	82.8	80.5	74.6	80.0	68.7
San Diego	0.0	100.0	100.0	100.0	100.0	100.0
San Francisco	3.3	96.7	90.5	90.5	90.8	81.3
Local Median	2.2	77.0	72.4	75.3	73.5	61.2
Local Range	0.0-11.6	61.2-100.0	58.7-100.0	57.4-100.0	57.4-100.0	45.5-100.0
TERRITORIAL SURVEYS						
American Samoa	57.7	42.3	38.5	23.1	38.5	3.8
Guam	7.7	84.6	84.6	84.6	84.6	76.9
Marshall Islands	7.6	54.3	49.2	40.2	37.2	31.7
Northern Mariana Islands	0.0	62.5	62.5	62.5	62.5	62.5
Palau	18.2	0.0	0.0	0.0	9.1	0.0
Territorial Median	7.7	54.3	49.2	40.2	38.5	31.7
Territorial Range	0.0-57.7	0.0-84.6	0.0-84.6	0.0-84.6	9.1-84.6	0.0-76.9
TRIBAL SURVEYS						
Cherokee Nation	5.5	55.1	46.9	62.8	52.7	39.1
Nez Perce	0.0	57.1	57.1	85.7	85.7	57.1

*Such as t-shirts, hats, and book covers.

†Prohibited all advertisements for candy, fast food restaurants, and soft drinks in school buildings, on school grounds, on school buses or other vehicles used to transport students, in school publications, and through sponsorship of school events, and did not promote candy, meals from fast food restaurants, or soft drinks through the distribution of products to students.

‡Including on the outside of the school building, on playing fields, or other areas of the campus.

TABLE 30. Percentage of Secondary Schools That Had a Policy Prohibiting Tobacco Use and the Percentage That Prohibited All Tobacco Use in All Locations,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Had a policy prohibiting tobacco use	Prohibited all tobacco use at all times in all locations
STATE SURVEYS		
Alabama	95.2	63.9
Alaska	99.3	46.6
Arizona	95.4	57.8
Arkansas	98.7	57.9
California	97.9	65.7
Colorado	97.4	57.9
Connecticut	95.6	53.7
Delaware	100.0	69.7
Florida	95.5	51.9
Georgia	97.7	63.4
Hawaii	98.9	76.1
Idaho	98.4	44.7
Indiana	99.3	53.2
Iowa	99.6	60.5
Kansas	97.8	41.3
Kentucky	95.8	23.5
Louisiana	99.6	60.0
Maine	99.6	64.7
Maryland	96.6	64.0
Massachusetts	95.9	58.7
Michigan	95.3	55.1
Minnesota	97.1	51.0
Mississippi	99.5	71.8
Missouri	96.7	33.0
Montana	98.4	50.3
Nebraska	97.1	31.4
Nevada	96.9	52.3
New Hampshire	98.8	61.2
New Jersey	96.3	51.3
New Mexico	99.1	55.4
New York	90.2	56.9
North Carolina	98.7	76.9
North Dakota	98.9	42.0
Ohio	98.4	43.1
Oklahoma	96.9	42.9
Oregon	96.5	58.8
Pennsylvania	97.7	59.9
Rhode Island	96.9	61.7
South Carolina	97.3	65.0
South Dakota	97.3	32.0
Tennessee	99.0	45.0
Texas	97.9	65.9
Utah	98.9	57.4
Vermont	100.0	60.8
Virginia	97.5	54.3
Washington	99.0	68.4
West Virginia	99.2	79.3
Wisconsin	98.0	55.3
Wyoming	100.0	42.0
State Median	97.9	57.4
State Range	90.2-100.0	23.5-79.3

TABLE 30. Percentage of Secondary Schools That Had a Policy Prohibiting Tobacco Use and the Percentage That Prohibited All Tobacco Use in All Locations,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Had a policy prohibiting tobacco use	Prohibited all tobacco use at all times in all locations
LOCAL SURVEYS		
Albuquerque	97.4	47.1
Baltimore	83.5	44.3
Broward County	95.5	43.6
Charlotte-Mecklenberg County	98.0	77.0
Chicago	72.9	29.3
Detroit	88.7	49.5
District of Columbia	85.8	32.4
Fresno	96.2	76.0
Houston	97.2	75.6
Los Angeles	98.4	66.8
Memphis	91.9	66.1
Miami-Dade County	97.1	41.6
Milwaukee	87.6	33.9
New York City	78.0	33.9
Newark	90.1	45.5
Orange County	97.8	62.1
Philadelphia	81.0	43.4
San Diego	98.5	95.5
San Francisco	100.0	67.0
Local Median	95.5	47.1
Local Range	72.9-100.0	29.3-95.5
TERRITORIAL SURVEYS		
American Samoa	100.0	0.0
Guam	100.0	50.0
Marshall Islands	95.2	14.9
Northern Mariana Islands	100.0	60.0
Palau	100.0	0.0
Territorial Median	100.0	14.9
Territorial Range	95.2-100.0	0.0-60.0
TRIBAL SURVEYS		
Cherokee Nation	96.4	38.4
Nez Perce	100.0	40.0

*Prohibited the use of all tobacco, including cigarettes, smokeless tobacco, cigars, and pipes, by students, faculty and school staff, and visitors, in school buildings, outside on school grounds, on school buses or other vehicles used to transport students, and at off-campus, school-sponsored events during school hours and non-school hours.

TABLE 31a. Percentage of Secondary Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Notified parents or guardians	Referred to a school counselor	Referred to a school administrator	Encouraged to participate in an assistance, education, or cessation program	Required to participate in an assistance, education, or cessation program
STATE SURVEYS					
Alabama	98.7	69.0	95.2	45.9	29.6
Alaska	99.0	64.3	98.7	69.2	42.0
Arizona	99.7	68.1	99.7	54.9	34.2
Arkansas	99.5	72.2	98.6	48.5	28.6
California	99.8	81.7	97.7	70.2	50.5
Colorado	95.5	77.4	97.7	69.8	36.5
Connecticut	98.3	88.2	98.1	69.3	34.7
Delaware	100.0	82.1	96.9	74.3	36.8
Florida	99.7	80.9	98.8	64.4	46.6
Georgia	100.0	76.4	100.0	45.4	24.9
Hawaii	100.0	91.7	100.0	70.7	33.8
Idaho	98.4	86.1	98.8	69.9	61.9
Indiana	99.7	71.8	99.3	63.7	46.0
Iowa	99.2	78.8	99.2	63.2	30.8
Kansas	99.6	73.6	99.6	52.9	30.7
Kentucky	98.8	68.9	99.6	60.8	43.6
Louisiana	99.6	72.7	99.2	40.0	28.5
Maine	99.2	91.9	99.2	83.5	55.7
Maryland	98.7	92.9	99.1	84.4	85.8
Massachusetts	98.7	85.5	99.3	66.6	35.0
Michigan	99.3	72.5	99.6	63.8	32.1
Minnesota	99.6	77.9	99.6	66.8	37.5
Mississippi	99.1	68.6	99.1	38.7	20.4
Missouri	99.3	64.9	99.3	40.9	17.0
Montana	100.0	84.6	99.1	65.2	52.0
Nebraska	100.0	75.7	100.0	54.2	30.6
Nevada	100.0	68.2	99.4	67.5	44.3
New Hampshire	100.0	88.9	100.0	73.5	51.2
New Jersey	98.6	91.7	98.6	72.6	48.6
New Mexico	99.5	75.5	99.5	58.6	45.0
New York	98.5	90.2	98.2	71.2	36.3
North Carolina	98.3	74.6	98.3	58.3	43.0
North Dakota	99.3	76.4	98.6	64.8	35.0
Ohio	99.4	73.6	99.4	68.4	44.6
Oklahoma	99.4	66.1	99.1	46.9	22.5
Oregon	97.2	77.1	97.5	75.9	49.6
Pennsylvania	99.1	86.3	98.5	73.2	46.3
Rhode Island	100.0	93.7	100.0	77.8	51.6
South Carolina	99.7	78.6	100.0	62.7	44.2
South Dakota	99.3	73.2	99.3	52.9	26.2
Tennessee	100.0	72.0	99.7	54.4	38.7
Texas	99.7	76.3	99.7	46.9	22.3
Utah	100.0	80.2	100.0	85.1	77.1
Vermont	98.5	94.0	99.3	81.7	58.4
Virginia	99.7	80.4	99.7	62.8	47.9
Washington	98.6	88.5	100.0	80.6	62.0
West Virginia	100.0	84.8	99.4	76.3	66.5
Wisconsin	99.7	85.0	98.9	73.8	40.5
Wyoming	100.0	78.2	100.0	71.1	55.5
State Median	99.5	77.9	99.3	66.6	42.0
State Range	95.5-100.0	64.3-94.0	95.2-100.0	38.7-85.1	17.0-85.8

TABLE 31a. Percentage of Secondary Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Notified parents or guardians	Referred to a school counselor	Referred to a school administrator	Encouraged to participate in an assistance, education, or cessation program	Required to participate in an assistance, education, or cessation program
LOCAL SURVEYS					
Albuquerque	97.1	80.7	97.1	68.1	53.3
Baltimore	95.1	79.0	96.4	53.9	32.9
Broward County	98.3	89.9	100.0	64.7	30.2
Charlotte-Mecklenberg County	100.0	88.4	100.0	69.8	75.2
Chicago	98.9	79.6	96.7	49.5	29.6
Detroit	98.8	84.6	98.8	48.6	28.3
District of Columbia	100.0	84.7	100.0	51.3	36.5
Fresno	100.0	87.4	95.6	62.9	29.7
Houston	100.0	78.9	98.7	45.5	27.5
Los Angeles	99.3	90.1	84.8	89.4	78.8
Memphis	100.0	82.1	100.0	71.4	64.6
Miami-Dade County	100.0	99.2	100.0	78.2	48.7
Milwaukee	97.2	71.4	97.2	51.7	31.2
New York City	96.2	95.1	96.2	65.2	39.6
Newark	97.8	97.8	97.7	76.3	81.7
Orange County	100.0	88.8	100.0	70.7	55.6
Philadelphia	100.0	87.3	91.2	59.6	30.6
San Diego	100.0	100.0	100.0	83.6	67.7
San Francisco	97.0	94.0	94.0	78.5	84.5
Local Median	99.3	87.4	97.7	65.2	39.6
Local Range	95.1-100.0	71.4-100.0	84.8-100.0	45.5-89.4	27.5-84.5
TERRITORIAL SURVEYS					
American Samoa	100.0	100.0	100.0	80.8	76.9
Guam	100.0	84.6	100.0	84.6	61.5
Marshall Islands	97.1	89.7	95.5	71.6	65.2
Northern Mariana Islands	100.0	87.5	100.0	57.1	66.7
Palau	100.0	100.0	100.0	9.1	9.1
Territorial Median	100.0	89.7	100.0	71.6	65.2
Territorial Range	97.1-100.0	84.6-100.0	95.5-100.0	9.1-84.6	9.1-76.9
TRIBAL SURVEYS					
Cherokee Nation	98.5	67.8	97.3	47.5	18.8
Nez Perce	100.0	71.4	100.0	85.7	57.1

TABLE 31b. Percentage of Secondary Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Referred to legal authorities	Placed in detention	Not allowed to participate in extra-curricular activities or interscholastic sports	Given in-school suspension	Suspended from school	Expelled from school	Reassigned to an alternative school
STATE SURVEYS							
Alabama	38.3	59.3	67.9	68.0	83.7	24.2	46.0
Alaska	59.9	67.7	90.3	67.5	79.1	5.7	3.8
Arizona	51.2	64.4	71.8	69.5	85.1	19.2	17.2
Arkansas	35.7	59.1	56.0	74.8	76.9	13.3	19.8
California	48.5	70.7	69.7	72.1	88.2	16.3	17.0
Colorado	50.8	73.8	79.0	76.2	74.2	12.2	7.0
Connecticut	36.8	69.8	58.8	79.9	54.2	4.7	3.2
Delaware	16.6	67.8	57.5	76.1	79.0	8.4	13.0
Florida	66.2	65.4	54.6	70.6	78.4	11.9	16.4
Georgia	27.1	55.7	61.1	85.7	79.2	16.8	29.2
Hawaii	12.7	78.9	57.5	63.1	75.3	1.4	2.4
Idaho	92.0	67.0	85.8	73.2	81.0	18.7	17.0
Indiana	63.7	53.9	83.7	63.1	83.9	26.2	21.1
Iowa	80.4	64.6	94.0	78.4	68.0	4.4	5.6
Kansas	71.2	61.8	89.5	73.4	76.9	14.8	6.7
Kentucky	23.6	78.8	57.0	85.1	55.4	3.1	17.8
Louisiana	25.2	58.3	55.0	63.1	86.6	24.5	31.7
Maine	73.2	57.1	91.7	62.8	77.9	6.5	2.1
Maryland	76.1	89.8	81.4	78.8	86.4	62.9	64.8
Massachusetts	22.2	70.2	69.6	63.8	73.2	2.0	2.6
Michigan	64.5	51.3	82.7	53.2	92.4	11.0	6.7
Minnesota	76.9	62.7	89.6	72.8	80.1	3.3	3.3
Mississippi	33.7	64.0	60.3	69.7	83.3	15.8	36.8
Missouri	46.8	77.8	80.7	82.7	70.1	10.9	12.5
Montana	84.7	69.0	95.7	76.5	77.7	9.7	4.4
Nebraska	66.0	73.8	92.0	77.7	68.1	11.4	6.7
Nevada	28.6	73.0	79.8	72.4	77.2	15.5	20.9
New Hampshire	82.5	64.2	83.8	71.6	85.2	2.8	1.1
New Jersey	41.5	69.7	67.8	63.2	75.7	3.8	4.1
New Mexico	20.4	78.9	72.4	76.2	60.5	7.7	9.5
New York	20.3	71.7	69.9	79.9	60.7	5.2	6.7
North Carolina	21.9	62.7	48.9	73.4	76.3	8.3	13.1
North Dakota	62.9	63.3	96.0	75.0	62.3	8.0	0.6
Ohio	43.1	50.9	82.8	57.8	87.4	16.4	19.8
Oklahoma	44.8	73.5	73.6	75.9	74.0	20.6	14.9
Oregon	62.3	62.0	74.2	72.5	76.2	16.7	13.2
Pennsylvania	73.6	57.9	67.2	69.5	65.8	6.8	10.9
Rhode Island	27.6	75.8	62.1	65.0	74.5	5.6	4.4
South Carolina	51.2	58.3	57.0	72.6	91.8	18.2	26.6
South Dakota	73.8	74.7	96.5	82.9	62.8	7.3	1.6
Tennessee	81.7	53.1	57.4	67.5	69.0	16.7	33.7
Texas	69.6	74.3	75.8	90.6	55.9	9.3	34.7
Utah	97.2	57.6	77.4	59.7	87.2	13.9	14.0
Vermont	68.3	56.9	83.2	73.3	63.2	2.3	2.6
Virginia	68.4	52.8	76.5	66.5	87.2	9.3	14.0
Washington	44.8	55.8	82.9	59.2	83.5	9.6	6.7
West Virginia	76.8	57.5	53.1	56.9	71.8	5.9	13.4
Wisconsin	85.6	63.9	91.7	70.9	74.7	5.4	3.0
Wyoming	92.8	61.3	91.0	63.7	74.0	9.0	2.6
State Median	59.9	64.2	75.8	72.4	76.9	9.6	12.5
State Range	12.7-97.2	50.9-89.8	48.9-96.5	53.2-90.6	54.2-92.4	1.4-62.9	0.6-64.8

TABLE 31b. Percentage of Secondary Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Referred to legal authorities	Placed in detention	Not allowed to participate in extra-curricular activities or interscholastic sports	Given in-school suspension	Suspended from school	Expelled from school	Reassigned to an alternative school
LOCAL SURVEYS							
Albuquerque	33.8	78.1	75.2	80.9	64.3	8.1	5.5
Baltimore	23.4	72.5	58.8	43.3	61.2	7.9	7.8
Broward County	49.4	78.4	48.5	89.1	74.0	9.1	11.0
Charlotte-Mecklenberg County	37.8	79.0	61.7	79.8	68.6	7.9	9.9
Chicago	29.4	64.3	59.7	79.4	77.0	9.7	8.6
Detroit	36.7	51.5	66.5	50.8	94.9	23.8	22.5
District of Columbia	46.9	75.8	68.7	84.9	81.2	24.9	18.8
Fresno	20.6	91.8	82.6	76.4	87.9	0.0	0.0
Houston	87.3	89.9	79.3	86.6	90.0	28.3	58.4
Los Angeles	59.1	74.4	52.3	72.3	72.7	11.2	18.2
Memphis	60.7	50.0	61.4	55.9	90.6	35.8	30.7
Miami-Dade County	47.8	81.8	77.9	89.9	82.5	20.3	16.8
Milwaukee	35.0	46.2	58.5	41.3	86.8	13.9	12.6
New York City	26.9	72.5	51.1	75.9	46.3	7.5	10.7
Newark	35.5	75.4	52.4	70.7	77.3	6.4	15.2
Orange County	61.1	65.9	66.7	80.4	82.6	8.7	15.0
Philadelphia	25.3	86.9	58.7	71.0	57.5	8.1	9.7
San Diego	54.5	66.2	71.6	67.2	92.4	16.4	7.5
San Francisco	13.0	54.3	54.3	54.3	41.1	0.0	0.0
Local Median	36.7	74.4	61.4	75.9	77.3	9.1	11.0
Local Range	13.0-87.3	46.3-91.8	48.5-82.6	41.3-89.9	41.1-94.9	0.0-35.8	0.0-58.4
TERRITORIAL SURVEYS							
American Samoa	46.2	88.5	96.2	96.2	57.7	23.1	15.4
Guam	23.1	50.0	84.6	30.8	100.0	0.0	53.8
Marshall Islands	36.5	61.0	60.7	59.0	39.8	22.5	17.9
Northern Mariana Islands	50.0	87.5	100.0	85.7	100.0	37.5	25.0
Palau	18.2	90.9	100.0	100.0	90.9	81.8	72.7
Territorial Median	36.5	87.5	96.2	85.7	90.9	23.1	25.0
Territorial Range	18.2-50.0	50.0-90.9	60.7-100.0	30.8-100.0	39.8-100.0	0.0-81.8	15.4-72.7
TRIBAL SURVEYS							
Cherokee Nation	48.2	72.8	67.1	78.8	72.0	24.1	14.9
Nez Perce	100.0	28.6	100.0	50.0	100.0	0.0	0.0

TABLE 32. Percentage of Secondary Schools That Took Specific Actions to Support a Tobacco-Free Environment Policy, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Informed groups about policy prohibiting their use*			Included guidelines on what actions should be taken**	Always or almost always notified parents or guardians†	Used effect or severity of the violation or repeat offender status to determine actions taken**	Used remedial rather than punitive sanctions for violators‡	Had an individual responsible for enforcing policy§	Posted signs marking a tobacco-free school zone¶	Met all 7 criteria**
	Students	Faculty and staff	Visitors							
STATE SURVEYS										
Alabama	100.0	98.2	93.4	95.9	88.1	77.4	34.2	48.0	82.0	5.0
Alaska	98.4	95.3	86.1	98.3	97.1	86.0	48.6	69.2	82.3	17.3
Arizona	97.2	95.5	91.3	96.2	96.7	85.2	42.7	55.6	88.3	15.1
Arkansas	99.6	98.3	94.3	97.6	95.8	82.0	32.5	60.1	94.1	9.9
California	98.9	95.6	92.0	96.4	98.8	85.1	61.0	52.6	88.0	22.3
Colorado	99.1	96.4	89.1	96.2	92.3	86.8	42.0	54.2	94.3	13.0
Connecticut	96.5	93.6	84.3	92.5	91.3	78.4	62.9	43.8	40.9	5.2
Delaware	100.0	96.8	84.8	96.8	96.9	77.6	58.9	46.0	66.6	17.1
Florida	98.0	93.0	86.0	97.5	98.3	75.6	56.8	42.0	72.5	11.0
Georgia	99.6	98.7	93.3	98.0	98.0	90.1	29.7	46.5	82.9	6.3
Hawaii	100.0	98.5	95.8	93.1	94.5	83.9	46.2	55.2	84.7	11.4
Idaho	97.6	94.3	77.2	98.8	95.5	86.5	65.8	59.9	70.1	19.7
Indiana	100.0	97.0	92.9	94.9	99.0	87.2	46.3	52.2	75.3	14.4
Iowa	98.8	96.6	94.9	94.2	97.9	80.8	43.9	57.7	87.0	16.7
Kansas	98.8	97.5	84.7	96.1	98.6	84.5	39.1	60.4	73.4	12.2
Kentucky	99.6	90.9	83.8	97.6	94.3	85.1	44.0	49.4	61.9	7.5
Louisiana	100.0	98.2	91.0	98.7	98.3	72.5	45.7	56.5	87.3	15.6
Maine	97.6	96.3	89.5	97.6	97.5	90.4	75.9	60.7	84.0	29.6
Maryland	98.2	92.8	87.8	99.1	96.5	89.6	84.7	50.0	80.1	6.8
Massachusetts	97.8	95.7	84.8	96.8	92.9	79.8	56.0	54.5	75.4	13.1
Michigan	98.5	95.9	83.8	98.1	97.9	86.8	44.6	55.0	66.1	10.9
Minnesota	98.5	95.6	84.6	96.5	96.1	87.4	54.0	57.4	73.4	17.3
Mississippi	98.2	97.2	95.7	97.6	92.2	74.1	32.9	51.3	95.4	11.7
Missouri	99.7	96.4	81.4	98.6	96.1	90.6	25.7	56.2	69.3	7.7
Montana	98.9	97.3	88.9	96.8	99.5	87.2	59.5	60.9	92.2	23.9
Nebraska	99.0	92.9	73.4	96.9	98.1	86.3	37.9	63.0	70.5	11.2
Nevada	98.6	94.6	84.9	97.3	94.4	85.8	42.3	50.2	65.4	9.1
New Hampshire	99.4	98.2	93.3	96.4	99.4	74.6	69.8	56.7	88.1	20.7
New Jersey	98.2	96.1	88.7	96.8	96.6	80.3	81.2	54.1	72.7	20.6
New Mexico	98.6	96.2	86.3	95.2	90.9	82.6	51.0	54.0	76.8	15.5
New York	98.4	97.5	92.3	95.9	94.6	83.2	62.2	42.3	75.1	11.2
North Carolina	99.2	99.2	98.4	97.9	94.0	82.3	46.0	48.4	92.5	13.7
North Dakota	99.4	96.8	89.9	93.2	99.3	71.3	60.4	61.6	71.4	17.2
Ohio	96.9	93.5	87.2	94.3	96.4	85.7	49.0	60.2	75.6	16.3
Oklahoma	99.7	98.1	90.9	94.2	94.9	81.8	27.7	60.0	79.6	12.0
Oregon	99.2	97.3	89.9	94.6	91.6	89.7	55.0	50.7	82.8	15.1
Pennsylvania	97.4	93.9	87.9	98.0	97.3	79.0	61.5	58.7	75.1	23.1
Rhode Island	100.0	96.7	81.9	95.5	94.8	80.6	71.7	48.1	76.2	14.7
South Carolina	99.4	98.5	94.3	98.5	95.2	82.0	44.9	44.4	69.3	13.4
South Dakota	99.4	94.6	84.6	99.5	97.4	88.2	42.3	70.5	80.7	13.0
Tennessee	100.0	97.5	94.9	98.9	97.9	79.6	42.3	60.1	88.6	12.7
Texas	99.2	97.4	93.3	95.5	98.0	79.4	31.2	43.8	86.4	6.7
Utah	99.5	93.9	76.4	99.5	99.5	90.7	64.4	54.5	68.6	15.5
Vermont	99.2	98.3	91.5	97.0	97.6	83.5	85.2	66.2	68.4	30.9
Virginia	100.0	98.5	91.2	98.8	99.2	84.3	48.2	43.2	73.1	11.7
Washington	100.0	98.2	89.5	97.6	97.2	94.6	73.6	58.6	91.7	29.6
West Virginia	99.4	99.4	98.3	97.4	98.2	85.4	76.3	57.4	92.6	34.0
Wisconsin	97.9	94.7	87.6	94.9	96.8	80.6	52.2	59.5	82.4	18.4
Wyoming	96.6	94.3	81.8	95.6	97.5	88.7	62.7	65.5	84.7	24.5
State Median	99.1	96.6	89.1	96.8	96.9	84.3	49.0	55.2	79.6	14.4
State Range	96.5-100.0	90.9-99.4	73.4-98.4	92.5-99.5	88.1-99.5	71.3-94.6	25.7-85.2	42.0-70.5	40.9-95.4	5.0-34.0

TABLE 32. Percentage of Secondary Schools That Took Specific Actions to Support a Tobacco-Free Environment Policy, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Informed groups about policy prohibiting their use*			Included guidelines on what actions should be taken*†	Always or almost always notified parents or guardians†	Used effect or severity of the violation or repeat offender status to determine actions taken*†	Used remedial rather than punitive sanctions for violators‡	Had an individual responsible for enforcing policy§	Posted signs marking a tobacco-free school zone¶	Met all 7 criteria**
	Students	Faculty and staff	Visitors							
LOCAL SURVEYS										
Albuquerque	100.0	94.6	82.4	97.1	88.8	77.7	54.5	40.7	86.4	11.3
Baltimore	92.3	86.2	79.0	96.9	91.2	87.9	49.2	62.8	69.0	10.8
Broward County	100.0	92.1	78.8	100.0	96.7	79.1	56.7	41.4	61.7	8.1
Charlotte-Mecklenberg County	100.0	97.9	89.4	100.0	96.2	87.5	73.1	47.8	86.3	23.2
Chicago	95.6	94.3	74.4	95.8	97.8	82.7	45.1	49.5	46.5	8.0
Detroit	98.6	96.0	92.9	98.7	98.8	77.9	59.9	58.8	63.9	9.0
District of Columbia	93.1	82.2	72.9	93.3	100.0	86.7	56.0	32.2	57.4	9.1
Fresno	100.0	96.0	95.8	100.0	100.0	96.2	65.7	46.3	92.0	20.3
Houston	98.5	91.4	88.4	100.0	93.8	76.3	60.8	45.8	78.5	10.8
Los Angeles	98.4	97.4	93.5	95.8	96.7	83.1	82.6	45.6	91.4	24.8
Memphis	100.0	98.2	98.2	98.4	96.4	75.9	63.4	48.5	91.4	22.4
Miami-Dade County	98.0	98.0	85.7	98.0	100.0	82.7	84.4	40.7	87.1	19.5
Milwaukee	98.5	98.5	89.4	95.4	94.3	70.2	48.6	59.8	70.5	18.2
New York City	95.3	93.1	87.1	93.1	91.2	77.2	75.5	43.2	57.3	7.0
Newark	100.0	100.0	81.1	100.0	97.8	94.9	91.5	43.4	62.7	13.4
Orange County	97.6	97.6	90.5	97.7	100.0	89.2	53.3	30.3	84.8	14.1
Philadelphia	96.6	92.9	82.7	94.7	92.6	80.9	63.2	45.8	48.8	11.4
San Diego	100.0	100.0	100.0	98.5	100.0	92.5	94.0	37.9	98.5	30.3
San Francisco	100.0	97.0	94.0	96.8	85.0	90.7	97.0	59.7	81.3	32.3
Local Median	98.5	96.0	88.4	97.7	96.7	82.7	63.2	45.8	78.5	13.4
Local Range	92.3-100.0	82.2-100.0	72.9-100.0	93.1-100.0	85.0-100.0	70.2-96.2	45.1-97.0	30.3-62.8	46.5-98.5	7.0-32.3
TERRITORIAL SURVEYS										
American Samoa	100.0	100.0	100.0	100.0	96.2	100.0	100.0	57.7	26.9	7.7
Guam	100.0	92.3	92.3	100.0	100.0	92.3	53.8	63.6	100.0	33.3
Marshall Islands	100.0	92.9	91.8	96.5	88.3	83.0	88.3	97.7	69.8	39.3
Northern Mariana Islands	100.0	85.7	71.4	100.0	100.0	100.0	85.7	50.0	57.1	16.7
Palau	100.0	60.0	40.0	100.0	100.0	100.0	54.5	88.9	18.2	0.0
Territorial Median	100.0	92.3	91.8	100.0	100.0	100.0	85.7	63.6	57.1	16.7
Territorial Range	100.0-100.0	60.0-100.0	40.0-100.0	96.5-100.0	88.3-100.0	83.0-100.0	53.8-100.0	50.0-97.7	18.2-100.0	0.0-39.3
TRIBAL SURVEYS										
Cherokee Nation	100.0	96.3	85.0	93.9	93.1	82.7	33.5	67.5	81.9	11.4
Nez Perce	100.0	60.0	40.0	100.0	71.4	57.1	57.1	100.0	100.0	33.3

*Among schools that have adopted a policy prohibiting tobacco use.

†When students are caught smoking cigarettes.

‡Always or almost always took at least one of the following actions when students are caught smoking cigarettes: referred to a school counselor, encouraged to participate in an assistance, education, or cessation program, or required to participate in an assistance, education, or cessation program.

§Principal, assistant principal, other school administrator, or other school faculty or staff member.

¶A specified distance from school grounds where tobacco use is not allowed.

**Informed all groups (students, faculty and staff, and visitors) counts as 1 of the 7 criteria.

TABLE 33. Percentage of Secondary Schools That Provided Tobacco Cessation Services for Specific Groups and the Percentage That Had Arrangements with Organizations or Health Care Professionals Not on School Property to Provide Tobacco Cessation Services for Specific Groups, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Provided services		Had arrangements with organizations or health care professionals		Provided services or had arrangements for all groups
	Faculty and staff	Students	Faculty and staff	Students	
STATE SURVEYS					
Alabama	21.3	25.6	31.0	33.4	31.6
Alaska	13.1	24.2	29.0	39.7	28.3
Arizona	11.8	17.0	21.3	24.3	17.1
Arkansas	14.8	28.6	17.2	25.3	19.7
California	14.4	34.0	25.6	38.9	23.8
Colorado	11.0	28.7	27.7	36.8	24.4
Connecticut	14.0	26.2	22.7	32.6	18.8
Delaware	18.4	33.1	18.6	32.0	22.7
Florida	27.4	29.7	40.4	40.2	35.4
Georgia	13.7	13.0	20.4	21.6	18.7
Hawaii	5.6	37.2	12.9	50.7	15.9
Idaho	11.1	36.1	26.0	43.9	27.2
Indiana	15.2	31.8	31.0	48.2	28.0
Iowa	12.7	22.0	28.6	41.6	26.7
Kansas	11.6	13.5	18.6	19.6	16.3
Kentucky	14.4	37.1	25.5	45.0	24.5
Louisiana	12.1	11.5	14.6	19.8	15.7
Maine	20.4	30.8	37.8	44.0	41.3
Maryland	21.8	39.4	26.2	43.0	25.4
Massachusetts	18.0	24.2	31.9	34.7	30.3
Michigan	9.2	19.1	21.3	35.3	19.6
Minnesota	12.9	23.4	30.7	33.7	25.2
Mississippi	17.9	22.5	20.8	27.4	25.4
Missouri	14.5	16.7	22.8	23.6	20.1
Montana	20.1	28.9	20.3	33.5	25.7
Nebraska	9.2	12.9	16.6	19.6	16.5
Nevada	12.7	25.3	20.5	31.5	20.0
New Hampshire	16.6	19.2	44.2	33.9	28.5
New Jersey	16.2	34.5	25.5	36.6	26.6
New Mexico	13.7	33.9	26.6	42.5	29.8
New York	16.1	27.3	25.6	31.8	26.0
North Carolina	33.1	37.5	46.5	46.7	43.1
North Dakota	18.8	26.5	36.2	41.4	35.1
Ohio	15.7	23.1	32.4	49.5	30.2
Oklahoma	11.8	21.8	17.2	28.2	19.3
Oregon	19.4	28.4	31.3	46.5	29.6
Pennsylvania	19.8	34.2	35.2	48.3	32.6
Rhode Island	22.3	44.9	27.0	32.8	31.5
South Carolina	26.9	26.4	40.2	39.9	34.6
South Dakota	11.5	21.8	20.5	36.7	24.9
Tennessee	21.2	25.1	33.1	32.9	26.1
Texas	8.2	15.0	13.2	17.4	13.8
Utah	17.5	47.2	35.3	61.5	36.9
Vermont	22.7	48.6	43.1	42.6	38.8
Virginia	18.1	37.0	30.1	36.0	25.8
Washington	21.9	48.9	37.5	55.8	35.0
West Virginia	31.6	71.6	35.3	54.5	38.8
Wisconsin	18.3	26.0	35.3	31.0	27.3
Wyoming	24.6	40.6	34.9	53.8	36.0
State Median	16.1	27.3	27.0	36.6	26.1
State Range	5.6-33.1	11.5-71.6	12.9-46.5	17.4-61.5	13.8-43.1

TABLE 33. Percentage of Secondary Schools That Provided Tobacco Cessation Services for Specific Groups and the Percentage That Had Arrangements with Organizations or Health Care Professionals Not on School Property to Provide Tobacco Cessation Services for Specific Groups, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Provided services		Had arrangements with organizations or health care professionals		Provided services or had arrangements for all groups
	Faculty and staff	Students	Faculty and staff	Students	
LOCAL SURVEYS					
Albuquerque	13.6	30.7	24.3	24.5	21.7
Baltimore	13.5	17.1	23.3	31.9	19.8
Broward County	21.2	34.3	42.1	41.9	32.5
Charlotte-Mecklenberg County	21.4	49.8	28.7	47.9	30.7
Chicago	6.9	8.9	12.7	21.5	13.8
Detroit	15.9	17.0	24.3	23.1	25.7
District of Columbia	13.8	22.2	15.1	18.2	17.6
Fresno	31.9	28.6	23.6	32.4	24.2
Houston	19.4	19.3	22.1	22.0	23.3
Los Angeles	23.7	56.6	33.6	40.0	37.6
Memphis	24.1	41.1	41.1	53.8	43.0
Miami-Dade County	28.5	40.2	35.8	39.4	41.1
Milwaukee	13.4	23.6	22.4	21.0	17.9
New York City	9.6	20.5	14.8	23.5	15.8
Newark	23.1	47.1	36.4	72.7	39.2
Orange County	32.8	39.5	50.5	53.5	42.7
Philadelphia	9.4	11.1	13.3	20.8	13.3
San Diego	7.5	49.3	23.9	38.8	23.9
San Francisco	29.7	72.1	30.3	60.8	36.1
Local Median	19.4	30.7	24.3	32.4	24.2
Local Range	6.9-32.8	8.9-72.1	12.7-50.5	18.2-72.7	13.3-43.0
TERRITORIAL SURVEYS					
American Samoa	61.5	88.5	88.5	92.3	92.3
Guam	15.4	25.0	38.5	69.2	38.5
Marshall Islands	50.8	67.6	57.5	63.5	59.7
Northern Mariana Islands	42.9	62.5	42.9	50.0	57.1
Palau	9.1	9.1	63.6	72.7	72.7
Territorial Median	42.9	62.5	57.5	69.2	59.7
Territorial Range	9.1-61.5	9.1-88.5	38.5-88.5	50.0-92.3	38.5-92.3
TRIBAL SURVEYS					
Cherokee Nation	11.8	18.8	20.7	31.9	20.7
Nez Perce	0.0	42.9	28.6	42.9	28.6

TABLE 34. Percentage of Secondary Schools That Coordinated Tobacco-Use Prevention Messages and Programs with Community and Mass Media Efforts During the 2 Years Before the Survey, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Gathered and shared information with students and families about mass-media messages or community-based tobacco-use prevention efforts	Worked with local agencies or organizations to plan and implement events or programs intended to reduce tobacco use	Conducted both activities
STATE SURVEYS			
Alabama	55.4	49.5	33.9
Alaska	54.1	57.9	43.5
Arizona	47.9	49.9	35.4
Arkansas	57.3	60.9	45.9
California	46.0	52.6	37.2
Colorado	42.6	47.1	31.7
Connecticut	40.4	44.4	29.4
Delaware	58.3	56.8	47.4
Florida	54.5	60.2	45.6
Georgia	35.3	36.8	24.8
Hawaii	43.8	54.8	36.3
Idaho	50.4	51.3	39.5
Indiana	45.6	57.1	36.9
Iowa	47.7	61.3	40.6
Kansas	40.4	43.4	27.6
Kentucky	45.8	65.4	41.2
Louisiana	45.4	45.0	34.2
Maine	51.6	52.2	36.4
Maryland	44.1	47.9	33.8
Massachusetts	40.8	40.1	26.9
Michigan	33.3	35.5	23.6
Minnesota	41.9	45.2	29.3
Mississippi	58.7	59.9	47.5
Missouri	47.8	48.6	37.2
Montana	63.2	64.8	51.6
Nebraska	37.1	41.0	24.7
Nevada	42.2	46.7	31.4
New Hampshire	45.1	46.9	34.8
New Jersey	59.5	56.5	45.6
New Mexico	48.6	52.4	39.4
New York	52.4	48.9	38.3
North Carolina	50.1	53.6	41.3
North Dakota	47.9	62.3	39.0
Ohio	34.1	42.5	26.0
Oklahoma	49.8	58.6	42.1
Oregon	33.5	33.9	19.2
Pennsylvania	39.6	47.1	30.2
Rhode Island	37.4	42.8	28.0
South Carolina	41.9	42.5	29.0
South Dakota	49.3	58.0	36.5
Tennessee	48.3	52.0	37.7
Texas	44.5	48.2	34.5
Utah	58.8	58.7	44.2
Vermont	70.8	76.0	60.4
Virginia	44.6	44.0	30.0
Washington	45.0	54.1	36.5
West Virginia	74.4	82.9	68.4
Wisconsin	53.5	53.6	38.1
Wyoming	53.2	71.6	46.4
State Median	47.7	52.0	36.5
State Range	33.3-74.4	33.9-82.9	19.2-68.4

TABLE 34. Percentage of Secondary Schools That Coordinated Tobacco-Use Prevention Messages and Programs with Community and Mass Media Efforts During the 2 Years Before the Survey, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Gathered and shared information with students and families about mass-media messages or community-based tobacco-use prevention efforts	Worked with local agencies or organizations to plan and implement events or programs intended to reduce tobacco use	Conducted both activities
LOCAL SURVEYS			
Albuquerque	41.0	41.0	30.0
Baltimore	40.5	39.7	29.4
Broward County	56.0	53.7	45.1
Charlotte-Mecklenberg County	39.9	39.1	24.4
Chicago	34.8	27.0	21.8
Detroit	55.6	47.6	41.3
District of Columbia	34.3	40.0	28.6
Fresno	36.8	41.8	29.1
Houston	36.1	30.5	25.2
Los Angeles	63.1	62.0	45.6
Memphis	46.3	27.2	27.2
Miami-Dade County	60.5	56.5	46.0
Milwaukee	49.3	50.7	34.8
New York City	43.6	27.9	23.3
Newark	48.3	30.1	23.7
Orange County	56.7	39.5	37.3
Philadelphia	35.5	22.6	16.3
San Diego	29.9	32.8	16.4
San Francisco	81.6	69.2	66.3
Local Median	43.6	39.7	29.1
Local Range	29.9-81.6	22.6-69.2	16.3-66.3
TERRITORIAL SURVEYS			
American Samoa	69.2	96.2	69.2
Guam	33.3	69.2	16.7
Marshall Islands	64.2	59.6	47.8
Northern Mariana Islands	57.1	71.4	57.1
Palau	100.0	100.0	100.0
Territorial Median	64.2	71.4	57.1
Territorial Range	33.3-100.0	59.6-100.0	16.7-100.0
TRIBAL SURVEYS			
Cherokee Nation	54.1	62.2	44.2
Nez Perce	71.4	85.7	71.4

TABLE 35. Percentage of Secondary Schools with a Policy That Addressed Specific Issues on HIV* or AIDS,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Worksite safety†	Confidential counseling for HIV-infected students	Communication of the policy to students, school staff, and parents	Adequate training about HIV infection for school staff	Procedures for implementing the policy	Attendance of students with HIV infection	Procedures to protect HIV-infected students and staff from discrimination	Maintaining confidentiality of HIV-infected students and staff	All of the last 3 issues
STATE SURVEYS									
Alabama	77.7	64.0	63.1	58.5	57.1	65.3	68.5	74.4	51.4
Alaska	72.4	42.1	47.7	56.1	50.3	48.0	50.5	61.9	44.3
Arizona	80.2	52.9	56.5	62.5	59.2	58.4	65.7	72.2	56.9
Arkansas	73.9	59.8	55.9	44.2	50.7	51.5	59.0	70.5	50.9
California	79.6	57.3	57.0	57.9	56.1	53.6	63.6	73.5	51.2
Colorado	79.8	54.5	50.5	45.6	50.5	53.7	60.3	71.2	50.2
Connecticut	90.9	69.1	64.8	73.1	70.3	66.0	76.8	85.9	64.3
Delaware	72.7	53.2	43.8	37.1	37.9	38.3	50.6	72.9	36.7
Florida	83.7	66.4	62.9	67.8	63.3	58.0	71.0	83.3	56.5
Georgia	81.8	64.3	61.3	60.4	64.0	60.1	65.4	78.6	57.3
Hawaii	90.6	59.1	61.6	68.9	66.3	57.1	68.6	79.0	55.7
Idaho	78.2	64.8	61.1	51.5	60.4	62.0	66.1	73.3	60.8
Indiana	91.4	59.0	60.5	75.0	68.6	59.2	69.7	81.4	57.3
Iowa	83.0	58.1	58.6	67.1	64.0	56.4	69.0	75.8	53.6
Kansas	78.6	53.8	57.2	68.7	60.7	53.2	62.8	70.0	51.4
Kentucky	69.2	40.0	36.8	52.0	41.4	26.1	38.5	54.1	25.4
Louisiana	66.7	48.0	45.5	40.0	42.1	47.0	49.4	61.8	44.1
Maine	90.7	73.5	70.6	80.2	76.1	77.1	83.4	86.6	77.0
Maryland	82.3	63.8	61.3	64.5	65.4	56.2	66.3	77.7	54.1
Massachusetts	83.1	64.9	62.3	55.6	60.8	62.1	71.2	80.3	60.2
Michigan	82.1	61.3	51.9	66.4	58.6	49.8	63.5	74.2	48.5
Minnesota	80.5	56.5	64.1	67.7	64.0	57.1	67.0	74.6	56.4
Mississippi	58.4	50.3	39.3	30.9	34.9	38.6	47.8	56.5	36.7
Missouri	82.0	61.8	60.4	61.2	63.8	54.9	66.6	79.5	53.1
Montana	85.5	57.0	60.5	63.6	60.1	62.0	67.1	77.6	58.6
Nebraska	76.5	50.1	58.5	56.7	56.5	58.8	65.4	72.3	56.0
Nevada	92.4	71.2	72.3	76.2	73.7	69.4	77.5	87.8	68.7
New Hampshire	91.1	66.3	73.4	68.7	71.8	73.9	78.8	84.7	73.8
New Jersey	82.7	67.3	67.2	70.2	68.7	61.4	68.4	79.1	59.7
New Mexico	79.5	66.0	59.3	61.2	59.6	53.6	64.7	77.5	52.8
New York	87.1	71.4	67.0	76.1	71.8	60.2	73.3	84.4	59.2
North Carolina	86.0	64.2	58.7	62.4	56.7	54.2	65.5	76.2	51.1
North Dakota	72.6	46.5	53.1	56.1	51.6	49.3	54.7	62.8	46.7
Ohio	81.0	56.6	56.9	64.1	60.5	58.8	65.6	72.9	55.8
Oklahoma	91.2	70.8	76.7	90.6	79.2	69.9	78.3	85.9	68.9
Oregon	94.3	72.5	77.7	84.6	80.4	77.7	84.9	87.2	76.0
Pennsylvania	85.9	65.6	70.2	59.9	65.6	67.0	75.6	85.6	64.3
Rhode Island	90.5	72.9	73.1	77.6	73.5	75.7	84.7	88.3	70.9
South Carolina	91.5	73.5	68.7	79.1	74.7	69.2	75.7	86.8	67.6
South Dakota	67.1	52.9	59.5	52.3	55.9	59.2	61.5	68.2	56.3
Tennessee	91.2	75.0	74.7	80.8	77.4	72.3	79.3	85.8	71.8
Texas	77.6	56.0	55.6	56.4	55.2	51.1	59.4	75.8	47.8
Utah	84.1	62.1	60.2	60.9	64.0	65.3	68.8	80.1	61.7
Vermont	94.3	73.0	83.5	83.5	80.2	85.1	89.4	93.5	85.1
Virginia	87.9	67.1	66.5	75.8	73.2	68.9	78.7	86.2	67.5
Washington	91.3	68.8	69.1	81.6	72.9	66.0	79.5	87.5	64.4
West Virginia	76.4	63.7	58.4	55.2	56.4	54.4	63.5	73.6	53.0
Wisconsin	88.5	59.1	58.0	70.1	64.9	56.3	69.4	78.6	54.2
Wyoming	89.5	65.5	60.1	73.6	66.1	59.4	67.5	77.0	55.9
State Median	82.7	63.7	60.5	64.1	63.8	58.8	67.1	77.6	56.3
State Range	58.4-94.3	40.0-75.0	36.8-83.5	30.9-90.6	34.9-80.4	26.1-85.1	38.5-89.4	54.1-93.5	25.4-85.1

TABLE 35. Percentage of Secondary Schools with a Policy That Addressed Specific Issues on HIV* or AIDS,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Worksite safety*	Confidential counseling for HIV-infected students	Communication of the policy to students, school staff, and parents	Adequate training about HIV infection for school staff	Procedures for implementing the policy	Attendance of students with HIV infection	Procedures to protect HIV-infected students and staff from discrimination	Maintaining confidentiality of HIV-infected students and staff	All of the last 3 issues
LOCAL SURVEYS									
Albuquerque	89.0	70.0	53.8	73.1	62.1	65.0	67.6	86.4	65.0
Baltimore	62.3	42.1	46.9	44.6	44.4	37.1	42.9	60.0	33.3
Broward County	94.6	83.0	80.8	89.1	86.4	77.0	91.4	94.4	77.0
Charlotte-Mecklenberg County	84.0	71.6	58.7	62.5	52.8	53.0	64.9	84.0	53.0
Chicago	70.0	60.8	57.0	48.2	52.9	52.8	60.6	71.4	51.8
Detroit	63.8	52.8	50.6	36.5	41.9	32.9	42.3	59.4	31.7
District of Columbia	64.8	66.8	42.9	37.2	37.2	34.3	60.1	72.4	34.3
Fresno	82.7	61.9	53.6	61.9	61.9	51.4	61.9	74.4	51.4
Houston	77.8	68.9	64.2	37.0	53.1	36.0	70.3	80.1	36.0
Los Angeles	89.0	75.5	73.8	74.8	71.6	69.5	79.8	84.8	67.8
Memphis	86.6	75.3	79.4	75.9	75.6	60.5	79.3	83.9	60.5
Miami-Dade County	93.1	85.6	80.5	72.1	76.8	67.6	79.8	91.3	66.7
Milwaukee	75.7	54.4	51.3	44.8	54.4	47.3	58.9	71.4	44.4
New York City	85.4	74.4	71.2	79.0	72.4	53.6	69.0	82.8	53.1
Newark	77.4	56.7	53.2	47.3	42.8	35.3	52.2	73.0	31.4
Orange County	84.1	72.7	66.0	75.1	70.5	66.0	70.5	79.5	63.8
Philadelphia	74.1	63.8	63.4	47.6	60.1	54.4	68.1	76.2	53.7
San Diego	100.0	98.5	98.5	100.0	98.5	98.5	100.0	100.0	98.5
San Francisco	96.7	93.3	80.3	71.7	81.3	74.0	93.5	96.7	74.0
Local Median	84.0	70.0	63.4	62.5	61.9	53.6	68.1	80.1	53.1
Local Range	62.3-100.0	42.1-98.5	42.9-98.5	36.5-100.0	37.2-98.5	32.9-98.5	42.3-100.0	59.4-100.0	31.4-98.5
TERRITORIAL SURVEYS									
American Samoa	80.8	68.0	42.3	23.1	15.4	84.6	88.5	80.8	76.9
Guam	38.5	23.1	16.7	15.4	15.4	23.1	15.4	30.8	15.4
Marshall Islands	83.9	86.5	89.9	79.5	84.0	79.5	89.9	86.9	78.1
Northern Mariana Islands	87.5	85.7	87.5	62.5	87.5	75.0	75.0	75.0	62.5
Palau	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	83.9	85.7	87.5	62.5	84.0	79.5	88.5	80.8	76.9
Territorial Range	38.5-100.0	23.1-100.0	16.7-100.0	15.4-100.0	15.4-100.0	23.1-100.0	15.4-100.0	30.8-100.0	15.4-100.0
TRIBAL SURVEYS									
Cherokee Nation	93.5	77.2	86.9	94.2	89.5	80.1	86.1	90.7	79.1
Nez Perce	100.0	71.4	71.4	71.4	71.4	57.1	71.4	85.7	57.1

*Human immunodeficiency virus.

†Acquired immunodeficiency syndrome.

‡Universal precautions for all school staff.

TABLE 36. Percentage of Secondary Schools That Required School Staff to Receive Professional Development* on HIV,† STD,‡ or Pregnancy Prevention Issues and Resources for Specific Groups, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Youth who participate in drop-out prevention, alternative education, or GED§ programs	Ethnic/racial minority youth¶ at high risk
STATE SURVEYS		
Alabama	33.5	46.1
Alaska	28.5	32.9
Arizona	19.6	21.3
Arkansas	42.1	29.8
California	24.4	32.0
Colorado	25.1	20.3
Connecticut	18.3	18.9
Delaware	33.1	24.6
Florida	37.0	39.8
Georgia	39.3	37.0
Hawaii	27.1	27.9
Idaho	24.4	20.9
Indiana	23.1	13.3
Iowa	32.0	21.8
Kansas	15.1	12.1
Kentucky	28.5	22.0
Louisiana	33.3	26.7
Maine	21.0	8.7
Maryland	38.4	42.5
Massachusetts	22.9	19.9
Michigan	22.1	26.6
Minnesota	21.7	19.0
Mississippi	47.6	35.5
Missouri	32.2	19.0
Montana	25.0	23.6
Nebraska	21.7	14.6
Nevada	30.7	28.7
New Hampshire	13.0	2.7
New Jersey	20.1	24.5
New Mexico	34.8	40.3
New York	29.4	33.0
North Carolina	39.7	37.9
North Dakota	14.9	13.8
Ohio	17.7	19.5
Oklahoma	51.6	50.6
Oregon	21.6	21.8
Pennsylvania	24.2	20.6
Rhode Island	24.5	18.0
South Carolina	47.3	49.4
South Dakota	10.6	6.2
Tennessee	33.5	34.4
Texas	44.8	44.6
Utah	42.7	43.7
Vermont	16.3	12.8
Virginia	33.1	26.8
Washington	33.2	33.3
West Virginia	41.3	26.7
Wisconsin	27.3	22.6
Wyoming	22.3	20.7
State Median	27.3	24.5
State Range	10.6-51.6	2.7-50.6

TABLE 36. Percentage of Secondary Schools That Required School Staff to Receive Professional Development* on HIV,† STD,‡ or Pregnancy Prevention Issues and Resources for Specific Groups, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Youth who participate in drop-out prevention, alternative education, or GED§ programs	Ethnic/racial minority youth¶ at high risk
LOCAL SURVEYS		
Albuquerque	33.1	44.1
Baltimore	36.8	41.5
Broward County	48.2	57.5
Charlotte-Mecklenberg County	44.1	40.5
Chicago	24.4	42.9
Detroit	39.0	60.0
District of Columbia	37.3	64.0
Fresno	20.9	20.3
Houston	43.3	42.0
Los Angeles	54.0	73.3
Memphis	42.8	63.6
Miami-Dade County	62.1	72.5
Milwaukee	29.5	50.9
New York City	44.2	65.3
Newark	53.3	64.0
Orange County	45.9	58.6
Philadelphia	25.0	37.1
San Diego	0.0	100.0
San Francisco	63.7	70.0
Local Median	42.8	58.6
Local Range	0.0-63.7	20.3-100.0
TERRITORIAL SURVEYS		
American Samoa	23.1	57.7
Guam	46.2	23.1
Marshall Islands	55.3	36.4
Northern Mariana Islands	42.9	50.0
Palau	81.8	72.7
Territorial Median	46.2	50.0
Territorial Range	23.1-81.8	23.1-72.7
TRIBAL SURVEYS		
Cherokee Nation	55.1	55.0
Nez Perce	42.9	57.1

*Workshops, conferences, continuing education, or any other kind of in-service.

†Human immunodeficiency virus.

‡Sexually transmitted disease.

§General Equivalency Diploma.

¶Such as black, Hispanic, or American Indian youth.

TABLE 37. Percentage of Secondary Schools That Provided HIV,* STD,† or Pregnancy Prevention Programs for Ethnic/Racial Minority Youth‡ at High Risk That Did Each Specific Activity, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010

Site	Provided curricula or materials that reflect life experiences of these youth	Provided curricula or materials in the primary languages of these youth and families	Facilitated access to health services [§]	Facilitated access to social and psychological services [§]	Required professional development for school staff on issues and resources for these youth and did all 4 activities
STATE SURVEYS					
Alabama	52.7	37.5	41.2	38.1	12.0
Alaska	24.0	20.5	27.9	27.2	4.0
Arizona	21.4	14.9	23.5	25.6	3.4
Arkansas	33.5	30.4	33.7	33.0	6.9
California	39.3	33.5	40.0	39.4	12.5
Colorado	NA	NA	NA	NA	3.8
Connecticut	23.8	19.0	22.1	21.0	2.4
Delaware	33.8	30.0	46.7	46.7	5.5
Florida	32.2	27.2	31.1	29.7	9.8
Georgia	30.8	28.2	30.2	26.7	6.4
Hawaii	43.8	27.0	46.3	41.5	5.1
Idaho	20.8	18.2	26.1	24.1	2.9
Indiana	19.8	13.4	19.1	19.6	1.8
Iowa	22.4	19.1	23.3	24.3	4.9
Kansas	18.5	12.9	15.6	17.9	1.6
Kentucky	32.2	25.5	32.7	32.2	6.2
Louisiana	23.4	19.3	19.4	20.0	4.7
Maine	13.8	12.2	13.4	13.7	0.7
Maryland	40.2	32.9	34.0	32.0	7.5
Massachusetts	25.0	16.2	25.9	27.4	4.7
Michigan	23.7	20.2	19.2	22.2	4.7
Minnesota	19.6	15.2	22.5	22.6	2.7
Mississippi	37.2	31.4	28.7	32.1	12.4
Missouri	26.0	23.8	25.0	26.2	3.7
Montana	22.4	22.6	23.5	25.2	3.8
Nebraska	26.7	20.1	18.0	18.0	3.2
Nevada	43.7	40.5	41.6	36.9	6.9
New Hampshire	9.4	5.1	8.9	9.9	0.5
New Jersey	31.0	24.5	31.8	33.3	3.8
New Mexico	NA	NA	NA	NA	12.3
New York	37.0	31.1	35.1	36.3	10.9
North Carolina	30.3	26.6	30.3	30.1	8.2
North Dakota	19.2	18.4	20.4	20.2	4.5
Ohio	27.2	21.2	23.9	24.3	3.4
Oklahoma	36.6	30.2	35.5	38.5	15.3
Oregon	24.1	21.6	26.6	25.8	3.3
Pennsylvania	24.3	18.8	25.8	25.0	2.5
Rhode Island	18.7	16.6	21.9	22.9	4.9
South Carolina	34.1	29.2	28.1	27.1	8.0
South Dakota	20.2	16.5	13.7	15.2	1.4
Tennessee	32.1	28.6	34.5	33.7	11.0
Texas	31.6	27.1	28.3	27.2	9.6
Utah	16.3	16.8	16.4	13.4	1.5
Vermont	10.8	6.1	12.9	15.7	0.0
Virginia	24.6	21.4	23.1	23.4	2.8
Washington	31.9	28.2	36.4	32.8	6.3
West Virginia	29.4	27.6	24.7	24.7	6.6
Wisconsin	30.2	27.0	28.6	28.6	5.3
Wyoming	12.8	11.8	19.0	15.7	0.7
State Median	26.0	21.6	25.9	25.8	4.7
State Range	9.4-52.7	5.1-40.5	8.9-46.7	9.9-46.7	0.0-15.3

TABLE 37. Percentage of Secondary Schools That Provided HIV,* STD,† or Pregnancy Prevention Programs for Ethnic/Racial Minority Youth‡ at High Risk That Did Each Specific Activity, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010 (continued)

Site	Provided curricula or materials that reflect life experiences of these youth	Provided curricula or materials in the primary languages of these youth and families	Facilitated access to health services§	Facilitated access to social and psychological services§	Required professional development for school staff on issues and resources for these youth and did all 4 activities
LOCAL SURVEYS					
Albuquerque	32.2	35.3	48.2	58.0	17.1
Baltimore	45.5	33.3	34.6	32.9	11.4
Broward County	39.4	36.0	39.8	45.7	13.3
Charlotte-Mecklenberg County	39.9	35.8	40.5	38.5	0.0
Chicago	35.9	32.9	40.0	37.9	16.3
Detroit	41.0	33.9	31.3	34.9	21.1
District of Columbia	91.1	81.8	65.6	65.8	38.2
Fresno	35.2	26.0	29.9	31.4	7.7
Houston	68.0	58.8	56.6	49.2	19.7
Los Angeles	71.3	53.1	63.4	65.8	33.3
Memphis	46.6	46.9	38.5	38.9	23.1
Miami-Dade County	51.7	48.8	48.6	50.7	37.3
Milwaukee	59.6	54.7	58.5	53.3	21.8
New York City	56.8	48.6	49.3	50.6	24.5
Newark	50.2	40.5	48.8	44.0	8.7
Orange County	57.9	48.7	51.1	53.5	21.5
Philadelphia	45.8	38.7	46.3	42.0	12.8
San Diego	100.0	98.3	100.0	100.0	98.3
San Francisco	81.6	63.2	81.8	81.9	47.0
Local Median	50.2	46.9	48.6	49.2	21.1
Local Range	32.2-100.0	26.0-98.3	29.9-100.0	31.4-100.0	0.0-98.3
TERRITORIAL SURVEYS					
American Samoa	65.4	38.5	53.8	42.3	7.4
Guam	30.8	0.0	30.8	30.8	0.0
Marshall Islands	NA	NA	NA	NA	6.0
Northern Mariana Islands	75.0	50.0	62.5	62.5	25.0
Palau	100.0	90.9	81.8	90.9	63.6
Territorial Median	70.2	44.3	58.2	52.4	7.4
Territorial Range	30.8-100.0	0.0-90.9	30.8-81.8	30.8-90.9	0.0-63.6
TRIBAL SURVEYS					
Cherokee Nation	39.1	30.7	45.5	45.8	17.5
Nez Perce	28.6	14.3	14.3	14.3	14.3

NA=Data not available.

*Human immunodeficiency virus.

†Sexually transmitted disease.

‡Such as black, Hispanic, or American Indian youth.

§Direct services or arrangements with providers not on school property who have experience in serving these youth in the community.

TABLE 38. Percentage of Secondary Schools that Provide Curricula or Supplementary Materials* that Include HIV,† STD,‡ or Pregnancy Prevention Information Relevant to Lesbian, Gay, Bisexual, Transgender, or Questioning (LGBTQ) Youth; the Percentage that Engage in the Following Practices Related to LGBTQ Youth; and the Percentage That Had a Gay/Straight Alliance or Similar Club,§ Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010

Site	Practices related to LGBTQ Youth						Schools that provide curricula or supplementary materials and engage in all 5 practices related to LGBTQ youth	Had a gay/straight alliance or similar club
	Provide curricula or supplementary materials	Identify safe spaces ¹	Prohibit harassment ^{**}	Encourage staff to attend professional development on safe and supportive school environments for all students ^{††}	Facilitate access to providers not on school property who have experience in providing health services ^{††} to LGBTQ youth	Facilitate access to providers not on school property who have experience in providing social and psychological services to LGBTQ youth		
STATE SURVEYS								
Alabama	39.4	32.6	77.2	38.4	34.6	30.0	2.1	20.5
Alaska	18.4	34.8	71.9	48.7	40.1	39.8	4.3	20.3
Arizona	10.5	48.0	82.1	44.6	37.8	39.9	2.0	24.1
Arkansas	23.3	41.4	84.0	49.7	38.1	40.9	7.8	15.0
California	33.7	57.5	89.8	60.1	53.9	55.7	13.3	38.6
Colorado	NA	56.8	85.4	50.9	42.4	48.5	5.1	40.1
Connecticut	28.9	62.1	90.5	66.7	52.0	58.8	11.0	42.7
Delaware	45.6	39.8	79.2	48.1	44.8	42.1	8.7	27.8
Florida	17.5	57.4	89.9	62.6	48.1	49.1	5.7	33.5
Georgia	7.1	45.5	84.5	44.2	36.6	34.5	0.7	15.2
Hawaii	50.8	69.6	98.9	66.2	62.5	59.5	14.1	28.1
Idaho	10.7	48.2	88.0	51.1	42.5	42.2	1.9	22.8
Indiana	8.7	52.3	90.4	40.2	38.1	42.1	2.5	26.0
Iowa	23.2	61.4	95.2	64.2	50.0	50.6	8.9	22.2
Kansas	11.7	44.4	83.2	44.4	32.5	33.6	0.7	18.5
Kentucky	15.7	48.0	88.5	48.8	43.8	44.5	3.0	16.6
Louisiana	11.2	40.5	80.5	50.6	36.4	40.1	2.8	9.3
Maine	33.1	61.7	93.1	73.0	61.3	58.0	13.4	46.4
Maryland	30.8	54.1	91.2	66.8	47.1	50.7	6.8	35.8
Massachusetts	40.8	71.5	93.0	75.2	58.5	65.3	18.0	50.1
Michigan	21.6	48.6	84.1	49.4	44.4	46.2	5.9	27.2
Minnesota	20.5	57.6	93.0	65.0	54.1	55.7	8.2	27.1
Mississippi	19.0	42.2	77.0	49.8	37.2	41.4	2.4	15.2
Missouri	16.1	41.9	81.6	49.1	40.3	41.3	3.2	19.1
Montana	9.8	42.1	86.2	56.5	37.8	41.3	2.2	23.4
Nebraska	17.2	39.6	79.7	50.1	34.0	38.3	5.8	16.6
Nevada	30.4	55.2	87.4	53.9	38.6	39.8	6.2	31.3
New Hampshire	31.4	63.2	89.8	66.2	59.5	60.2	12.6	36.7
New Jersey	34.5	65.1	91.1	69.5	57.4	56.0	13.0	37.7
New Mexico	NA	48.3	88.3	54.4	63.7	61.6	5.3	23.0
New York	42.8	66.0	90.1	73.0	63.2	64.1	21.4	41.9
North Carolina	12.0	51.6	85.8	52.8	40.2	40.2	2.9	19.3
North Dakota	15.3	29.6	74.1	46.2	30.5	31.1	3.0	15.8
Ohio	22.7	45.6	86.1	46.8	38.5	40.7	5.3	21.5
Oklahoma	22.3	40.8	77.7	51.4	39.9	39.6	4.9	11.4
Oregon	25.6	58.9	90.5	61.1	51.0	56.7	8.2	26.8
Pennsylvania	21.4	61.8	93.4	66.0	54.4	57.6	9.8	29.8
Rhode Island	31.6	66.2	93.3	68.3	65.3	65.6	15.3	41.2
South Carolina	11.5	42.6	84.0	54.1	36.9	37.6	1.1	15.2
South Dakota	16.0	28.4	72.4	40.6	28.4	33.6	1.9	11.6
Tennessee	20.1	56.2	89.0	64.4	48.3	49.9	5.7	18.5
Texas	16.3	41.1	78.6	47.5	35.0	36.2	4.4	20.1
Utah	6.1	53.7	90.6	56.3	38.8	42.6	0.9	27.0
Vermont	42.7	66.8	96.4	80.2	54.2	62.1	19.1	32.4
Virginia	13.2	53.8	89.4	60.0	38.8	42.4	3.4	25.8
Washington	27.7	61.9	92.4	67.3	58.2	54.2	7.1	43.0
West Virginia	21.7	50.1	91.2	65.2	46.4	50.7	6.6	24.0
Wisconsin	26.0	63.9	90.8	70.9	52.6	57.0	9.9	28.0
Wyoming	13.4	40.1	86.1	49.9	46.8	46.9	3.2	25.5
State Median	21.4	51.6	88.3	54.1	43.8	44.5	5.7	25.5
State Range	6.1-50.8	28.4-71.5	71.9-98.9	38.4-80.2	28.4-65.3	30.0-65.6	0.7-21.4	9.3-50.1

TABLE 38. Percentage of Secondary Schools that Provide Curricula or Supplementary Materials* that Include HIV,[†] STD,[‡] or Pregnancy Prevention Information Relevant to Lesbian, Gay, Bisexual, Transgender, or Questioning (LGBTQ) Youth; the Percentage that Engage in the Following Practices Related to LGBTQ Youth; and the Percentage That Had a Gay/Straight Alliance or Similar Club,[§] Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2010 (*continued*)

Site	Practices related to LGBTQ Youth						Schools that provide curricula or supplementary materials and engage in all 5 practices related to LGBTQ youth	Had a gay/straight alliance or similar club
	Provide curricula or supplementary materials	Identify safe spaces [†]	Prohibit harassment ^{**}	Encourage staff to attend professional development on safe and supportive school environments for all students ^{††}	Facilitate access to providers not on school property who have experience in providing health services ^{††} to LGBTQ youth	Facilitate access to providers not on school property who have experience in providing social and psychological services to LGBTQ youth		
LOCAL SURVEYS								
Albuquerque	46.3	92.2	97.4	73.3	76.5	73.6	27.7	60.3
Baltimore	17.0	40.7	76.2	56.5	40.0	41.2	4.1	16.4
Broward County	30.1	76.1	93.2	79.9	51.6	54.9	14.2	42.3
Charlotte-Mecklenberg County	15.7	57.3	83.7	53.6	35.8	42.3	3.5	30.5
Chicago	35.4	51.7	75.1	61.6	45.9	40.8	12.6	28.3
Detroit	28.7	44.5	74.3	57.9	36.7	40.5	11.2	30.9
District of Columbia	66.9	58.4	86.2	69.6	54.4	52.9	17.7	27.7
Fresno	26.0	68.0	88.0	80.0	55.4	42.9	12.5	34.1
Houston	37.3	31.5	47.7	35.4	29.0	28.9	8.7	22.3
Los Angeles	69.0	83.5	97.4	86.0	79.0	76.4	48.1	62.0
Memphis	31.2	61.9	81.8	69.7	41.6	45.2	5.8	33.9
Miami-Dade County	40.1	82.1	95.2	83.0	70.5	68.7	25.8	53.6
Milwaukee	39.0	48.3	70.0	51.3	47.9	51.0	13.5	18.6
New York City	49.3	72.2	88.4	79.9	69.3	65.4	32.6	45.4
Newark	32.3	56.4	70.9	56.4	56.7	56.7	6.9	43.3
Orange County	62.7	80.3	95.6	77.8	75.2	70.6	26.2	43.0
Philadelphia	30.8	73.3	90.6	75.9	64.9	61.7	17.2	33.5
San Diego	100.0	94.0	100.0	82.1	95.5	94.0	75.4	31.3
San Francisco	83.7	100.0	100.0	100.0	100.0	100.0	81.6	90.8
Local Median	37.3	68.0	88.0	73.3	55.4	54.9	14.2	33.9
Local Range	15.7-100.0	31.5-100.0	47.7-100.0	35.4-100.0	29.0-100.0	28.9-100.0	3.5-81.6	16.4-90.8
TERRITORIAL SURVEYS								
American Samoa	54.2	69.2	76.9	76.9	57.7	61.5	28.0	46.2
Guam	44.4	61.5	76.9	38.5	30.8	30.8	7.7	41.7
Marshall Islands	NA	27.9	29.8	47.0	24.9	19.1	5.6	26.7
Northern Mariana Islands	50.0	75.0	87.5	87.5	75.0	50.0	12.5	50.0
Palau	18.2	18.2	9.1	18.2	9.1	9.1	0.0	81.8
Territorial Median	47.2	61.5	76.9	47.0	30.8	30.8	7.7	46.2
Territorial Range	18.2-54.2	18.2-75.0	9.1-87.5	18.2-87.5	9.1-75.0	9.1-61.5	0.0-28.0	26.7-81.8
TRIBAL SURVEYS								
Cherokee Nation	19.2	44.2	78.0	56.2	44.7	46.1	4.0	15.1
Nez Perce	16.7	33.3	100.0	100.0	71.4	71.4	16.7	0.0

NA=Data not available.

*Such as curricula or materials that use inclusive language or terminology.

†Human immunodeficiency virus.

‡Sexually transmitted disease.

§A student-led club that aims to create a safe, welcoming, and accepting school environment for all youth, regardless of sexual orientation or gender identity.

¶Such as a counselor's office, designated classroom, or student organization where LGBTQ youth can receive support from administrators, teachers, or other school staff.

**Based on a student's perceived or actual sexual orientation or gender identity.

††Regardless of sexual orientation or gender identity.

‡‡Including HIV/STD testing and counseling.

TABLE 39. Percentage of Secondary Schools That Have or Participate in the Following Programs, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Student mentoring program	Safe passages to school program	Program to prevent bullying	Program to prevent dating violence	Youth development program
STATE SURVEYS					
Alabama	63.3	20.9	72.7	27.9	39.6
Alaska	41.4	13.1	69.7	26.3	47.7
Arizona	59.7	27.2	79.4	28.3	46.0
Arkansas	57.0	23.4	80.6	31.5	39.7
California	67.1	25.8	74.0	26.0	48.0
Colorado	78.4	29.9	85.5	32.2	58.5
Connecticut	70.9	12.7	89.3	30.1	52.8
Delaware	84.2	16.5	90.7	26.5	51.6
Florida	80.2	25.6	91.9	31.6	49.6
Georgia	83.2	19.2	79.0	23.0	50.2
Hawaii	53.2	19.2	75.3	38.0	59.7
Idaho	63.4	26.1	81.6	61.5	52.1
Indiana	69.3	18.6	78.6	33.1	54.3
Iowa	58.6	18.0	80.7	29.2	54.9
Kansas	60.9	14.7	95.9	26.4	40.6
Kentucky	73.6	25.8	83.8	29.3	52.0
Louisiana	58.1	23.2	75.8	27.3	36.3
Maine	56.0	13.5	80.8	37.0	32.1
Maryland	85.4	64.2	87.8	61.7	76.7
Massachusetts	65.7	20.6	83.6	43.0	46.4
Michigan	63.4	24.5	73.9	27.8	46.4
Minnesota	68.3	22.0	83.8	33.6	60.3
Mississippi	61.3	30.9	69.0	27.7	48.6
Missouri	69.0	19.3	81.5	35.6	46.3
Montana	61.3	26.0	79.9	36.8	52.6
Nebraska	67.9	20.6	83.6	51.6	55.1
Nevada	71.5	27.9	77.5	20.2	44.4
New Hampshire	54.7	18.7	85.2	37.4	46.8
New Jersey	74.3	27.3	94.5	44.9	60.9
New Mexico	59.0	27.1	80.0	48.2	50.2
New York	72.7	28.3	87.2	45.3	68.2
North Carolina	73.8	20.7	82.9	27.3	47.9
North Dakota	41.5	15.4	72.0	21.8	36.8
Ohio	60.4	21.4	77.4	36.8	49.8
Oklahoma	46.9	21.9	83.3	32.9	44.4
Oregon	59.2	19.7	70.8	23.1	41.0
Pennsylvania	72.0	21.2	90.3	30.9	49.4
Rhode Island	71.1	20.3	81.9	77.5	49.7
South Carolina	78.5	25.8	80.5	28.0	53.0
South Dakota	50.5	16.7	81.2	21.2	35.9
Tennessee	67.5	28.6	88.5	34.7	59.9
Texas	71.4	26.7	83.1	64.3	51.5
Utah	80.8	56.2	90.4	40.9	61.9
Vermont	66.1	20.2	92.6	34.8	54.2
Virginia	76.9	28.1	83.2	34.9	51.4
Washington	62.6	16.2	73.4	24.4	38.8
West Virginia	72.5	31.1	89.2	40.6	50.6
Wisconsin	67.5	20.5	82.9	41.0	53.5
Wyoming	48.5	18.6	84.7	25.1	38.2
State Median	67.1	21.4	81.9	32.2	49.8
State Range	41.4-85.4	12.7-64.2	69.0-95.9	20.2-77.5	32.1-76.7

TABLE 39. Percentage of Secondary Schools That Have or Participate in the Following Programs, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Student mentoring program	Safe passages to school program	Program to prevent bullying	Program to prevent dating violence	Youth development program
LOCAL SURVEYS					
Albuquerque	67.9	29.8	81.2	41.2	43.8
Baltimore	78.5	30.5	74.5	24.8	65.0
Broward County	95.5	38.2	98.3	47.6	65.4
Charlotte-Mecklenberg County	84.0	15.6	94.2	23.5	42.8
Chicago	75.5	68.0	76.0	29.1	60.3
Detroit	81.3	62.3	74.5	25.5	66.7
District of Columbia	75.0	43.0	65.6	41.8	79.9
Fresno	80.2	40.1	84.6	29.1	56.0
Houston	86.4	37.8	54.1	41.5	67.5
Los Angeles	77.8	65.1	86.7	51.1	78.9
Memphis	93.0	43.3	89.6	33.9	86.6
Miami-Dade County	84.9	47.6	97.2	47.6	74.9
Milwaukee	71.6	25.1	78.8	25.5	49.7
New York City	75.7	40.7	89.6	51.6	80.1
Newark	81.6	57.8	91.3	36.0	67.7
Orange County	93.6	44.7	97.9	39.7	67.2
Philadelphia	83.3	37.1	86.1	26.9	52.5
San Diego	59.7	32.8	74.6	88.1	44.8
San Francisco	88.4	46.7	88.4	62.2	78.0
Local Median	81.3	40.7	86.1	39.7	66.7
Local Range	59.7-95.5	15.6-68.0	54.1-98.3	23.5-88.1	42.8-86.6
TERRITORIAL SURVEYS					
American Samoa	57.7	19.2	92.3	73.1	46.2
Guam	69.2	30.8	84.6	53.8	84.6
Marshall Islands	63.6	70.1	71.2	53.0	58.2
Northern Mariana Islands	87.5	62.5	100.0	62.5	100.0
Palau	100.0	45.5	90.9	54.5	100.0
Territorial Median	69.2	45.5	90.9	54.5	84.6
Territorial Range	57.7-100.0	19.2-70.1	71.2-100.0	53.0-73.1	46.2-100.0
TRIBAL SURVEYS					
Cherokee Nation	48.7	32.4	80.4	30.8	49.3
Nez Perce	42.9	28.6	71.4	57.1	42.9

TABLE 40. Percentage of Secondary Schools That Had a Full-Time* Registered Nurse Who Provided Health Services to Students; the Percentage That Had an Asthma Action Plan on File for All Students with Known Asthma;† the Percentage That Had a Designated, Secure, and Accessible Storage Location for Medications; and the Percentage in Which School Staff Members Were Required to Receive Training on Recognizing and Responding to Severe Asthma Symptoms at Least Once Per Year, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Full-time registered nurse	Asthma action plan for all students with known asthma	Annual training for school staff members
STATE SURVEYS			
Alabama	70.4	87.6	48.0
Alaska	19.9	31.9	15.9
Arizona	41.3	44.7	26.0
Arkansas	71.6	66.0	36.1
California	21.8	53.5	24.5
Colorado	24.5	67.6	35.7
Connecticut	88.9	63.9	30.8
Delaware	98.5	70.3	26.6
Florida	43.9	65.1	36.2
Georgia	55.9	59.8	34.6
Hawaii	17.8	44.8	23.4
Idaho	16.2	49.0	19.2
Indiana	63.5	62.9	25.2
Iowa	47.7	51.2	23.8
Kansas	38.5	52.9	22.1
Kentucky	56.1	63.6	48.6
Louisiana	25.6	72.5	40.9
Maine	46.5	59.4	27.5
Maryland	68.6	69.9	32.6
Massachusetts	96.1	61.7	33.3
Michigan	12.8	31.0	22.7
Minnesota	41.4	56.9	26.6
Mississippi	47.9	46.7	29.8
Missouri	75.8	63.9	40.6
Montana	11.9	40.9	13.7
Nebraska	32.9	67.0	65.8
Nevada	19.3	68.2	32.9
New Hampshire	94.4	58.5	25.5
New Jersey	99.4	75.3	61.2
New Mexico	56.1	53.2	36.9
New York	94.8	54.1	28.7
North Carolina	24.2	72.5	55.8
North Dakota	6.3	32.2	11.6
Ohio	39.3	50.4	23.5
Oklahoma	26.6	56.0	37.0
Oregon	8.1	53.1	28.9
Pennsylvania	84.6	55.2	26.8
Rhode Island	87.4	70.8	37.6
South Carolina	82.5	76.6	46.2
South Dakota	23.5	34.3	17.1
Tennessee	49.1	69.8	42.0
Texas	77.3	56.8	40.7
Utah	4.9	49.8	28.5
Vermont	76.3	49.1	26.1
Virginia	74.2	66.3	43.6
Washington	20.1	82.3	62.0
West Virginia	27.2	77.2	41.0
Wisconsin	22.5	53.2	25.9
Wyoming	37.5	44.8	32.2
State Median	43.9	58.5	30.8
State Range	4.9-99.4	31.0-87.6	11.6-65.8

TABLE 40. Percentage of Secondary Schools That Had a Full-Time* Registered Nurse Who Provided Health Services to Students; the Percentage That Had an Asthma Action Plan on File for All Students with Known Asthma;† the Percentage That Had a Designated, Secure, and Accessible Storage Location for Medications; and the Percentage in Which School Staff Members Were Required to Receive Training on Recognizing and Responding to Severe Asthma Symptoms at Least Once Per Year, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Full-time registered nurse	Asthma action plan for all students with known asthma	Annual training for school staff members
LOCAL SURVEYS			
Albuquerque	89.3	72.6	79.1
Baltimore	68.8	62.7	27.2
Broward County	43.3	56.9	51.3
Charlotte-Mecklenberg County	46.9	72.4	94.3
Chicago	16.9	55.8	38.6
Detroit	31.4	42.1	31.8
District of Columbia	100.0	45.7	28.6
Fresno	30.2	55.5	36.3
Houston	91.0	45.5	26.9
Los Angeles	92.0	63.7	39.1
Memphis	14.7	57.7	38.6
Miami-Dade County	42.4	35.6	23.6
Milwaukee	31.2	56.8	37.0
New York City	88.4	52.2	35.7
Newark	97.9	87.0	62.4
Orange County	48.6	63.0	26.3
Philadelphia	74.6	69.2	26.9
San Diego	58.2	31.3	0.0
San Francisco	33.9	58.1	31.2
Local Median	48.6	56.9	35.7
Local Range	14.7-100.0	31.3-87.0	0.0-94.3
TERRITORIAL SURVEYS			
American Samoa	0.0	0.0	46.2
Guam	91.7	30.8	7.7
Marshall Islands	29.3	4.9	30.4
Northern Mariana Islands	12.5	12.5	50.0
Palau	0.0	0.0	0.0
Territorial Median	12.5	4.9	30.4
Territorial Range	0.0-91.7	0.0-30.8	0.0-50.0
TRIBAL SURVEYS			
Cherokee Nation	44.6	61.4	37.2
Nez Perce	0.0	71.4	28.6

*A nurse is at the school during all school hours, 5 days per week.

†Those identified by the school to have a current diagnosis of asthma as reported on student emergency cards, medication records, health room visit information, emergency care plans, physical exam forms, parent notes, and other forms of health care clinician notification.

TABLE 41. Percentage of Secondary Schools That Used Specific Types of Information to Identify Students with Poorly Controlled Asthma, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Frequent absences from school	Frequent visits to the school health office due to asthma	Frequent asthma symptoms at school	Frequent non-participation in physical education due to asthma	Students sent home early due to asthma	Calls from school to 911* due to asthma	At least 3 of these 6 types of information
STATE SURVEYS							
Alabama	30.9	58.5	48.0	38.5	34.0	18.3	48.5
Alaska	19.7	26.0	30.4	22.7	20.5	10.4	50.5
Arizona	39.7	61.0	56.5	39.3	39.8	26.2	63.4
Arkansas	26.0	63.7	54.6	27.2	31.8	17.3	47.6
California	44.3	67.2	61.3	54.5	51.7	37.5	72.3
Colorado	36.9	65.6	59.5	46.6	44.6	29.5	64.2
Connecticut	47.2	78.9	67.8	54.3	54.0	34.5	71.5
Delaware	46.2	78.4	67.7	51.3	55.0	31.7	66.8
Florida	34.7	62.8	51.6	37.9	43.6	31.7	62.1
Georgia	40.3	71.2	67.2	47.1	46.9	32.3	71.3
Hawaii	54.7	75.8	60.9	49.4	50.2	33.1	69.0
Idaho	36.0	33.2	46.9	35.8	32.0	20.4	58.0
Indiana	46.2	71.9	60.2	45.7	43.8	25.5	67.0
Iowa	35.0	64.9	50.1	38.7	31.6	15.6	57.9
Kansas	30.2	51.8	51.1	41.0	31.8	18.6	59.3
Kentucky	41.0	56.3	49.4	30.2	33.4	18.4	54.1
Louisiana	33.3	45.4	56.6	32.7	33.3	16.8	53.0
Maine	43.8	67.2	62.9	45.7	46.1	23.5	72.2
Maryland	52.7	78.6	62.1	44.9	58.3	44.0	74.1
Massachusetts	50.2	80.4	66.2	49.2	55.1	30.6	70.7
Michigan	35.9	26.2	50.2	34.7	33.6	17.9	59.2
Minnesota	44.9	69.1	56.9	46.2	39.6	25.6	62.8
Mississippi	32.1	50.3	58.3	33.4	34.6	19.8	51.2
Missouri	46.4	74.6	66.2	48.1	48.7	30.2	67.0
Montana	29.7	32.7	45.0	38.3	28.5	12.0	54.9
Nebraska	32.4	52.8	57.7	46.0	36.8	23.0	59.1
Nevada	36.7	69.3	55.6	49.7	50.0	37.2	65.9
New Hampshire	52.7	82.1	67.2	50.1	52.0	27.9	68.8
New Jersey	54.2	82.1	67.0	52.9	54.5	30.7	73.2
New Mexico	40.0	71.0	66.8	50.8	49.7	31.2	72.5
New York	54.7	76.8	63.4	52.2	52.5	33.2	72.8
North Carolina	45.5	64.8	64.0	47.8	51.0	38.4	70.6
North Dakota	18.1	17.9	37.7	29.5	14.3	9.4	44.7
Ohio	32.1	55.9	51.8	34.6	35.4	22.6	62.5
Oklahoma	30.2	33.0	49.7	30.9	28.2	10.4	48.1
Oregon	34.7	50.3	56.7	44.8	38.8	24.7	65.9
Pennsylvania	44.1	76.5	64.4	49.0	51.5	31.9	71.9
Rhode Island	52.3	75.1	62.3	50.9	56.6	29.5	75.9
South Carolina	41.3	81.4	62.0	42.0	50.9	28.7	63.9
South Dakota	20.9	25.6	39.8	27.6	20.8	7.2	40.7
Tennessee	39.0	65.6	58.8	35.9	42.3	26.3	54.9
Texas	40.3	74.2	59.0	44.7	44.6	25.8	61.0
Utah	36.7	34.6	58.5	46.8	34.4	24.8	60.3
Vermont	59.4	84.8	75.9	60.2	55.7	27.2	75.3
Virginia	47.7	79.6	68.1	52.9	59.0	42.2	74.9
Washington	37.5	67.1	65.4	49.0	42.5	37.4	75.6
West Virginia	32.9	56.7	58.3	36.3	36.3	22.3	45.9
Wisconsin	35.9	57.8	57.3	40.9	36.9	21.2	63.5
Wyoming	31.5	56.4	54.1	42.0	32.5	19.0	66.1
State Median	39.0	65.6	58.5	44.9	42.5	25.8	63.9
State Range	18.1-59.4	17.9-84.8	30.4-75.9	22.7-60.2	14.3-59.0	7.2-44.0	40.7-75.9

TABLE 41. Percentage of Secondary Schools That Used Specific Types of Information to Identify Students with Poorly Controlled Asthma, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Frequent absences from school	Frequent visits to the school health office due to asthma	Frequent asthma symptoms at school	Frequent non-participation in physical education due to asthma	Students sent home early due to asthma	Calls from school to 911* due to asthma	At least 3 of these 6 types of information
LOCAL SURVEYS							
Albuquerque	64.8	92.2	86.7	40.5	81.7	41.0	91.7
Baltimore	53.3	75.3	48.2	27.6	43.4	25.4	61.2
Broward County	39.9	73.8	57.8	31.7	50.4	55.6	64.5
Charlotte-Mecklenberg County	37.8	81.2	70.1	47.4	58.7	53.3	67.0
Chicago	52.2	48.3	58.3	35.6	40.3	26.9	60.1
Detroit	49.4	32.9	50.6	34.4	45.3	18.3	53.9
District of Columbia	27.6	69.4	47.2	16.7	19.5	19.5	44.8
Fresno	56.0	84.1	80.8	59.9	72.0	48.4	75.8
Houston	50.6	60.7	52.9	51.9	38.9	17.3	57.7
Los Angeles	49.6	76.5	59.1	57.9	54.5	41.0	74.2
Memphis	56.6	43.2	54.8	25.4	47.2	43.2	53.4
Miami-Dade County	39.7	45.1	47.6	36.8	43.5	41.8	58.6
Milwaukee	52.6	64.6	54.6	34.6	41.0	32.1	61.2
New York City	56.5	72.7	58.8	45.5	49.5	34.2	68.1
Newark	57.6	80.4	57.0	38.0	46.1	30.8	62.7
Orange County	32.9	73.8	56.5	41.5	47.9	43.7	70.4
Philadelphia	55.9	83.3	64.3	45.6	57.5	41.7	68.8
San Diego	77.6	91.0	85.1	70.1	71.6	40.3	90.3
San Francisco	58.2	72.4	69.7	70.0	57.9	48.7	74.3
Local Median	52.6	73.8	57.8	40.5	47.9	41.0	64.5
Local Range	27.6-77.6	32.9-92.2	47.2-86.7	16.7-70.1	19.5-81.7	17.3-55.6	44.8-91.7
TERRITORIAL SURVEYS							
American Samoa	80.8	50.0	53.8	42.3	80.0	0.0	91.7
Guam	15.4	76.9	61.5	53.8	38.5	46.2	61.5
Marshall Islands	6.4	8.0	4.6	4.9	5.0	1.5	29.3
Northern Mariana Islands	50.0	37.5	37.5	37.5	25.0	37.5	60.0
Palau	0.0	0.0	0.0	0.0	9.1	0.0	0.0
Territorial Median	15.4	37.5	37.5	37.5	25.0	1.5	60.0
Territorial Range	0.0-80.8	0.0-76.9	0.0-61.5	0.0-53.8	5.0-80.0	0.0-46.2	0.0-91.7
TRIBAL SURVEYS							
Cherokee Nation	35.6	44.8	48.4	33.7	32.9	13.5	54.6
Nez Perce	14.3	0.0	28.6	42.9	14.3	14.3	66.7

*Or other local emergency numbers.

TABLE 42a. Percentage of Secondary Schools That Provided Specific Services for Students with Poorly Controlled Asthma, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Provided referrals to primary healthcare clinicians or child health insurance programs	Ensured an appropriate written asthma action plan is obtained	Ensured access to and appropriate use of asthma medications, spacers, and peak flow meters at school	Offered asthma education for students with asthma	Minimized asthma triggers in the school environment
STATE SURVEYS					
Alabama	64.9	80.2	78.4	59.4	73.4
Alaska	56.6	48.6	52.9	36.8	52.4
Arizona	52.5	64.1	75.2	34.0	62.1
Arkansas	54.8	85.8	90.2	57.2	75.8
California	67.7	75.3	83.0	44.3	65.4
Colorado	64.4	86.1	86.3	53.0	69.7
Connecticut	84.5	91.9	93.9	70.6	78.9
Delaware	82.3	91.8	98.3	82.4	93.7
Florida	45.1	80.0	79.6	43.9	68.1
Georgia	50.6	84.0	88.3	51.4	73.0
Hawaii	41.8	87.4	82.4	47.6	58.6
Idaho	53.5	69.5	77.2	39.9	66.1
Indiana	59.0	83.8	88.0	56.7	72.7
Iowa	66.6	77.4	86.5	46.4	71.2
Kansas	49.0	73.9	78.9	45.3	61.8
Kentucky	55.7	76.5	86.0	47.9	70.0
Louisiana	48.1	81.4	82.9	42.0	62.0
Maine	77.7	85.9	88.4	68.7	86.5
Maryland	63.0	89.9	86.4	63.2	67.6
Massachusetts	81.1	86.9	93.6	68.8	76.9
Michigan	37.6	61.2	74.0	27.6	62.1
Minnesota	67.1	83.3	88.5	63.9	80.7
Mississippi	51.4	66.0	77.4	46.9	69.6
Missouri	64.0	83.8	88.6	60.5	76.0
Montana	46.5	70.0	73.3	41.1	68.4
Nebraska	53.2	84.2	81.9	55.7	79.2
Nevada	65.3	82.7	82.4	56.5	75.3
New Hampshire	83.2	92.6	97.8	75.3	85.8
New Jersey	79.1	95.4	95.7	78.8	88.5
New Mexico	78.7	86.1	88.9	62.3	72.8
New York	76.2	81.9	87.8	65.9	75.0
North Carolina	65.1	87.7	85.7	57.3	73.3
North Dakota	34.0	47.7	60.2	21.9	52.7
Ohio	51.9	70.2	80.2	42.9	63.6
Oklahoma	35.9	65.8	77.9	36.1	64.3
Oregon	62.3	78.8	81.2	42.2	66.2
Pennsylvania	71.2	83.8	90.3	66.8	77.8
Rhode Island	83.4	91.0	94.4	72.3	81.5
South Carolina	69.1	90.9	93.4	68.2	82.6
South Dakota	49.0	52.1	58.9	29.0	56.2
Tennessee	54.4	90.9	89.1	58.4	82.5
Texas	60.5	78.4	89.8	55.9	69.6
Utah	33.8	75.8	73.3	40.5	65.5
Vermont	95.2	87.1	94.2	79.8	89.4
Virginia	64.6	95.4	95.4	71.9	83.2
Washington	70.2	92.7	85.4	48.5	72.4
West Virginia	66.0	89.5	88.4	68.8	82.1
Wisconsin	59.3	79.1	87.8	54.9	76.6
Wyoming	70.5	71.4	87.7	54.2	67.8
State Median	63.0	83.3	86.4	55.7	72.7
State Range	33.8-95.2	47.7-95.4	52.9-98.3	21.9-82.4	52.4-93.7

TABLE 42a. Percentage of Secondary Schools That Provided Specific Services for Students with Poorly Controlled Asthma, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Provided referrals to primary healthcare clinicians or child health insurance programs	Ensured an appropriate written asthma action plan is obtained	Ensured access to and appropriate use of asthma medications, spacers, and peak flow meters at school	Offered asthma education for students with asthma	Minimized asthma triggers in the school environment
LOCAL SURVEYS					
Albuquerque	94.8	100.0	97.1	97.4	81.2
Baltimore	68.4	82.7	86.3	64.2	72.2
Broward County	49.9	75.0	76.2	62.2	67.2
Charlotte-Mecklenberg County	66.6	94.2	94.3	76.7	73.0
Chicago	73.2	86.8	85.2	67.4	73.4
Detroit	58.8	71.9	72.2	57.3	69.5
District of Columbia	79.4	75.7	81.3	58.1	62.5
Fresno	96.2	92.3	96.2	80.8	71.4
Houston	69.2	63.0	75.1	38.9	40.5
Los Angeles	82.9	86.5	93.4	68.4	76.9
Memphis	55.6	82.9	84.3	58.0	81.6
Miami-Dade County	54.5	66.3	66.4	40.7	53.6
Milwaukee	70.0	87.0	91.3	72.3	77.1
New York City	77.6	75.9	79.5	63.2	67.3
Newark	92.9	97.7	97.5	86.4	90.8
Orange County	58.5	88.2	90.5	53.9	68.4
Philadelphia	85.8	87.4	90.7	75.0	76.0
San Diego	92.5	94.0	95.5	79.1	71.2
San Francisco	93.8	96.7	93.8	77.3	87.5
Local Median	73.2	86.8	90.5	67.4	72.2
Local Range	49.9-96.2	63.0-100.0	66.4-97.5	38.9-97.4	40.5-90.8
TERRITORIAL SURVEYS					
American Samoa	69.2	80.8	42.3	73.1	76.9
Guam	100.0	61.5	75.0	61.5	69.2
Marshall Islands	29.8	16.3	17.8	20.8	16.3
Northern Mariana Islands	50.0	37.5	57.1	25.0	50.0
Palau	100.0	0.0	0.0	0.0	0.0
Territorial Median	69.2	37.5	42.3	25.0	50.0
Territorial Range	29.8-100.0	0.0-80.8	0.0-75.0	0.0-73.1	0.0-76.9
TRIBAL SURVEYS					
Cherokee Nation	48.5	78.4	86.5	45.4	73.8
Nez Perce	28.6	42.9	71.4	28.6	28.6

TABLE 42b. Percentage of Secondary Schools That Provided Specific Services for Students with Poorly Controlled Asthma, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Addressed social and emotional issues related to asthma	Provided additional psychosocial counseling or support services as needed	Ensured access to safe, enjoyable physical education and activity opportunities	Ensured access to preventive medications before physical activity	Provided all 9 services
STATE SURVEYS					
Alabama	53.1	45.3	78.1	76.6	22.0
Alaska	36.8	30.8	67.1	62.4	18.2
Arizona	40.3	35.0	77.8	76.4	17.7
Arkansas	54.0	52.5	91.1	87.6	25.6
California	46.8	46.7	89.2	83.8	24.3
Colorado	53.6	54.1	89.1	85.8	21.7
Connecticut	70.1	72.3	94.1	95.1	34.6
Delaware	75.9	67.5	98.4	98.4	48.1
Florida	48.6	50.0	90.6	85.0	16.1
Georgia	49.7	52.8	91.5	92.1	20.9
Hawaii	48.1	65.0	89.7	80.9	19.3
Idaho	47.6	52.0	84.4	80.5	19.6
Indiana	47.5	44.3	91.0	89.8	23.4
Iowa	48.6	42.9	87.6	85.5	21.9
Kansas	42.9	41.1	88.5	88.1	17.8
Kentucky	52.0	56.0	89.9	89.1	26.1
Louisiana	40.2	41.8	85.9	81.0	18.3
Maine	60.7	56.7	94.9	92.1	34.5
Maryland	78.3	74.1	74.1	71.6	38.2
Massachusetts	70.3	70.9	95.6	95.1	41.5
Michigan	39.2	45.2	82.5	81.5	9.8
Minnesota	59.1	54.0	92.3	90.1	29.8
Mississippi	43.3	49.4	87.9	82.2	25.7
Missouri	55.7	51.9	90.3	90.8	26.2
Montana	34.1	47.5	84.2	80.5	13.7
Nebraska	47.4	35.6	93.1	90.5	21.2
Nevada	50.9	49.5	90.5	86.8	27.0
New Hampshire	71.0	60.3	97.8	95.0	41.8
New Jersey	76.1	71.9	97.3	97.3	46.9
New Mexico	60.3	60.3	90.8	93.3	36.4
New York	61.7	67.3	92.9	85.5	33.6
North Carolina	53.9	56.6	89.0	86.0	30.0
North Dakota	30.9	30.2	76.1	72.9	11.2
Ohio	42.4	39.7	79.4	76.2	17.9
Oklahoma	36.1	41.8	83.9	80.7	16.5
Oregon	41.7	44.7	88.4	84.2	20.9
Pennsylvania	63.4	61.0	93.0	90.4	31.3
Rhode Island	72.3	79.7	97.6	95.3	50.6
South Carolina	70.3	57.9	93.2	91.5	39.7
South Dakota	35.3	30.9	71.5	68.3	11.1
Tennessee	55.1	58.4	94.4	90.7	29.0
Texas	51.9	53.2	91.6	91.9	29.0
Utah	43.7	46.5	91.0	76.1	10.0
Vermont	81.6	67.4	96.8	95.0	50.6
Virginia	60.6	58.8	97.2	95.8	32.4
Washington	54.6	47.7	86.4	83.8	26.9
West Virginia	66.9	58.0	89.2	91.8	39.3
Wisconsin	55.5	49.8	93.1	90.5	26.0
Wyoming	46.8	48.2	89.8	83.0	24.6
State Median	52.0	52.0	90.3	86.8	25.7
State Range	30.9-81.6	30.2-79.7	67.1-98.4	62.4-98.4	9.8-50.6

TABLE 42b. Percentage of Secondary Schools That Provided Specific Services for Students with Poorly Controlled Asthma, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Addressed social and emotional issues related to asthma	Provided additional psychosocial counseling or support services as needed	Ensured access to safe, enjoyable physical education and activity opportunities	Ensured access to preventive medications before physical activity	Provided all 9 services
LOCAL SURVEYS					
Albuquerque	86.9	76.2	97.4	100.0	70.5
Baltimore	60.8	58.1	86.3	70.0	27.9
Broward County	55.4	64.4	93.7	82.7	26.3
Charlotte-Mecklenberg County	66.6	64.4	92.4	94.3	35.0
Chicago	60.6	52.9	90.8	78.5	30.2
Detroit	64.6	72.4	86.7	78.1	24.3
District of Columbia	50.1	50.1	84.4	74.2	31.4
Fresno	68.1	84.6	100.0	100.0	48.4
Houston	29.3	39.0	80.2	80.3	19.5
Los Angeles	64.3	74.6	88.5	93.6	42.2
Memphis	50.1	62.6	91.6	82.8	28.0
Miami-Dade County	52.3	67.8	89.5	71.7	16.8
Milwaukee	55.2	50.0	92.6	86.8	25.7
New York City	59.8	65.1	88.2	72.7	30.9
Newark	82.0	74.5	93.0	95.1	58.8
Orange County	64.5	69.2	90.4	83.2	22.0
Philadelphia	64.7	66.7	92.4	85.9	41.3
San Diego	65.2	66.7	93.9	92.5	47.0
San Francisco	84.0	77.5	90.8	84.6	54.6
Local Median	64.3	66.7	90.8	83.2	30.9
Local Range	29.3-86.9	39.0-84.6	80.2-100.0	70.0-100.0	16.8-70.5
TERRITORIAL SURVEYS					
American Samoa	57.7	57.7	80.8	73.1	26.9
Guam	61.5	61.5	76.9	76.9	25.0
Marshall Islands	28.3	28.3	30.2	23.7	10.4
Northern Mariana Islands	37.5	50.0	75.0	75.0	12.5
Palau	0.0	0.0	0.0	0.0	0.0
Territorial Median	37.5	50.0	75.0	73.1	12.5
Territorial Range	0.0-61.5	0.0-61.5	0.0-80.8	0.0-76.9	0.0-26.9
TRIBAL SURVEYS					
Cherokee Nation	49.0	43.1	90.0	84.3	21.5
Nez Perce	28.6	42.9	71.4	71.4	0.0

TABLE 43. Percentage of Secondary Schools That Had Adopted a Policy Stating That Students are Permitted to Carry and Self-Administer Asthma Medications and, Among Those Schools, the Percentage That Had Procedures to Inform Specific Groups About the Policy, the Percentage That Designated an Individual Responsible for Implementing the Policy,* and the Percentage of All Schools That Had a Fully Implemented Policy,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Had adopted a policy	Had procedures to inform†		Designated an individual responsible for implementing the policy†	Had a fully implemented policy
		Students	Parents and families		
STATE SURVEYS					
Alabama	71.5	96.4	92.6	79.5	52.2
Alaska	46.4	80.4	80.5	77.8	29.4
Arizona	57.3	87.4	87.3	78.6	38.3
Arkansas	77.6	93.3	94.8	78.7	55.4
California	66.1	89.2	89.2	79.2	44.8
Colorado	67.3	92.8	92.2	80.8	48.4
Connecticut	60.7	92.2	94.6	91.2	49.2
Delaware	70.9	93.1	92.8	86.4	55.8
Florida	73.4	91.1	93.4	79.7	50.7
Georgia	84.7	95.0	94.6	71.2	53.9
Hawaii	69.8	88.1	84.1	77.9	43.1
Idaho	77.8	85.2	82.2	68.5	42.8
Indiana	83.6	94.9	90.8	86.6	64.1
Iowa	74.5	91.0	89.7	82.4	52.6
Kansas	71.9	94.1	91.5	81.1	52.1
Kentucky	73.9	90.8	89.1	74.3	46.3
Louisiana	67.5	94.1	93.4	81.7	50.1
Maine	83.5	89.7	91.9	83.3	61.6
Maryland	55.5	91.3	92.8	83.7	41.2
Massachusetts	65.3	93.6	93.1	93.5	54.3
Michigan	75.9	85.8	87.8	63.8	40.7
Minnesota	73.3	92.3	91.9	83.7	54.4
Mississippi	70.6	92.5	91.1	75.6	45.5
Missouri	68.3	94.4	92.6	83.6	51.1
Montana	78.9	93.1	89.5	65.7	42.9
Nebraska	71.7	94.6	95.1	78.5	52.5
Nevada	84.3	90.6	89.2	82.7	59.5
New Hampshire	86.3	92.1	92.8	89.5	69.6
New Jersey	74.5	93.8	95.3	90.3	59.8
New Mexico	73.1	90.4	92.9	84.9	53.6
New York	67.1	89.5	91.8	87.0	50.4
North Carolina	84.2	95.2	93.8	80.9	62.9
North Dakota	55.9	89.3	87.2	73.6	33.1
Ohio	75.8	91.1	90.0	73.1	47.9
Oklahoma	78.2	94.8	94.4	75.4	54.5
Oregon	69.6	88.9	91.6	75.6	43.9
Pennsylvania	74.4	97.6	96.8	81.1	57.1
Rhode Island	75.3	95.4	96.8	95.3	66.7
South Carolina	78.3	96.2	97.4	87.3	65.2
South Dakota	39.9	91.1	92.3	70.3	21.9
Tennessee	83.5	97.3	95.5	82.8	64.8
Texas	75.3	96.2	96.2	82.2	58.6
Utah	86.7	87.6	86.8	65.2	45.3
Vermont	76.0	92.9	94.7	92.4	64.1
Virginia	82.5	98.2	98.2	76.7	60.7
Washington	80.0	95.8	91.7	77.8	56.6
West Virginia	79.9	97.1	93.9	83.5	62.4
Wisconsin	73.5	89.4	88.8	80.2	50.1
Wyoming	71.1	89.2	87.9	89.3	53.1
State Median	73.9	92.5	92.3	80.9	52.5
State Range	39.9-86.7	80.4-98.2	80.5-98.2	63.8-95.3	21.9-69.6

TABLE 43. Percentage of Secondary Schools That Had Adopted a Policy Stating That Students are Permitted to Carry and Self-Administer Asthma Medications and, Among Those Schools, the Percentage That Had Procedures to Inform Specific Groups About the Policy, the Percentage That Designated an Individual Responsible for Implementing the Policy,* and the Percentage of All Schools That Had a Fully Implemented Policy,† Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Had adopted a policy	Had procedures to inform‡		Designated an individual responsible for implementing the policy‡	Had a fully implemented policy
		Students	Parents and families		
LOCAL SURVEYS					
Albuquerque	91.9	94.3	94.3	90.9	78.0
Baltimore	39.5	86.3	86.8	78.8	27.2
Broward County	73.6	87.6	85.1	67.9	41.2
Charlotte-Mecklenberg County	70.2	91.1	94.1	76.6	49.3
Chicago	63.7	86.5	91.7	76.2	36.9
Detroit	62.8	86.5	84.0	76.8	37.8
District of Columbia	54.3	94.8	100.0	94.5	47.1
Fresno	84.6	85.6	90.4	85.6	59.8
Houston	38.4	90.3	87.1	92.6	28.1
Los Angeles	64.9	90.7	89.7	91.4	50.4
Memphis	63.6	97.7	96.9	68.0	34.3
Miami-Dade County	49.7	90.6	92.5	55.2	23.8
Milwaukee	55.3	91.6	83.2	83.7	35.2
New York City	57.4	81.2	85.3	79.8	34.0
Newark	36.7	93.8	93.4	93.4	30.4
Orange County	80.2	94.2	91.4	81.6	56.6
Philadelphia	39.2	94.2	94.3	91.5	30.1
San Diego	100.0	98.5	100.0	83.3	81.8
San Francisco	77.8	88.2	88.5	88.2	59.3
Local Median	63.6	90.7	91.4	85.3	37.8
Local Range	36.7-100.0	81.2-98.5	83.2-100.0	55.2-94.5	23.8-81.8
TERRITORIAL SURVEYS					
American Samoa	76.9	95.0	100.0	35.0	26.9
Guam	38.5	100.0	80.0	75.0	23.1
Marshall Islands	19.3	61.9	67.1	100.0	9.3
Northern Mariana Islands	37.5	100.0	100.0	100.0	28.6
Palau	0.0	NA	NA	NA	0.0
Territorial Median	37.5	97.5	90.0	87.5	23.1
Territorial Range	0.0-76.9	61.9-100.0	67.1-100.0	35.0-100.0	0.0-28.6
TRIBAL SURVEYS					
Cherokee Nation	75.9	96.6	92.2	80.2	55.2
Nez Perce	85.7	100.0	100.0	50.0	42.9

NA=Data not available.

*Principal, assistant principal, school nurse, or other school faculty or staff member.

†Have adopted a policy, have procedures to inform students and parents/families about the policy, and have designated an individual responsible for implementing the policy.

‡Among schools that have adopted a policy stating that students are permitted to carry and self-administer asthma medications.

TABLE 44. Percentage of Secondary Schools That Had Someone Who Oversees or Coordinates School Health and Safety Programs and Activities and the Percentage That Ever Used the School Health Index or Other Self-Assessment Tool to Assess School Policies, Activities, and Programs in Specific Areas, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Had someone who oversees or coordinates school health and safety programs and activities	Ever used School Health Index or other self-assessment tool					
		Asthma	Injury and violence prevention	Physical activity	Nutrition	Tobacco-use prevention	Physical activity, nutrition, and tobacco-use prevention
STATE SURVEYS							
Alabama	83.0	18.7	33.8	49.0	31.6	47.7	24.7
Alaska	71.0	16.2	24.6	28.3	26.3	27.4	21.9
Arizona	74.5	11.8	24.7	25.8	24.9	30.1	17.7
Arkansas	90.5	52.3	62.8	79.5	79.5	73.6	69.4
California	77.7	16.1	31.3	40.5	34.2	40.3	28.1
Colorado	78.3	24.7	43.6	49.5	45.7	49.9	35.5
Connecticut	81.8	22.2	30.7	44.1	41.0	36.4	31.3
Delaware	88.2	26.2	42.0	61.5	51.3	56.3	45.1
Florida	87.2	23.4	40.8	46.1	44.2	45.8	33.9
Georgia	87.3	31.0	42.3	52.4	53.8	54.6	43.9
Hawaii	87.0	39.3	47.5	54.2	49.5	57.3	43.9
Idaho	84.6	15.1	28.4	35.2	37.9	36.5	27.4
Indiana	87.1	18.2	29.4	35.5	37.1	47.7	28.7
Iowa	82.3	16.0	26.7	33.3	34.6	31.2	24.8
Kansas	80.9	18.1	28.7	43.5	42.6	43.4	35.6
Kentucky	85.1	20.3	35.3	48.7	45.5	44.0	35.6
Louisiana	85.6	19.4	34.1	37.1	37.8	38.3	29.2
Maine	78.7	40.0	45.4	55.4	54.1	53.1	49.5
Maryland	83.4	29.6	39.3	47.7	44.2	50.8	36.8
Massachusetts	84.8	26.4	37.5	45.3	45.4	41.3	37.2
Michigan	77.1	23.4	31.6	40.0	40.9	38.3	30.7
Minnesota	85.6	20.6	25.7	34.6	37.0	31.9	25.3
Mississippi	90.5	39.1	57.1	66.2	66.3	64.4	58.5
Missouri	90.4	31.7	38.5	50.3	47.0	45.4	38.6
Montana	84.7	36.5	45.4	55.0	56.7	58.3	50.6
Nebraska	86.5	22.9	28.4	33.6	33.1	29.8	27.2
Nevada	87.7	14.3	23.0	26.8	29.7	25.4	20.6
New Hampshire	83.5	26.3	36.6	45.0	44.2	42.4	36.7
New Jersey	92.8	29.7	36.5	42.3	41.4	39.1	33.1
New Mexico	85.9	33.3	44.8	46.5	47.6	46.2	38.9
New York	90.4	32.9	40.8	53.6	48.2	44.9	39.2
North Carolina	84.1	19.2	28.9	34.6	33.1	35.1	25.5
North Dakota	75.2	21.0	32.5	43.9	48.5	50.7	37.7
Ohio	76.3	18.0	25.9	37.2	36.7	32.9	26.8
Oklahoma	92.0	28.2	42.4	50.1	50.8	50.8	39.2
Oregon	79.6	16.3	32.0	34.1	34.4	34.4	26.4
Pennsylvania	88.4	21.7	32.5	47.0	47.7	35.2	30.0
Rhode Island	81.8	24.4	36.7	43.0	39.7	35.1	29.4
South Carolina	86.5	24.7	36.7	54.7	45.9	42.4	36.9
South Dakota	73.1	12.2	28.0	37.7	34.5	36.2	27.5
Tennessee	89.6	59.3	70.0	82.4	77.9	75.0	71.9
Texas	90.9	27.3	41.6	54.7	43.8	41.8	35.3
Utah	86.9	20.3	38.1	45.1	42.8	47.7	32.5
Vermont	79.7	39.7	49.7	58.0	61.1	58.6	54.7
Virginia	89.3	23.9	38.2	47.3	43.5	40.8	34.2
Washington	80.2	17.6	30.8	34.6	35.3	44.6	26.2
West Virginia	91.1	29.4	47.6	73.1	62.0	64.8	53.7
Wisconsin	85.0	23.8	40.2	51.2	50.2	53.3	41.5
Wyoming	79.0	18.2	31.0	35.7	32.1	42.6	26.8
State Median	85.0	23.4	36.6	45.3	43.8	43.4	34.2
State Range	71.0-92.8	11.8-59.3	23.0-70.0	25.8-82.4	24.9-79.5	25.4-75.0	17.7-71.9

TABLE 44. Percentage of Secondary Schools That Had Someone Who Oversees or Coordinates School Health and Safety Programs and Activities and the Percentage That Ever Used the School Health Index or Other Self-Assessment Tool to Assess School Policies, Activities, and Programs in Specific Areas, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Had someone who oversees or coordinates school health and safety programs and activities	Ever used School Health Index or other self-assessment tool					
		Asthma	Injury and violence prevention	Physical activity	Nutrition	Tobacco-use prevention	Physical activity, nutrition, and tobacco-use prevention
LOCAL SURVEYS							
Albuquerque	93.8	37.9	46.0	43.1	51.2	40.7	37.9
Baltimore	73.8	32.2	31.3	34.9	35.4	33.7	26.1
Broward County	86.7	45.9	49.6	52.5	45.9	47.5	41.0
Charlotte-Mecklenberg County	89.9	26.2	31.9	40.4	38.1	29.2	25.3
Chicago	90.0	31.9	31.3	31.3	35.0	26.0	20.2
Detroit	90.5	56.0	46.7	53.1	51.2	41.4	37.7
District of Columbia	77.7	22.8	20.5	34.2	28.5	17.5	14.6
Fresno	91.8	25.3	25.3	26.5	25.3	25.3	25.3
Houston	57.1	42.7	50.1	59.5	53.2	50.2	47.6
Los Angeles	98.2	32.1	55.1	56.1	49.3	55.8	41.9
Memphis	76.3	44.2	42.6	52.0	45.4	47.7	39.8
Miami-Dade County	96.9	45.8	71.4	77.5	73.6	61.1	58.4
Milwaukee	95.8	52.5	44.9	61.4	56.6	48.1	42.4
New York City	90.3	32.9	35.2	48.3	36.7	27.4	24.6
Newark	95.8	36.4	36.9	33.8	47.6	33.1	23.7
Orange County	100.0	25.6	41.8	52.3	48.8	44.0	32.6
Philadelphia	93.6	34.3	41.8	41.9	49.6	29.8	25.4
San Diego	71.2	11.9	20.9	25.4	16.4	22.4	14.9
San Francisco	96.7	53.1	74.8	71.8	68.8	65.0	56.1
Local Median	90.5	34.3	41.8	48.3	47.6	40.7	32.6
Local Range	57.1-100.0	11.9-56.0	20.5-74.8	25.4-77.5	16.4-73.6	17.5-65.0	14.6-58.4
TERRITORIAL SURVEYS							
American Samoa	100.0	69.2	92.3	96.2	96.2	96.2	96.2
Guam	91.7	54.5	66.7	50.0	54.5	66.7	45.5
Marshall Islands	58.3	29.2	62.1	80.8	81.9	73.9	65.2
Northern Mariana Islands	100.0	25.0	62.5	62.5	62.5	62.5	62.5
Palau	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Territorial Median	100.0	29.2	62.5	62.5	62.5	66.7	62.5
Territorial Range	58.3-100.0	0.0-69.2	0.0-92.3	0.0-96.2	0.0-96.2	0.0-96.2	0.0-96.2
TRIBAL SURVEYS							
Cherokee Nation	95.6	38.6	47.6	59.6	59.3	59.9	49.3
Nez Perce	71.4	14.3	14.3	85.7	85.7	85.7	85.7

TABLE 45. Percentage of Secondary Schools With a School Improvement Plan (SIP) That Includes Health-Related Goals and Objectives on Specific Topics and the Percentage of Schools With an SIP That Includes Any Health-Related Goals and Objectives,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Health education	Physical education and physical activity	Nutrition services and foods and beverages available at school	Health services	Mental health and social services	Healthy and safe school environment	Family and community involvement	Faculty and staff health promotion	SIP includes any health-related goals
STATE SURVEYS									
Alabama	36.1	28.2	23.5	26.0	29.5	55.4	55.4	18.5	82.4
Alaska	19.9	21.2	19.1	15.0	18.5	36.8	52.3	15.6	56.2
Arizona	17.0	18.3	14.6	14.2	15.7	45.5	53.6	14.2	58.9
Arkansas	80.1	80.1	72.4	74.9	64.0	80.5	80.2	59.6	86.3
California	33.5	42.8	25.0	29.8	36.2	61.9	63.6	19.9	69.9
Colorado	28.2	27.6	23.9	22.5	27.7	64.1	64.3	21.7	74.7
Connecticut	28.1	34.4	25.8	23.9	33.5	56.6	56.4	17.5	69.1
Delaware	35.1	29.0	23.1	27.3	34.4	74.4	77.3	20.2	86.4
Florida	30.3	37.3	21.5	23.3	18.2	67.5	77.2	22.2	84.3
Georgia	31.8	31.4	24.6	23.7	27.0	63.3	71.8	23.7	78.6
Hawaii	53.4	57.5	42.6	49.6	66.7	85.2	92.2	34.3	94.3
Idaho	43.9	42.3	38.0	33.5	34.2	62.7	61.4	36.2	71.3
Indiana	26.4	28.0	22.4	21.6	21.0	64.3	62.9	17.4	72.0
Iowa	54.2	56.6	52.6	42.2	32.7	72.1	56.7	34.0	80.0
Kansas	38.6	39.4	34.8	28.4	21.2	44.0	41.7	33.4	57.7
Kentucky	39.0	41.6	29.1	29.6	29.8	60.7	69.2	21.0	78.4
Louisiana	17.6	20.8	14.0	14.4	17.9	60.2	79.6	15.8	83.6
Maine	24.3	24.2	23.9	22.2	20.5	25.8	23.7	21.4	28.7
Maryland	31.8	31.4	18.1	30.4	42.1	84.0	88.6	30.3	93.2
Massachusetts	33.8	30.1	23.9	27.3	38.4	76.4	76.7	20.6	86.9
Michigan	24.5	27.4	19.9	16.4	20.3	47.5	53.4	14.4	65.2
Minnesota	22.6	24.4	24.4	18.6	23.5	39.6	38.2	25.6	48.4
Mississippi	55.8	55.1	56.8	53.0	49.6	66.8	67.5	48.5	72.0
Missouri	50.3	51.5	44.8	48.6	38.1	74.2	76.4	43.1	83.6
Montana	43.2	45.0	40.5	33.1	36.9	60.5	56.2	30.4	65.7
Nebraska	34.5	35.6	33.0	31.0	25.0	50.4	49.7	31.6	58.7
Nevada	10.1	12.7	5.7	8.1	9.6	39.6	60.8	9.6	62.4
New Hampshire	26.1	27.9	27.7	26.0	21.3	33.7	32.8	22.5	38.3
New Jersey	35.2	34.9	32.8	32.6	28.8	39.1	33.9	24.0	45.5
New Mexico	41.0	46.5	45.7	44.6	46.3	65.8	72.7	38.5	78.4
New York	36.6	39.8	33.1	34.5	34.8	43.8	43.4	27.5	51.7
North Carolina	53.4	55.9	32.1	34.3	29.0	83.0	81.5	38.3	88.4
North Dakota	41.6	41.4	43.8	26.1	26.9	51.2	45.9	36.2	63.3
Ohio	38.2	40.0	39.9	35.8	32.5	56.1	53.6	31.8	68.2
Oklahoma	54.1	58.3	55.7	43.1	40.0	61.7	55.2	35.7	68.1
Oregon	19.0	15.9	15.7	15.0	21.1	61.0	59.2	11.8	72.6
Pennsylvania	29.4	31.8	29.5	33.1	35.7	45.5	45.8	26.7	53.3
Rhode Island	36.8	33.7	31.9	30.4	35.1	62.2	69.2	27.5	74.8
South Carolina	43.4	46.6	35.3	41.2	42.0	69.2	73.1	37.6	78.5
South Dakota	27.1	27.5	27.9	23.7	22.0	42.8	44.8	24.4	55.6
Tennessee	41.6	45.5	30.5	37.2	37.0	69.9	77.2	33.6	81.2
Texas	56.8	58.9	46.2	48.4	40.8	80.4	80.4	42.2	85.2
Utah	37.0	39.9	32.8	26.7	28.6	69.7	65.4	33.5	79.3
Vermont	29.7	36.3	35.6	32.5	35.4	57.5	48.3	36.2	63.2
Virginia	39.3	45.3	27.6	27.9	27.1	68.2	68.5	31.1	78.0
Washington	27.5	28.5	9.6	12.8	17.1	56.6	65.2	9.7	78.3
West Virginia	34.9	40.1	34.6	29.3	27.3	48.6	45.3	33.2	54.3
Wisconsin	31.6	30.5	31.5	25.7	23.0	44.2	43.2	31.0	52.1
Wyoming	23.8	23.1	18.5	17.0	18.8	36.4	36.2	21.5	47.7
State Median	34.9	35.6	29.5	28.4	29.0	60.7	60.8	27.5	72.0
State Range	10.1-80.1	12.7-80.1	5.7-72.4	8.1-74.9	9.6-66.7	25.8-85.2	23.7-92.2	9.6-59.6	28.7-94.3

TABLE 45. Percentage of Secondary Schools With a School Improvement Plan (SIP) That Includes Health-Related Goals and Objectives on Specific Topics and the Percentage of Schools With an SIP That Includes Any Health-Related Goals and Objectives,* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Health education	Physical education and physical activity	Nutrition services and foods and beverages available at school	Health services	Mental health and social services	Healthy and safe school environment	Family and community involvement	Faculty and staff health promotion	SIP includes any health-related goals
LOCAL SURVEYS									
Albuquerque	54.5	57.2	51.9	51.9	57.2	69.9	77.1	47.7	83.2
Baltimore	40.5	39.3	23.7	39.9	50.9	86.2	89.0	28.8	91.4
Broward County	21.3	24.9	16.2	18.0	22.6	53.1	70.3	20.4	77.2
Charlotte-Mecklenberg County	27.9	33.3	10.1	18.0	22.3	70.3	88.3	22.6	88.3
Chicago	49.2	60.8	27.8	40.9	51.7	81.9	87.0	28.7	90.1
Detroit	63.4	63.0	56.5	53.3	62.8	76.6	81.1	49.9	84.6
District of Columbia	45.7	46.9	39.3	48.4	64.6	82.2	88.1	38.2	88.1
Fresno	35.1	39.3	26.8	35.1	38.7	43.5	38.7	22.0	43.5
Houston	47.9	49.1	54.2	59.5	58.7	72.9	70.3	46.6	80.2
Los Angeles	57.9	56.6	51.8	65.4	68.0	77.7	77.7	52.4	80.3
Memphis	50.8	50.9	41.9	43.4	53.3	83.1	86.3	38.1	89.5
Miami-Dade County	42.2	56.5	31.9	35.8	28.8	65.2	90.0	30.9	91.8
Milwaukee	31.3	28.7	17.5	26.9	34.4	84.9	93.2	21.1	94.5
New York City	37.2	40.6	30.3	35.1	37.0	41.2	42.2	25.4	48.5
Newark	37.2	38.7	43.6	46.4	43.8	50.0	57.3	23.1	57.3
Orange County	20.5	29.4	23.9	14.1	22.7	54.1	67.0	20.3	71.4
Philadelphia	43.5	49.3	40.2	50.4	67.5	84.4	92.3	31.2	94.4
San Diego	9.0	9.0	6.0	13.6	13.4	28.4	35.8	6.0	37.3
San Francisco	47.1	48.5	40.0	55.2	55.2	58.5	58.5	38.7	58.5
Local Median	42.2	46.9	31.9	40.9	50.9	70.3	77.7	28.8	83.2
Local Range	9.0-63.4	9.0-63.0	6.0-56.5	13.6-65.4	13.4-68.0	28.4-86.2	35.8-93.2	6.0-52.4	37.3-94.5
TERRITORIAL SURVEYS									
American Samoa	100.0	100.0	96.2	100.0	92.0	96.2	100.0	96.2	100.0
Guam	15.4	15.4	15.4	7.7	7.7	46.2	53.8	15.4	61.5
Marshall Islands	90.7	89.3	53.8	61.6	39.8	86.6	87.8	62.5	97.0
Northern Mariana Islands	87.5	85.7	75.0	37.5	37.5	75.0	75.0	62.5	87.5
Palau	100.0	90.9	81.8	63.6	27.3	90.9	81.8	100.0	100.0
Territorial Median	90.7	89.3	75.0	61.6	37.5	86.6	81.8	62.5	97.0
Territorial Range	15.4-100.0	15.4-100.0	15.4-96.2	7.7-100.0	7.7-92.0	46.2-96.2	53.8-100.0	15.4-100.0	61.5-100.0
TRIBAL SURVEYS									
Cherokee Nation	61.1	60.9	60.5	44.3	43.6	61.4	55.8	39.0	70.5
Nez Perce	71.4	57.1	71.4	71.4	57.1	85.7	71.4	42.9	100.0

*Among schools with a School Improvement Plan.

TABLE 46. Percentage of Secondary Schools Required to Report to their School District the Following Types of Information Regarding Implementation of the Local School Wellness Policy, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Number of minutes of physical education required in each grade	Rates of student participation in school meal programs	Revenue from sale of foods and beverages*	Number of minutes of physical activity outside of physical education [†]
STATE SURVEYS				
Alabama	89.0	82.0	68.8	32.6
Alaska	57.9	78.9	44.5	33.1
Arizona	50.5	73.9	52.9	34.7
Arkansas	89.6	92.2	73.0	59.8
California	90.5	91.6	74.0	54.6
Colorado	50.2	83.9	58.4	32.3
Connecticut	82.5	88.9	64.3	47.0
Delaware	72.6	86.4	62.3	33.1
Florida	76.2	91.2	77.4	37.3
Georgia	68.2	94.1	64.2	28.2
Hawaii	50.9	88.5	72.9	37.3
Idaho	62.7	87.6	58.7	34.8
Indiana	72.6	94.5	71.4	27.0
Iowa	89.8	88.6	67.0	61.2
Kansas	74.2	89.7	53.8	39.2
Kentucky	71.9	93.0	81.8	44.2
Louisiana	93.6	94.9	72.3	58.5
Maine	61.1	81.9	58.2	34.7
Maryland	65.9	90.7	79.9	22.6
Massachusetts	68.6	87.1	60.2	29.9
Michigan	53.7	86.6	59.2	25.7
Minnesota	58.3	79.9	60.7	29.0
Mississippi	92.3	94.5	83.6	69.1
Missouri	87.5	95.8	65.6	52.1
Montana	80.3	86.1	52.6	48.2
Nebraska	79.4	88.5	64.5	45.2
Nevada	81.8	89.8	60.5	63.5
New Hampshire	70.8	83.8	63.3	39.4
New Jersey	87.7	88.4	69.9	46.7
New Mexico	70.1	89.6	67.1	37.4
New York	84.8	89.2	75.9	44.5
North Carolina	68.9	87.1	72.3	45.4
North Dakota	82.6	81.2	38.2	43.5
Ohio	59.0	83.7	65.8	27.4
Oklahoma	74.4	90.6	66.6	58.3
Oregon	79.9	91.1	55.5	39.6
Pennsylvania	71.7	88.6	68.1	35.2
Rhode Island	83.0	93.7	67.4	32.8
South Carolina	80.8	94.6	74.9	44.3
South Dakota	56.2	74.8	50.4	36.2
Tennessee	93.6	98.2	82.5	85.0
Texas	83.2	92.4	74.0	37.5
Utah	68.8	93.5	73.6	30.3
Vermont	50.1	72.5	42.2	22.2
Virginia	78.7	90.2	73.5	32.6
Washington	73.0	88.1	68.3	31.5
West Virginia	92.3	96.3	75.2	53.9
Wisconsin	78.6	83.7	59.6	43.7
Wyoming	68.0	86.7	38.6	24.0
State Median	74.2	88.6	66.6	37.4
State Range	50.1-93.6	72.5-98.2	38.2-83.6	22.2-85.0

TABLE 46. Percentage of Secondary Schools Required to Report to their School District the Following Types of Information Regarding Implementation of the Local School Wellness Policy, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (*continued*)

Site	Number of minutes of physical education required in each grade	Rates of student participation in school meal programs	Revenue from sale of foods and beverages*	Number of minutes of physical activity outside of physical education†
LOCAL SURVEYS				
Albuquerque	78.3	91.7	75.0	34.7
Baltimore	54.2	87.9	77.6	21.2
Broward County	80.9	95.5	77.6	32.2
Charlotte-Mecklenberg County	75.3	92.3	80.1	59.5
Chicago	85.5	87.1	79.6	65.5
Detroit	78.2	96.5	87.2	46.0
District of Columbia	48.5	88.6	54.3	27.7
Fresno	96.2	91.8	72.0	59.3
Houston	86.4	88.8	82.5	39.5
Los Angeles	94.9	99.2	93.4	64.8
Memphis	78.3	94.7	87.9	56.9
Miami-Dade County	86.8	99.1	93.5	49.2
Milwaukee	74.4	93.1	64.0	49.5
New York City	78.7	89.2	70.8	42.7
Newark	97.8	91.9	85.4	57.1
Orange County	69.0	100.0	81.9	23.7
Philadelphia	75.9	97.2	60.6	43.9
San Diego	82.1	91.0	76.1	59.7
San Francisco	88.4	100.0	79.2	52.6
Local Median	78.7	92.3	79.2	49.2
Local Range	48.5-97.8	87.1-100.0	54.3-93.5	21.2-65.5
TERRITORIAL SURVEYS				
American Samoa	100.0	100.0	57.7	96.2
Guam	69.2	84.6	83.3	38.5
Marshall Islands	79.2	52.1	20.9	64.7
Northern Mariana Islands	100.0	100.0	100.0	100.0
Palau	100.0	100.0	18.2	81.8
Territorial Median	100.0	100.0	57.7	81.8
Territorial Range	69.2-100.0	52.1-100.0	18.2-100.0	38.5-100.0
TRIBAL SURVEYS				
Cherokee Nation	79.3	90.8	73.5	65.1
Nez Perce	57.1	85.7	42.9	28.6

*From sale of foods and beverages from school-sponsored fundraisers, vending machines, school stores, or a la carte lines in the school cafeteria.

†Such as classroom physical activity breaks, free time physical activity, or recess.

TABLE 47a. Percentage of Secondary Schools That Had One or More School Health Councils* and, Among Schools with Councils, the Percentage That Have Specific Groups Represented, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Groups represented†								
	School health council	School administrators	Health education teachers	Physical education teachers	Mental health or social services staff	Nutrition or food service staff	Health services staff‡	Maintenance and transportation staff	Technology staff
STATE SURVEYS									
Alabama	36.1	87.6	84.3	84.4	34.3	54.0	69.5	23.4	24.9
Alaska	33.1	85.7	69.5	72.6	61.8	47.1	49.8	25.5	35.2
Arizona	41.3	85.5	66.0	75.0	44.6	43.0	50.4	19.5	23.1
Arkansas	80.4	98.2	97.2	97.1	62.1	86.5	95.8	30.5	33.4
California	41.1	91.7	70.8	83.9	58.0	55.3	67.6	17.9	18.2
Colorado	50.7	90.2	84.4	90.0	66.1	59.7	81.0	14.1	17.2
Connecticut	55.1	97.0	91.5	91.8	81.0	64.9	86.0	30.3	17.4
Delaware	68.4	97.6	95.3	90.3	73.6	59.1	88.4	17.1	24.8
Florida	46.3	90.3	83.6	91.6	52.7	66.8	74.2	16.3	23.8
Georgia	51.3	91.6	98.6	96.3	58.3	75.1	74.9	21.8	22.1
Hawaii	64.6	94.1	89.0	86.7	46.8	63.0	60.6	30.2	30.8
Idaho	47.8	98.4	94.0	95.7	58.4	73.8	59.5	24.8	26.6
Indiana	63.6	96.9	94.9	93.4	53.9	81.3	86.0	19.8	18.3
Iowa	64.4	95.5	86.9	88.8	44.2	84.6	90.2	14.6	19.9
Kansas	63.7	94.3	86.0	91.7	34.3	85.2	79.8	19.1	16.5
Kentucky	49.1	95.8	88.6	91.5	57.3	70.3	70.6	18.0	20.3
Louisiana	44.0	93.4	82.2	88.4	48.0	54.1	72.6	15.7	29.1
Maine	66.0	92.4	89.9	85.5	60.7	71.8	86.1	28.7	19.2
Maryland	50.4	88.8	89.5	86.1	62.3	33.6	78.0	16.2	24.1
Massachusetts	64.0	95.5	89.6	89.9	76.7	62.5	87.5	12.4	14.0
Michigan	60.5	98.5	94.2	91.6	53.3	61.0	41.3	14.4	16.5
Minnesota	62.4	93.7	88.7	88.9	69.0	74.2	84.0	39.2	22.1
Mississippi	73.8	98.0	89.3	94.7	56.5	86.0	77.5	24.9	27.7
Missouri	62.8	94.7	86.9	86.9	50.3	75.6	89.5	21.0	21.8
Montana	53.2	91.8	96.3	94.4	38.2	70.5	54.0	32.2	31.5
Nebraska	52.6	96.9	87.1	90.0	33.9	65.3	76.8	30.3	26.8
Nevada	34.1	86.8	88.1	98.2	43.7	46.1	67.8	17.5	21.0
New Hampshire	71.3	96.6	90.7	90.5	48.1	85.1	94.0	29.2	21.2
New Jersey	57.6	94.5	94.5	93.7	65.7	59.5	85.7	18.9	25.6
New Mexico	59.9	95.5	75.2	77.7	87.8	60.8	85.7	20.0	25.9
New York	67.4	97.7	91.3	89.7	75.7	77.1	88.3	33.9	30.0
North Carolina	49.9	84.8	90.7	90.2	55.3	49.0	79.4	17.8	26.7
North Dakota	50.1	95.1	93.4	87.5	28.5	74.8	32.8	10.9	26.3
Ohio	52.5	93.7	92.5	96.0	55.1	80.2	81.9	24.8	24.1
Oklahoma	73.5	97.9	79.1	87.0	41.4	60.0	49.9	25.6	30.1
Oregon	36.4	91.8	80.3	84.6	60.4	54.2	49.3	20.5	17.9
Pennsylvania	77.4	93.9	89.1	91.6	63.0	80.3	90.1	27.7	33.9
Rhode Island	53.4	95.7	97.9	93.9	76.4	81.4	85.0	20.1	13.5
South Carolina	58.0	93.5	93.4	94.6	44.8	64.1	83.1	24.0	25.0
South Dakota	50.7	98.0	94.7	92.8	31.4	83.4	49.9	11.0	16.2
Tennessee	74.6	95.2	90.8	95.2	58.0	77.2	86.0	18.5	26.6
Texas	70.0	94.7	88.4	89.9	49.8	69.3	87.3	24.2	32.6
Utah	47.7	91.8	94.1	88.1	52.8	55.3	59.7	28.9	28.9
Vermont	73.5	94.4	81.5	79.9	74.2	68.8	92.2	25.0	11.0
Virginia	58.3	91.5	93.7	93.6	54.0	66.8	85.7	16.3	23.6
Washington	48.1	93.4	87.9	82.0	61.6	44.7	76.3	18.9	17.7
West Virginia	64.5	93.4	92.1	91.7	48.5	72.2	84.2	33.5	33.1
Wisconsin	63.1	95.4	92.4	94.0	55.1	78.5	78.1	21.5	25.9
Wyoming	52.2	94.3	95.5	94.0	68.8	62.9	95.4	13.8	23.1
State Median	57.6	94.3	89.6	90.3	55.3	66.8	79.8	20.5	24.1
State Range	33.1-80.4	84.8-98.5	66.0-98.6	72.6-98.2	28.5-87.8	33.6-86.5	32.8-95.8	10.9-39.2	11.0-35.2

TABLE 47a. Percentage of Secondary Schools That Had One or More School Health Councils* and, Among Schools with Councils, the Percentage That Have Specific Groups Represented, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Groups represented†								
	School health council	School administrators	Health education teachers	Physical education teachers	Mental health or social services staff	Nutrition or food service staff	Health services staff‡	Maintenance and transportation staff	Technology staff
LOCAL SURVEYS									
Albuquerque	81.7	96.8	73.8	56.9	96.8	33.8	96.8	6.7	13.4
Baltimore	41.2	90.1	77.3	70.8	86.4	60.4	76.5	30.0	50.1
Broward County	40.0	100.0	77.3	92.0	73.3	68.9	85.3	29.3	41.3
Charlotte-Mecklenberg County	50.4	82.3	100.0	100.0	54.7	41.0	86.0	18.0	32.0
Chicago	44.4	82.8	71.4	90.5	79.1	47.7	74.0	20.1	32.8
Detroit	62.8	90.2	86.1	96.0	79.5	71.5	63.1	16.5	49.0
District of Columbia	64.0	89.9	86.3	86.2	66.4	33.2	71.2	24.9	33.3
Fresno	41.7	100.0	80.8	100.0	90.4	71.2	90.4	9.6	19.2
Houston	50.2	77.0	97.4	92.4	55.0	63.4	73.1	50.3	52.6
Los Angeles	60.2	90.2	83.2	70.5	85.7	49.5	84.0	37.9	40.8
Memphis	45.1	89.3	96.6	88.3	63.5	64.5	76.7	52.3	61.2
Miami-Dade County	71.9	97.3	76.2	92.9	83.5	78.1	52.6	13.4	31.3
Milwaukee	60.2	85.9	71.7	85.9	70.6	56.9	75.8	13.4	27.0
New York City	57.4	97.3	85.2	87.7	83.3	70.6	81.5	37.4	35.3
Newark	90.1	88.1	86.0	89.9	87.1	76.1	88.5	19.7	36.2
Orange County	81.9	85.5	77.9	94.2	68.5	55.4	72.0	3.0	8.6
Philadelphia	49.3	94.0	80.6	85.3	67.2	69.7	84.0	29.5	42.7
San Diego	35.8	95.8	62.5	91.7	83.3	33.3	87.5	8.7	25.0
San Francisco	90.5	85.6	75.0	69.6	100.0	32.8	66.6	15.2	21.6
Local Median	57.4	90.1	80.6	89.9	79.5	60.4	76.7	19.7	33.3
Local Range	35.8-90.5	77.0-100.0	62.5-100.0	56.9-100.0	54.7-100.0	32.8-78.1	52.6-96.8	3.0-52.3	8.6-61.2
TERRITORIAL SURVEYS									
American Samoa	100.0	100.0	100.0	100.0	73.1	76.9	30.8	53.8	57.7
Guam	76.9	80.0	90.0	90.0	30.0	20.0	80.0	0.0	0.0
Marshall Islands	45.9	89.8	93.3	89.9	44.8	57.1	42.8	29.4	28.5
Northern Mariana Islands	75.0	100.0	100.0	100.0	60.0	80.0	60.0	60.0	60.0
Palau	63.6	100.0	100.0	100.0	14.3	85.7	28.6	28.6	85.7
Territorial Median	75.0	100.0	100.0	100.0	44.8	76.9	42.8	29.4	57.7
Territorial Range	45.9-100.0	80.0-100.0	90.0-100.0	89.9-100.0	14.3-73.1	20.0-85.7	28.6-80.0	0.0-60.0	0.0-85.7
TRIBAL SURVEYS									
Cherokee Nation	80.6	100.0	86.6	92.6	44.3	76.2	68.6	31.9	35.1
Nez Perce	57.1	100.0	100.0	100.0	75.0	100.0	75.0	25.0	25.0

*A group, committee, or team that offers guidance on the development of policies or coordinates activities on health topics.

†Among schools with school health councils.

‡For example, school nurse.

TABLE 47b. Percentage of Secondary Schools That Had One or More School Health Councils,* and Among Schools with Councils, the Percentage That Had Specific Groups Represented, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Groups represented†								
	Library/media center staff	Student body	Parents or families of students	Community members	Local health departments, agencies, or organizations	Faith-based organizations	Businesses	Local government agencies	10 or more groups represented
STATE SURVEYS									
Alabama	22.3	48.6	54.1	41.7	30.5	20.0	19.5	25.7	9.6
Alaska	25.4	53.3	57.3	61.7	56.2	18.8	26.4	38.1	11.7
Arizona	18.9	39.1	56.4	40.7	27.7	6.5	12.2	14.8	8.0
Arkansas	32.5	74.6	89.5	76.2	52.7	21.0	37.2	33.4	43.6
California	15.4	55.5	68.5	56.1	40.2	8.3	15.4	27.4	12.0
Colorado	16.9	44.4	60.9	57.3	50.5	10.8	21.5	20.7	13.6
Connecticut	13.6	40.5	53.0	46.8	39.4	7.0	18.7	19.0	13.7
Delaware	17.1	46.3	69.8	54.6	37.3	5.3	21.8	14.9	21.8
Florida	29.3	53.4	52.2	42.4	38.4	14.5	24.0	23.0	13.3
Georgia	32.1	51.9	59.3	52.0	43.9	17.4	31.6	28.1	16.8
Hawaii	25.0	66.2	58.7	42.8	17.9	2.5	8.4	6.4	17.7
Idaho	22.0	55.4	68.1	60.3	41.8	14.0	26.9	25.5	15.0
Indiana	19.6	59.6	73.0	69.1	46.1	13.5	30.6	25.9	26.0
Iowa	14.7	68.1	74.2	69.0	48.4	11.6	30.5	23.9	24.4
Kansas	11.5	50.2	59.5	56.5	39.3	5.9	21.3	13.8	15.3
Kentucky	23.7	54.3	74.7	58.5	55.9	12.6	20.9	26.1	15.9
Louisiana	22.1	45.0	53.8	40.5	34.4	10.3	19.9	21.3	10.2
Maine	21.2	47.8	63.5	65.7	56.4	9.6	17.7	17.4	20.9
Maryland	27.6	40.7	51.8	39.0	37.5	9.0	16.5	18.2	12.2
Massachusetts	13.5	50.9	67.4	63.3	53.1	6.5	12.5	24.7	19.1
Michigan	14.8	65.8	86.3	79.2	61.4	46.9	24.8	23.4	20.4
Minnesota	16.8	41.6	66.7	60.2	46.5	8.4	16.9	27.0	21.8
Mississippi	42.0	71.5	75.9	68.7	47.1	31.9	41.7	35.2	36.5
Missouri	23.4	60.9	71.4	65.9	56.3	16.5	35.4	30.3	26.4
Montana	24.6	48.5	65.3	63.7	49.2	11.3	22.6	28.1	16.7
Nebraska	19.4	54.3	60.7	64.4	42.3	9.1	18.2	25.5	14.8
Nevada	19.1	35.5	50.0	38.8	28.4	1.9	9.6	13.6	6.4
New Hampshire	17.3	37.6	68.6	57.1	31.9	2.5	9.4	17.0	17.8
New Jersey	21.1	58.6	60.3	53.5	37.7	11.1	15.7	23.0	16.8
New Mexico	22.7	59.4	68.5	56.6	63.7	18.3	28.7	32.9	23.7
New York	23.6	69.3	81.1	64.9	48.3	9.8	16.9	20.2	28.6
North Carolina	26.0	40.6	58.0	52.1	46.3	18.0	26.0	32.3	16.9
North Dakota	25.7	57.6	70.8	59.4	44.2	12.4	15.1	21.1	11.3
Ohio	17.2	51.0	64.1	57.3	41.3	9.4	15.8	20.1	16.2
Oklahoma	28.4	87.7	90.1	68.5	33.1	11.2	19.5	21.9	22.0
Oregon	12.7	45.3	60.3	53.2	42.4	10.0	22.6	27.0	8.8
Pennsylvania	20.6	56.9	68.5	61.4	47.9	12.5	24.6	20.3	32.5
Rhode Island	13.7	65.1	80.3	80.3	63.6	10.6	27.6	33.5	23.1
South Carolina	30.8	52.0	66.4	56.7	42.6	25.4	28.0	23.7	20.0
South Dakota	13.8	60.1	66.3	64.4	39.6	5.8	28.2	15.4	15.8
Tennessee	29.5	65.5	75.9	58.3	43.0	12.3	22.3	24.9	25.5
Texas	25.9	48.5	83.0	80.2	42.0	23.3	53.0	26.9	31.6
Utah	21.0	59.2	74.7	69.7	45.6	9.1	19.7	28.7	16.6
Vermont	13.4	40.8	62.5	57.7	36.2	4.7	15.8	18.1	18.2
Virginia	21.2	47.9	52.0	47.3	46.2	12.3	23.5	35.4	18.3
Washington	19.7	39.7	54.3	41.0	33.4	10.0	12.4	20.4	9.6
West Virginia	31.3	65.4	68.5	64.9	51.2	18.5	46.7	31.8	29.0
Wisconsin	21.0	54.2	72.3	63.0	46.4	16.3	18.4	17.1	21.4
Wyoming	21.2	65.0	77.4	65.0	47.1	17.8	25.6	29.4	20.6
State Median	21.2	53.4	66.7	58.5	43.9	11.2	21.5	23.9	17.7
State Range	11.5-42.0	35.5-87.7	50.0-90.1	38.8-80.3	17.9-63.7	1.9-46.9	8.4-53.0	6.4-38.1	6.4-43.6

TABLE 47b. Percentage of Secondary Schools That Had One or More School Health Councils,* and Among Schools with Councils, the Percentage That Had Specific Groups Represented, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Groups represented†								
	Library/media center staff	Student body	Parents or families of students	Community members	Local health departments, agencies, or organizations	Faith-based organizations	Businesses	Local government agencies	10 or more groups represented
LOCAL SURVEYS									
Albuquerque	13.4	33.5	44.0	44.0	33.2	3.2	10.5	19.8	11.2
Baltimore	40.1	53.7	69.2	80.6	74.6	36.9	58.8	36.8	21.0
Broward County	45.3	73.8	65.8	54.7	55.1	24.9	38.2	42.7	21.4
Charlotte-Mecklenberg County	32.3	32.3	19.5	13.7	40.7	4.7	4.7	9.0	4.5
Chicago	35.9	56.3	65.9	61.2	63.6	24.4	19.7	27.0	16.6
Detroit	33.2	71.3	83.9	81.8	79.1	56.2	56.2	53.0	41.9
District of Columbia	28.6	55.0	54.9	44.9	44.9	24.9	31.5	42.1	24.2
Fresno	19.2	61.6	80.8	39.7	12.1	0.0	0.0	0.0	8.0
Houston	47.6	63.5	61.2	58.6	54.9	37.8	42.7	40.1	27.3
Los Angeles	37.9	62.5	74.6	69.1	52.5	13.9	28.9	28.7	25.5
Memphis	52.4	73.3	81.1	77.6	37.4	50.8	45.1	33.0	29.5
Miami-Dade County	39.7	69.5	69.8	50.7	39.5	5.3	29.4	24.0	24.8
Milwaukee	17.7	29.6	36.7	32.0	22.0	7.4	5.2	10.3	7.7
New York City	27.0	69.4	74.0	47.8	53.4	9.7	15.9	22.7	22.4
Newark	26.8	87.8	82.0	60.8	38.4	12.7	21.4	12.8	35.4
Orange County	19.4	41.2	39.5	34.1	34.1	3.0	11.7	11.4	11.7
Philadelphia	16.9	69.5	67.0	49.2	58.0	25.1	20.3	20.4	15.6
San Diego	33.3	45.8	50.0	37.5	45.8	20.8	20.8	8.3	11.9
San Francisco	25.6	64.1	66.7	53.7	42.5	7.5	10.9	21.6	16.3
Local Median	32.3	62.5	66.7	50.7	44.9	13.9	20.8	22.7	21.0
Local Range	13.4-52.4	29.6-87.8	19.5-83.9	13.7-81.8	12.1-79.1	0.0-56.2	0.0-58.8	0.0-53.0	4.5-41.9
TERRITORIAL SURVEYS									
American Samoa	73.1	84.6	92.3	100.0	88.5	73.1	50.0	73.1	84.6
Guam	10.0	40.0	20.0	20.0	40.0	0.0	20.0	20.0	7.7
Marshall Islands	45.1	48.4	89.9	86.3	86.3	75.5	64.5	85.9	32.3
Northern Mariana Islands	60.0	100.0	100.0	100.0	80.0	60.0	60.0	80.0	42.9
Palau	28.6	85.7	100.0	57.1	42.9	100.0	57.1	85.7	54.5
Territorial Median	45.1	84.6	92.3	86.3	80.0	73.1	57.1	80.0	42.9
Territorial Range	10.0-73.1	40.0-100.0	20.0-100.0	20.0-100.0	40.0-88.5	0.0-100.0	20.0-64.5	20.0-85.9	7.7-84.6
TRIBAL SURVEYS									
Cherokee Nation	29.0	91.4	90.6	69.8	41.7	16.2	24.0	28.2	35.3
Nez Perce	50.0	100.0	100.0	100.0	50.0	25.0	75.0	25.0	57.1

*A group, committee, or team that offers guidance on the development of policies or coordinates activities on health topics.

†Among schools with school health councils.

TABLE 48a. Percentage of Secondary Schools in Which Students' Families and Community Members Helped Develop or Implement Policies and Programs Related to Specific Health Topics, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Students' families		Community members		Families and community members	
	HIV,* STD,† or teen pregnancy prevention	Asthma	HIV, STD, or teen pregnancy prevention	Asthma	HIV, STD, or teen pregnancy prevention	Asthma
STATE SURVEYS						
Alabama	26.9	18.5	34.6	19.5	23.9	14.8
Alaska	10.9	9.0	20.6	11.8	9.4	6.4
Arizona	12.9	10.2	18.4	11.8	10.6	7.5
Arkansas	20.3	29.3	24.5	29.3	18.4	24.8
California	14.9	11.9	23.7	15.6	12.8	9.0
Colorado	11.3	9.3	20.3	10.2	9.9	6.6
Connecticut	8.2	8.9	28.0	15.1	7.8	7.0
Delaware	17.8	27.1	22.3	19.4	8.2	14.5
Florida	14.5	12.7	26.2	18.2	12.4	9.6
Georgia	14.5	15.3	29.2	17.5	13.4	11.9
Hawaii	10.1	9.8	31.3	22.2	8.8	8.7
Idaho	11.7	11.5	22.1	14.1	9.6	9.5
Indiana	19.5	13.5	29.1	12.9	16.7	9.0
Iowa	11.8	8.6	18.8	11.6	11.4	7.8
Kansas	9.2	9.2	13.1	8.1	8.1	7.1
Kentucky	14.3	13.5	26.5	14.8	13.0	9.3
Louisiana	9.3	8.9	13.1	8.8	6.6	5.9
Maine	7.0	6.6	12.7	9.4	5.5	5.4
Maryland	17.3	16.7	26.8	18.4	14.9	12.3
Massachusetts	10.6	12.4	20.4	16.3	8.9	9.6
Michigan	40.9	15.4	45.2	16.0	37.1	12.5
Minnesota	17.0	15.2	23.1	15.9	15.4	12.4
Mississippi	15.4	16.2	26.4	17.0	14.0	13.2
Missouri	18.4	21.6	23.8	21.9	14.7	18.3
Montana	12.9	10.6	19.3	13.5	12.1	9.7
Nebraska	7.7	13.1	13.7	13.1	4.8	7.2
Nevada	10.6	7.9	22.0	14.2	10.5	6.7
New Hampshire	9.4	9.6	12.2	9.5	7.2	6.2
New Jersey	13.6	18.2	19.2	17.0	10.2	11.9
New Mexico	18.8	16.6	33.5	17.8	16.0	13.8
New York	20.8	18.7	28.3	23.0	17.2	15.2
North Carolina	12.6	16.3	26.7	23.3	11.2	15.4
North Dakota	10.0	11.0	11.7	8.7	6.5	7.6
Ohio	7.4	9.4	15.9	11.8	6.1	6.6
Oklahoma	29.8	25.1	32.3	23.1	25.5	20.7
Oregon	12.1	4.8	20.1	7.4	10.9	2.5
Pennsylvania	13.5	15.1	24.0	20.0	11.7	12.5
Rhode Island	9.4	9.5	20.0	13.9	9.4	7.3
South Carolina	18.6	17.1	31.7	16.9	16.1	13.1
South Dakota	11.6	8.6	12.9	9.9	9.9	7.6
Tennessee	19.1	21.3	30.4	25.0	17.4	17.7
Texas	38.3	26.3	42.6	28.7	36.0	24.2
Utah	13.2	11.5	17.7	14.4	10.5	9.9
Vermont	7.9	12.4	14.5	12.5	7.2	10.0
Virginia	17.7	15.6	31.4	19.9	15.2	11.4
Washington	16.5	11.0	22.3	10.6	13.4	7.2
West Virginia	12.5	18.8	23.7	19.9	9.9	15.3
Wisconsin	14.5	11.0	20.5	13.0	12.2	8.4
Wyoming	10.2	8.0	13.0	7.2	8.5	5.4
State Median	13.2	12.4	22.3	15.1	11.2	9.6
State Range	7.0-40.9	4.8-29.3	11.7-45.2	7.2-29.3	4.8-37.1	2.5-24.8

TABLE 48a. Percentage of Secondary Schools in Which Students’ Families and Community Members Helped Develop or Implement Policies and Programs Related to Specific Health Topics, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 *(continued)*

Site	Students’ families		Community members		Families and community members	
	HIV,* STD,† or teen pregnancy prevention	Asthma	HIV, STD, or teen pregnancy prevention	Asthma	HIV, STD, or teen pregnancy prevention	Asthma
LOCAL SURVEYS						
Albuquerque	21.4	18.8	21.4	11.0	10.7	8.1
Baltimore	10.1	21.1	27.2	29.7	10.0	16.2
Broward County	20.6	24.4	35.9	32.0	17.6	18.2
Charlotte-Mecklenberg County	5.8	21.7	11.7	23.3	3.9	17.6
Chicago	16.0	28.7	32.6	31.7	14.0	23.7
Detroit	28.3	39.5	40.8	44.8	24.9	33.8
District of Columbia	31.6	20.0	43.1	25.8	25.8	17.1
Fresno	12.1	24.7	20.3	20.9	12.1	12.6
Houston	26.4	24.0	33.6	22.6	21.4	17.6
Los Angeles	22.9	22.1	34.6	22.0	18.7	16.2
Memphis	30.1	25.6	42.6	27.5	28.4	22.0
Miami-Dade County	23.1	18.4	31.4	24.9	19.7	13.9
Milwaukee	17.7	19.7	37.5	39.1	13.4	16.7
New York City	28.9	28.2	41.8	33.8	22.3	22.2
Newark	21.8	16.1	21.8	14.4	14.5	8.1
Orange County	24.1	21.7	22.1	15.5	17.7	11.2
Philadelphia	15.3	19.0	29.9	25.8	12.3	15.0
San Diego	0.0	0.0	100.0	98.5	0.0	0.0
San Francisco	31.5	28.5	65.0	42.9	28.5	15.8
Local Median	21.8	21.7	33.6	25.8	17.6	16.2
Local Range	0.0-31.6	0.0-39.5	11.7-100.0	11.0-98.5	0.0-28.5	0.0-33.8
TERRITORIAL SURVEYS						
American Samoa	19.2	7.7	100.0	88.5	19.2	7.7
Guam	7.7	0.0	61.5	30.8	7.7	0.0
Marshall Islands	69.3	7.3	75.8	11.9	60.7	5.8
Northern Mariana Islands	37.5	37.5	75.0	25.0	37.5	25.0
Palau	100.0	20.0	100.0	11.1	100.0	11.1
Territorial Median	37.5	7.7	75.8	25.0	37.5	7.7
Territorial Range	7.7-100.0	0.0-37.5	61.5-100.0	11.1-88.5	7.7-100.0	0.0-25.0
TRIBAL SURVEYS						
Cherokee Nation	33.1	29.5	39.9	32.7	28.5	24.9
Nez Perce	0.0	0.0	42.9	14.3	0.0	0.0

*Human immunodeficiency virus.
 †Sexually transmitted disease.

TABLE 48b. Percentage of Secondary Schools in Which Students' Families and Community Members Helped Develop or Implement Policies and Programs Related to Specific Health Topics, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010

Site	Students' families			Community members			Families and community members			
	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	Tobacco-use prevention, physical activity, and nutrition and healthy eating
STATE SURVEYS										
Alabama	41.3	23.6	30.1	40.2	30.7	33.5	49.0	34.2	37.9	13.6
Alaska	28.9	32.1	36.5	38.9	36.5	43.3	42.2	41.1	47.2	17.0
Arizona	19.7	20.4	24.7	26.2	23.7	30.2	30.6	28.7	34.9	8.9
Arkansas	44.5	51.0	60.6	49.2	47.7	59.1	53.3	55.1	64.7	30.5
California	22.4	24.6	37.7	28.5	28.8	40.3	33.3	33.9	47.5	11.3
Colorado	19.0	29.6	38.8	30.3	27.9	36.7	32.4	34.3	45.0	12.2
Connecticut	12.5	22.8	34.7	32.5	31.9	44.3	33.4	35.6	49.6	7.7
Delaware	35.1	43.2	55.8	32.3	40.0	51.0	46.0	51.1	63.7	14.5
Florida	21.3	30.7	31.2	33.8	34.3	35.9	37.9	40.6	43.5	11.8
Georgia	18.5	22.4	30.4	25.9	28.2	36.6	29.2	31.4	40.6	10.9
Hawaii	16.0	23.0	30.7	34.5	32.4	44.7	37.6	38.3	48.6	12.5
Idaho	22.8	29.4	40.5	30.7	31.0	43.3	35.2	37.8	49.6	11.0
Indiana	28.7	26.2	42.9	40.8	32.4	45.2	43.8	36.0	51.9	15.4
Iowa	27.2	41.1	54.5	35.1	45.1	59.8	38.0	47.5	63.2	18.5
Kansas	22.4	31.2	41.2	26.9	35.1	43.3	30.6	38.9	47.6	14.1
Kentucky	31.1	24.4	30.8	41.1	30.4	41.2	44.8	34.4	45.7	13.2
Louisiana	19.3	17.4	23.3	24.2	19.2	22.9	29.2	25.6	31.0	7.1
Maine	18.1	29.3	39.2	30.6	35.9	44.5	30.6	37.8	48.3	13.3
Maryland	22.3	24.1	28.4	30.2	29.1	35.3	35.1	34.5	39.5	11.8
Massachusetts	16.6	30.7	42.7	27.1	36.2	46.8	29.7	42.3	54.4	10.9
Michigan	25.0	33.9	47.0	33.4	33.1	47.4	36.1	39.7	55.5	15.3
Minnesota	23.6	31.9	48.3	31.6	33.4	50.7	34.6	38.5	55.9	16.5
Mississippi	29.7	37.1	49.5	35.0	40.6	49.9	40.8	47.0	56.6	17.2
Missouri	30.3	39.1	48.1	34.0	40.0	50.7	38.8	46.5	56.8	19.9
Montana	26.8	26.9	35.0	33.0	24.9	34.6	36.8	29.6	40.7	14.2
Nebraska	13.1	20.7	29.5	19.8	24.6	34.6	24.7	29.8	38.3	4.4
Nevada	13.0	16.9	22.4	27.0	26.3	31.3	27.0	27.7	34.3	8.7
New Hampshire	12.6	28.1	44.8	18.8	28.7	47.3	21.5	34.2	52.8	5.6
New Jersey	23.4	27.2	39.9	28.2	31.1	40.9	33.9	36.6	48.1	10.6
New Mexico	28.1	31.2	36.2	38.4	36.3	38.9	42.2	41.0	45.3	17.1
New York	29.3	39.2	55.3	35.5	42.5	54.5	39.6	50.4	63.6	18.0
North Carolina	25.2	23.5	27.6	38.4	31.7	39.1	40.6	35.6	41.1	12.0
North Dakota	28.8	30.8	46.7	34.6	29.4	46.6	39.3	35.7	52.3	12.7
Ohio	15.8	24.9	31.4	23.5	30.0	35.6	26.1	32.6	38.9	8.6
Oklahoma	41.3	41.2	53.1	43.6	38.1	48.0	50.6	44.9	56.1	25.3
Oregon	15.3	17.0	30.2	25.3	23.6	36.6	27.0	26.8	40.5	6.8
Pennsylvania	17.6	34.2	47.0	26.8	38.5	55.3	29.8	44.3	60.5	13.4
Rhode Island	24.2	29.6	53.0	30.2	31.0	57.3	31.4	35.5	64.9	17.5
South Carolina	19.9	30.6	34.6	31.3	35.7	38.7	33.6	38.9	44.4	15.0
South Dakota	30.0	37.0	44.5	28.7	36.7	43.8	34.2	40.8	48.6	17.0
Tennessee	22.8	40.0	40.8	35.4	42.8	43.0	36.4	49.0	48.2	19.0
Texas	37.3	46.1	51.0	41.8	49.0	54.0	44.3	54.4	58.0	28.9
Utah	24.7	26.3	38.4	29.3	29.4	47.0	33.1	33.2	50.3	13.2
Vermont	19.8	26.6	49.3	27.5	31.9	47.4	31.2	37.1	58.2	11.2
Virginia	22.5	29.9	39.2	32.5	38.4	44.6	36.0	44.7	50.0	14.6
Washington	18.8	22.0	31.4	25.9	23.2	33.3	31.3	29.4	38.9	7.4
West Virginia	31.3	27.0	39.6	40.1	34.2	46.1	45.8	38.6	51.5	14.8
Wisconsin	19.1	30.6	47.4	26.3	32.4	50.6	29.4	36.6	54.8	11.6
Wyoming	22.4	23.3	35.5	36.4	24.1	38.4	38.0	29.6	43.2	11.0
State Median	22.8	29.4	39.2	31.6	32.4	43.8	35.1	37.1	48.6	13.2
State Range	12.5-44.5	16.9-51.0	22.4-60.6	18.8-49.2	19.2-49.0	22.9-59.8	21.5-53.3	25.6-55.1	31.0-64.9	4.4-30.5

TABLE 48b. Percentage of Secondary Schools in Which Students’ Families and Community Members Helped Develop or Implement Policies and Programs Related to Specific Health Topics, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2010 (continued)

Site	Students’ families			Community members			Families and community members			
	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	Tobacco-use prevention	Physical activity	Nutrition and healthy eating	Tobacco-use prevention, physical activity, and nutrition and healthy eating
LOCAL SURVEYS										
Albuquerque	16.2	29.5	37.9	16.2	16.2	26.9	24.3	29.5	43.1	8.1
Baltimore	18.6	25.6	33.8	27.6	37.4	40.2	33.8	43.6	46.4	11.2
Broward County	26.4	30.8	34.0	37.3	41.5	46.2	41.8	47.5	50.6	15.7
Charlotte-Mecklenberg County	19.4	24.3	13.7	30.9	27.6	29.4	32.8	34.3	31.3	6.0
Chicago	14.9	37.6	42.5	25.5	40.7	46.6	27.5	48.5	53.5	9.0
Detroit	35.6	44.4	47.9	48.8	58.4	58.1	53.3	62.9	64.2	23.0
District of Columbia	20.1	31.5	43.0	28.7	34.3	40.1	34.4	40.0	45.8	11.4
Fresno	20.3	24.7	28.6	24.7	25.3	33.5	28.6	33.0	41.2	12.6
Houston	27.7	35.3	38.6	30.1	41.7	43.5	35.2	44.2	48.5	22.7
Los Angeles	29.4	24.8	30.9	35.2	33.6	36.8	42.8	37.9	41.9	18.0
Memphis	30.1	37.2	41.1	39.0	44.1	45.2	42.4	49.1	51.9	23.5
Miami-Dade County	25.0	40.0	42.9	34.0	42.4	47.9	38.4	51.0	54.2	16.4
Milwaukee	22.4	29.8	29.5	41.3	40.9	50.4	46.4	48.3	56.3	9.1
New York City	25.4	35.8	45.9	31.8	41.6	45.2	38.6	50.9	57.4	16.1
Newark	15.2	22.8	48.9	17.5	27.6	43.0	24.4	33.6	55.3	6.5
Orange County	26.2	34.6	32.6	24.0	30.5	39.3	32.8	39.0	47.8	13.2
Philadelphia	14.2	28.3	37.2	27.6	34.4	51.8	29.7	43.0	58.1	10.0
San Diego	0.0	0.0	0.0	100.0	100.0	100.0	100.0	100.0	100.0	0.0
San Francisco	34.8	34.5	59.2	65.2	57.6	62.2	71.5	62.0	77.7	19.0
Local Median	22.4	30.8	37.9	30.9	40.7	45.2	35.2	44.2	51.9	12.6
Local Range	0.0-35.6	0.0-44.4	0.0-59.2	16.2-100.0	16.2-100.0	26.9-100.0	24.3-100.0	29.5-100.0	31.3-100.0	0.0-23.5
TERRITORIAL SURVEYS										
American Samoa	23.1	19.2	19.2	100.0	100.0	100.0	100.0	100.0	100.0	19.2
Guam	7.7	7.7	0.0	69.2	61.5	61.5	69.2	61.5	61.5	0.0
Marshall Islands	75.1	85.5	75.2	86.7	85.2	85.2	88.4	89.9	85.4	65.8
Northern Mariana Islands	62.5	62.5	75.0	75.0	75.0	87.5	75.0	75.0	87.5	50.0
Palau	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Territorial Median	62.5	62.5	75.0	86.7	85.2	87.5	88.4	89.9	87.5	50.0
Territorial Range	7.7-100.0	7.7-100.0	0.0-100.0	69.2-100.0	61.5-100.0	61.5-100.0	69.2-100.0	61.5-100.0	61.5-100.0	0.0-100.0
TRIBAL SURVEYS										
Cherokee Nation	38.8	45.5	53.8	49.1	45.1	53.7	51.9	52.8	61.0	27.5
Nez Perce	14.3	28.6	42.9	42.9	28.6	71.4	42.9	28.6	71.4	14.3

Suggested Citation

Brener ND, Demissie Z, Foti K, McManus T, Shanklin SL, Hawkins J, Kann L.
School Health Profiles 2010: Characteristics of Health Programs Among Secondary Schools. Atlanta: Centers for Disease Control and Prevention; 2011.

Ordering Information

For additional information about school health or to request free copies of this report, send an e-mail to HealthyYouth@cdc.gov, call 1-800-CDC-INFO (232-4636), or visit our Web site at <http://www.cdc.gov/schoolhealthprofiles>.