

Funding Opportunity Announcement (FOA)

PS17-1704: Comprehensive High-Impact HIV Prevention Projects for Young Men of Color Who Have Sex with Men and Young Transgender Persons of Color

Attachment L: High-Risk and Substantial Risk Definitions for HIV infection

Table 1 presents the screening criteria for identifying persons at high risk for acquiring HIV, and Table 2 provides the screening criteria for identifying persons in the subset designated by the PrEP guidelines as being at substantial risk, for whom PrEP may be indicated.

Table 1. Screening Criteria for Identifying HIV-negative Persons Who May Be At High Risk for HIV		
Use these criteria for MEN WHO HAVE SEX WITH MEN (MSM)	Use these criteria for HETEROSEXUALLY ACTIVE MEN AND WOMEN	Use these criteria for TRANSGENDER PERSONS
<p style="text-align: center;"><u>BOTH OF THESE:</u></p> <ul style="list-style-type: none"> • Adult or Adolescent male • Without acute or established HIV infection <p>Note: If also has sex with women, evaluate by criteria in Column 2, as well</p> <p style="text-align: center;"><u>AND</u></p> <p style="text-align: center;"><u>ONE OR MORE OF THE FOLLOWING:</u></p> <p>IN THE PAST 12 MONTHS:</p> <ul style="list-style-type: none"> • Had receptive or insertive anal sex with an HIV-positive male partner • Had receptive or insertive anal sex without a condom with a male partner who was not known to be HIV-negative • Had a bacterial sexually transmitted infection (STI) diagnosed or reported 	<p style="text-align: center;"><u>BOTH OF THESE:</u></p> <ul style="list-style-type: none"> • Adult or Adolescent person • Without acute or established HIV infection <p>Note: If person is a man who has sex with both women and men, evaluate by criteria in Column 1, as well</p> <p style="text-align: center;"><u>AND</u></p> <p style="text-align: center;"><u>ONE OR MORE OF THE FOLLOWING:</u></p> <p>IN THE PAST 12 MONTHS:</p> <ul style="list-style-type: none"> • Had vaginal or anal sex with an HIV-positive partner • Had vaginal or anal sex without a condom with at least one partner of unknown HIV status whose behavioral risks are not known or who: <ul style="list-style-type: none"> ○ Is an MSM, or ○ Is a person who injects drugs (PWID), or ○ Had a recent bacterial STI, or ○ Has an HIV-positive partner, or ○ Has a partner who is a PWID, or ○ Resides in or is socially connected to an area, population, or social, sexual, or drug using network presumed to have relatively high HIV prevalence, based on locally available information • Had a bacterial STI diagnosed or reported 	<p style="text-align: center;"><u>BOTH OF THESE:</u></p> <ul style="list-style-type: none"> • Adult or Adolescent person • Without acute or established HIV infection <p style="text-align: center;"><u>AND</u></p> <p style="text-align: center;"><u>ONE OR MORE OF THE FOLLOWING:</u></p> <p>IN THE PAST 12 MONTHS:</p> <ul style="list-style-type: none"> • Had vaginal or anal sex with an HIV-positive partner • Had vaginal or anal sex without a condom with at least one partner of unknown HIV status whose behavioral risks are not known or who: <ul style="list-style-type: none"> ○ Is an MSM, or ○ Is a PWID, or ○ Had a recent bacterial STI, or ○ Has an HIV-positive partner, or ○ Has a partner who is a PWID, or ○ Resides in or is socially connected to an area, population, or social, sexual, or drug using network presumed to have relatively high HIV prevalence, based on locally available information • Had a bacterial STI diagnosed or reported

Table 1. Screening Criteria for Identifying HIV-negative Persons Who May Be At High Risk for HIV

Use these criteria for
PERSONS WHO INJECT DRUGS (PWID)

ALL OF THESE:

- Adult or Adolescent person
- Without acute or established HIV infection
- **IN THE PAST 12 MONTHS**, any injection of drugs not prescribed to the person by a clinician for medical purposes

AND

- **IN THE PAST 12 MONTHS**, any sharing of injection or drug preparation equipment to inject drugs not prescribed to the person by a clinician for medical purposes

Note: Evaluate risk for sexual acquisition by criteria in Columns 1, 2, and 3, as well

Table 2. Screening Criteria for Identifying HIV-negative Persons Who May Be At Substantial Risk for HIV and Recommended Indications For PrEP Use¹

Use these criteria for MEN WHO HAVE SEX WITH MEN	Use these criteria for HETEROSEXUALLY ACTIVE MEN AND WOMEN	Use these criteria for PERSONS WHO INJECT DRUGS
<p><u>ALL OF THESE:</u></p> <ul style="list-style-type: none"> • Adult man² • Without acute or established HIV infection • Any male sex partners IN THE PAST 6 MONTHS (if also has sex with women, see Column 2) • Not in a monogamous partnership with a recently tested, HIV-negative man <p style="text-align: center;"><u>AND</u></p> <p><u>AT LEAST ONE OF THE FOLLOWING:</u></p> <ul style="list-style-type: none"> • Any anal sex without condoms (receptive or insertive) IN THE PAST 6 MONTHS • Any STI diagnosed or reported IN THE PAST 6 MONTHS • Is in an ongoing sexual relationship with an HIV-positive male partner 	<p><u>ALL OF THESE:</u></p> <ul style="list-style-type: none"> • Adult person² • Without acute or established HIV infection • Any sex with opposite sex partners IN THE PAST 6 MONTHS • Not in a monogamous partnership with a recently tested, HIV-negative partner <p style="text-align: center;"><u>AND</u></p> <p><u>AT LEAST ONE OF THE FOLLOWING:</u></p> <ul style="list-style-type: none"> • Is a man who has sex with both women and men (behaviorally bisexual) [also evaluate indications for PrEP use by Column 1 criteria] • Infrequently uses condoms during sex with 1 or more partners of unknown HIV status who are known to be at substantial risk of HIV infection (IDU or bisexual male partner) • Is in an ongoing sexual relationship with an HIV-positive partner 	<p><u>ALL OF THESE:</u></p> <ul style="list-style-type: none"> • Adult person² • Without acute or established HIV infection • Any injection of drugs not prescribed by a clinician IN THE PAST 6 MONTHS <p style="text-align: center;"><u>AND</u></p> <p><u>AT LEAST ONE OF THE FOLLOWING:</u></p> <ul style="list-style-type: none"> • Any sharing of injection or drug preparation equipment IN THE PAST 6 MONTHS • Been in a methadone, buprenorphine, or suboxone treatment program IN THE PAST 6 MONTHS • Risk of sexual acquisition (also evaluate by criteria in Column 1 or 2)