




# HIV Surveillance – Adolescents and Young Adults


# Diagnoses of HIV Infection among Adolescents and Young Adults Aged 13–24 years, by Race/Ethnicity, 2010–2016—United States and 6 Dependent Areas



<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Adolescents and Young Adults Aged 13–24 Years, by Transmission Category, 2010–2016—United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises less than 1% of cases.

<sup>a</sup> Heterosexual contact is with a person known to have, or to be at high risk for, HIV infection.



# Adolescents and Young Adults Aged 13–24 Years Living with Diagnosed HIV Infection, by Sex and Race/Ethnicity, Year-end 2016—United States and 6 Dependent Areas




<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Adolescents and Young Adults Aged 13–24 Years Living with Diagnosed HIV Infection by Sex and Transmission Category, Year-end 2016—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises 1% or less of cases.


<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.



# Rates of Adolescents Aged 13–19 Years Living with Diagnosed HIV Infection

## Year-end 2016—United States and 6 Dependent Areas




# Rates of Young Adults Aged 20–24 Years Living with Diagnosed HIV Infection

## Year-end 2016—United States and 6 Dependent Areas

N = 30,160


Total Rate = 133.1




Note. Data are based on address of residence as of December 31, 2016 (i.e., most recent known address).



# Stage 3 (AIDS) Classifications among Adolescents Aged 13–19 Years, by Sex and Year of Classification, 1985–2016—United States and 6 Dependent Areas




# Stage 3 (AIDS) Classifications Among Young Adults Aged 20–24 Years, by Sex and Year of Classification, 1985–2016—United States and 6 Dependent Areas



# Rates of Adolescents Aged 13–19 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2016—United States and 6 Dependent Areas

N = 951 Total Rate = 3.2




Note. Data are based on address of residence as of December 31, 2016 (i.e., most recent known address).




# Rates of Young Adults Aged 20–24 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2016—United States and 6 Dependent Areas

N = 6,046

Total Rate = 26.7




American Samoa	0.0
Guam	7.4
Northern Mariana Islands	0.0
Puerto Rico	31.6
Republic of Palau	0.0
U.S. Virgin Islands	32.3



Note. Data are based on address of residence as of December 31, 2016 (i.e., most recent known address).



# Diagnoses of HIV Infection among Persons Aged 13 Years and Older, by Sex and and Age Group, 2017—United States and 6 Dependent Areas



Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay.



# Diagnoses of HIV Infection Among Male Adolescents and Young Adults, by Age Group and Transmission Category, 2017—United States and 6 Dependent Areas

Transmission category	13–19 years		20–24 years	
	No.	%	No.	%
Male-to-male sexual contact	1,400	94.2	5,192	92.1
Injection drug use (IDU)	13	0.9	89	1.6
Male-to-male sexual contact and IDU	28	1.9	180	3.2
Heterosexual contact <sup>a</sup>	37	2.5	173	3.1
Other <sup>b</sup>	8	0.5	3	0.1
<b>Total<sup>c</sup></b>	<b>1,487</b>	<b>100</b>	<b>5,638</b>	<b>100</b>

*Note.* Data for the year 2017 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>c</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total



# Diagnoses of HIV Infection Among Female Adolescents and Young Adults, by Age Group and Transmission Category, 2017—United States and 6 Dependent Areas

Transmission category	13–19 years		20–24 years	
	No.	%	No.	%
Injection drug use	22	8.5	90	11.5
Heterosexual contact <sup>a</sup>	221	84.7	677	87.0
Other <sup>b</sup>	18	6.8	11	1.5
<b>Total<sup>c</sup></b>	<b>261</b>	<b>100</b>	<b>778</b>	<b>100</b>

*Note.* Data for the year 2017 are considered preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.


<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>c</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



# Diagnoses of HIV Infection and Population among Adolescents Aged 13–19 Years, by Race/Ethnicity, 2017—United States




Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection and Population among Young Adults Aged 20–24 Years, by Race/Ethnicity 2017—United States



Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay.


<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Diagnoses of HIV Infection among Adolescents Aged 13–19 Years 2017—United States and 6 Dependent Areas

**N = 1,748**

**Total Rate = 5.9**




*Note.* Data for the year 2017 are considered preliminary and based on 6 months reporting delay.

# Rates of Diagnoses of HIV Infection among Young Adults Aged 20–24 Years 2017—United States and 6 Dependent Areas

N = 6,416


Total Rate = 28.7



Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay.



# Stage 3 (AIDS) Classifications among Persons Aged 13 Years and Older with Diagnosed HIV Infection, by Race/Ethnicity and Age Group, 2017 United States and 6 Dependent Areas



Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay.


<sup>a</sup> Includes Asian/Pacific Islander legacy cases

<sup>b</sup> Hispanics/Latinos can be of any race.



# Rates of Diagnosed HIV Infection Classified as Stage 3 (AIDS) among Adolescents Aged 13–19 Years, 2017—United States and 6 Dependent Areas

N = 210 Total Rate = 0.7




Note. Data for the year 2017 are preliminary and based on 6 months reporting delay.



# Rates of Diagnosed HIV Infection Classified as Stage 3 (AIDS) among Young Adults Aged 20–24 Years, 2017—United States and 6 Dependent Areas

N = 1,158

Total Rate = 5.2



Note. Data for the year 2017 are considered preliminary and based on 6 months reporting delay.

