



# Epidemiology of HIV Infection through 2015

## Diagnoses of HIV Infection among Adults and Adolescents, by Sex 2010–2014—United States and 6 Dependent Areas




*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

# Rates of Diagnoses of HIV Infection among Adults and Adolescents by Age at Diagnosis, 2010–2014—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.


# Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

<sup>a</sup> Hispanics/Latinos can be of any race.

# Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. "Other" transmission category not displayed as it comprises less than 1% of cases.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

# Deaths of Persons with Diagnosed HIV Infection by Race/Ethnicity, 2014—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	55	2.3
Asian <sup>a</sup>	55	0.3
Black/African American	6,888	17.4
Hispanic/Latino <sup>b</sup>	2,525	4.6
Native Hawaiian/other Pacific Islander	7	1.3
White	4,722	2.4
Multiple races	867	13.6
<b>Total</b>	<b>15,119</b>	<b>4.7</b>


*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause. Rates are per 100,000 population.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Adults and Adolescents Living with Diagnosed HIV Infection, by Sex and Race/Ethnicity, Year-end 2014—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Adults and Adolescents Living with Diagnosed HIV Infection, by Sex and Transmission Category, Year-end 2014—United States and 6 Dependent Areas



*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises 1% or less of cases.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.






# Rates of Adults and Adolescents Living with Diagnosed HIV Infection

## Year-end 2014—United States and 6 Dependent Areas

N = 970,319


Total Rate = 360.0



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data are based on address of residence as of December 31, 2014 (i.e., most recent known address).

# Rates of Children Aged <13 Years Living with Diagnosed HIV Infection


## Year-end 2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data are based on address of residence as of December 31, 2014 (i.e., most recent known address).

# Stage 3 (AIDS) Classifications and Deaths of Persons with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), among Adults and Adolescents, 1985–2014


## US and 6 Dependent Areas



Note. Deaths of persons with HIV infection, stage 3 (AIDS) may be due to any cause.



# Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity and Year of Classification, 1985–2014—United States and 6 Dependent Areas




<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Includes Asian/Pacific Islander legacy cases.



# Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity and Year of Classification, 1985–2014—United States and 6 Dependent Areas




<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Includes Asian/Pacific Islander legacy cases.

# Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Transmission Category and Year of Classification, 1985–2014

## United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

# Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Transmission Category and Year of Classification 1985–2014—United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




## Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Sex, 1993–2014—United States and 6 Dependent Areas





# Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, 1993–2014—United States and 6 Dependent Areas




<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Includes Asian/Pacific Islander legacy cases.

# Rates of Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2014—United States and 6 Dependent Areas

N = 530,620

Total Rate = 196.9




Note. Data are based on address of residence as of December 31, 2014 (i.e., most recent known address).

# Rates of Children Aged <13 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2014—United States and 6 Dependent Areas

N = 311

Total Rate = 0.6



Note. Data are based on address of residence as of December 31, 2014 (i.e., most recent known address).

## Diagnoses of HIV Infection among Male Adults and Adolescents by Race/Ethnicity, 2015—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	152	16.2
Asian	816	11.7
Black/African American	13,070	84.8
Hispanic/Latino <sup>a</sup>	8,147	37.4
Native Hawaiian/other Pacific Islander	71	31.2
White	9,060	10.8
Multiple races	675	32.9
<b>Total</b>	<b>31,991</b>	<b>24.4</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Rates are per 100,000 population.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Female Adults and Adolescents by Race/Ethnicity, 2015—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	55	5.6
Asian	132	1.7
Black/African American	4,524	26.2
Hispanic/Latino <sup>a</sup>	1,131	5.3
Native Hawaiian/other Pacific Islander	8	3.5
White	1,431	1.6
Multiple races	121	5.5
<b>Total</b>	<b>7,402</b>	<b>5.4</b>


*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Rates are per 100,000 population.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Adults and Adolescents by Age at Diagnosis, 2015—United States


N = 39,393



*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay.



# Diagnoses of HIV Infection among Adults and Adolescents by Sex and Race/Ethnicity, 2015—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Diagnoses of HIV Infection among Adults and Adolescents by Sex and Race/Ethnicity, 2015—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay.


<sup>a</sup> Hispanics/Latinos can be of any race.





# Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2015—United States and 6 Dependent Areas

N = 39,920




*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises less than 1% of cases.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



# Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Transmission Category, 2015—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. "Other" transmission category not displayed as it comprises less than 1% of cases.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



# Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category 2015—United States and 6 Dependent Areas

Transmission category	No.	%
Male-to-male sexual contact	26,646	66.7
Injection drug use (IDU)	2,460	6.2
Male-to-male sexual contact and IDU	1,215	3.0
Heterosexual contact <sup>a</sup>	9,515	23.8
Other <sup>b</sup>	84	0.2
<b>Total<sup>c</sup></b>	<b>39,920</b>	<b>100</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>c</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



# Diagnosed HIV Infections Attributed to Male-to-Male Sexual Contact by Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	120	0.5
Asian	729	2.7
Black/African American	10,318	38.7
Hispanic/Latino <sup>a</sup>	7,271	27.3
Native Hawaiian/other Pacific Islander	64	0.2
White	7,572	28.4
Multiple races	570	2.1
<b>Total<sup>b</sup></b>	<b>26,646</b>	<b>100</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



# Diagnosed HIV Infections Attributed to Injection Drug Use, by Sex and Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	Males		Females		Total	
	No.	%	No.	%	No.	%
American Indian/Alaska Native	10	0.7	15	1.5	25	1.0
Asian	20	1.4	6	0.6	26	1.1
Black/African American	539	36.6	363	36.8	902	36.7
Hispanic/Latino <sup>a</sup>	388	26.3	121	12.3	509	20.7
Native Hawaiian/other Pacific Islander	3	0.2	2	0.2	5	0.2
White	493	33.4	458	46.5	951	38.7
Multiple races	22	1.5	20	2.0	42	1.7
<b>Total<sup>b</sup></b>	<b>1,474</b>	<b>100</b>	<b>986</b>	<b>100</b>	<b>2,460</b>	<b>100</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



# Diagnosed HIV Infections Attributed to Male-to-Male Sexual Contact and Injection Drug Use, by Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	11	0.9
Asian	17	1.4
Black/African American	262	21.6
Hispanic/Latino <sup>a</sup>	276	22.7
Native Hawaiian/other Pacific Islander	1	0.1
White	606	49.9
Multiple races	42	3.5
<b>Total<sup>b</sup></b>	<b>1,215</b>	<b>100</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



# Diagnosed HIV Infections Attributed to Heterosexual Contact<sup>a</sup>, by Sex and Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	Males		Females		Total	
	No.	%	No.	%	No.	%
American Indian/Alaska Native	11	0.4	40	0.6	51	0.5
Asian	53	1.7	125	1.9	178	1.9
Black/African American	1,926	63.5	4,146	64.0	6,072	63.8
Hispanic/Latino <sup>b</sup>	619	20.4	1,096	16.9	1,715	18.0
Native Hawaiian/other Pacific Islander	5	0.2	6	0.1	11	0.1
White	381	12.6	968	14.9	1,349	14.2
Multiple races	40	1.3	100	1.5	140	1.5
<b>Total<sup>c</sup></b>	<b>3,034</b>	<b>100</b>	<b>6,481</b>	<b>100</b>	<b>9,515</b>	<b>100</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Hispanics/Latinos can be of any race.


<sup>c</sup> Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



# Rates of Diagnoses of HIV Infection among Adults and Adolescents 2015—United States and 6 Dependent Areas

N = 39,920

Total Rate = 14.7





## Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity, 2015—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	96	5.0
Asian <sup>a</sup>	323	2.2
Black/African American	8,683	26.6
Hispanic/Latino <sup>b</sup>	3,865	8.9
Native Hawaiian/other Pacific Islander	22	4.8
White	4,666	2.7
Multiple races	619	14.6
<b>Total</b>	<b>18,274</b>	<b>6.8</b>


Note. Data for the year 2015 are preliminary and based on 6 months reporting delay. Rates are per 100,000 population.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Sex and Transmission Category, 2015—United States and 6 Dependent Areas



*Note.* Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Rates of Diagnosed HIV Infection Classified as Stage 3 (AIDS) 2015—United States and 6 Dependent Areas

N = 18,538

Total Rate = 5.7



Note. Data for the year 2015 are preliminary and based on 6 months reporting delay.