


Pediatric HIV Surveillance

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection among Children Aged <13 Years, by Age at Diagnosis, 2010–2014 United States and 6 Dependent Areas N= 1,059




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of HIV Infection among Children Aged <13 Years, by Race/Ethnicity, 2010–2014—United States and 6 Dependent Areas

N= 1,059



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.




Diagnoses of HIV Infection and Population in Children Aged <13 Years, by Race/Ethnicity, 2014—United States

Diagnoses of HIV infection
N = 174



Population, United States
N = 52,666,129



■ American Indian/Alaska Native
■ Asian
■ Black/African American

■ Hispanic/Latino^a
■ Native Hawaiian/Other Pacific Islander
■ White


■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^bHispanics/Latinos can be of any race




Stage 3 (AIDS) Classifications among Persons with Perinatally Acquired HIV Infection, 1985–2014—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Stage 3 (AIDS) Classifications among Children Aged <13 Years, by Race/Ethnicity, 1985–2014—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.
^a Hispanics/Latinos can be of any race.



Persons Living with Diagnosed Perinatally Acquired HIV Infection, Year-end 2013—United States and 6 Dependent Areas

N = 11,435




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Persons Living with Diagnosed Perinatally Acquired HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2013 United States and 6 Dependent Areas

N = 5,490




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Age Distribution of Persons Living with Diagnosed Perinatally Acquired HIV Infection, Year-end 2013— United States and 6 dependent areas

N = 11,435




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting



Rates of Females Aged 15–44 Years Living with HIV Infection, by Area of Residence, 2013

United States and Puerto Rico

N = 101,046 Total rate = 158.2




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of Perinatally Acquired HIV Infection among Children Born During 2012—United States and 6 Dependent Areas

N = 120




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of Perinatally Acquired HIV Infection among Children Born During 2008–2012, by Area of Residence, United States and Puerto Rico

N=1,003




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of Perinatally Acquired HIV infection among Children Born in the United States and Puerto Rico During 2008-2012, by Area of Residence, United States and Puerto Rico

N=672




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Time of Maternal HIV Testing among Children with Diagnosed Perinatally Acquired HIV infection and Children Exposed to HIV, Birth Years 2008-2012—United States and Puerto Rico


Infected Infants

N = 582



Exposed but not infected

N = 13,334



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have not been statistically adjusted.

Note: Exposure data from 47 areas.

Time of Antiretroviral (ARV) Administration among Pregnant Women with HIV or Perinatally Exposed Infants Birth Years 2008–2012—United States and Puerto Rico

Time of ARV Administration	Year of birth									
	2008		2009		2010		2011		2012	
	No.	%	No.	%	No.	%	No.	%	No.	%
During Pregnancy (DP) only	47	1.5	24	0.8	24	0.9	15	0.6	22	0.9
During Labor and Delivery (L&D) only	15	0.5	14	0.5	12	0.4	15	0.6	14	0.6
Infant received ARV after birth (Infant ARV) only	217	7.0	171	5.7	176	6.3	151	5.7	167	6.9
DP and L&D	79	2.6	99	3.3	69	2.5	48	1.8	37	1.5
DP and Infant ARV	187	6.1	185	6.2	175	6.3	168	6.4	163	6.8
L&D and Infant ARV	229	7.4	219	7.3	176	6.3	188	7.1	154	6.4
DP and L&D and Infant ARV	2,004	65.0	2,071	69.4	1,951	69.9	1,866	70.6	1,640	68.0
No known ARVs	307	10.0	202	6.8	208	7.5	192	7.3	215	8.9
Total	3,085	100	2,985	100	2,791	100	2,643	100	2,412	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have not been statistically adjusted. Exposure data from 47 areas.



HIV Diagnostic Outcomes among Infants, by Time of Antiretroviral (ARV) Administration Birth Years 2008–2012—United States and Puerto Rico

Time of ARV Administration	Infected		Exposed but not infected		Total
	No.	% of	No.	% of	No.
		Row Total		Row Total	
During Pregnancy (DP) only	6	4.5	126	95.5	132
During Labor and Delivery (L&D) only	5	7.1	65	92.9	70
Infant received ARV after birth (Infant ARV) only	85	9.6	797	90.4	882
DP and L&D	7	2.1	325	97.9	332
DP and Infant ARV	27	3.1	851	96.9	878
L&D and Infant ARV	50	5.2	916	94.8	966
DP and L&D and Infant ARV	123	1.3	9,409	98.7	9,532
No known ARVs	279	24.8	845	75.2	1,124
Total	582	4.2	13,334	95.8	13,916

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have not been statistically adjusted. Exposure data from 47 areas.

