




HIV Surveillance – Men Who Have Sex with Men (MSM) through 2016

Diagnoses of HIV Infection among Male Adults and Adolescents, by Transmission Category, 2010–2015—United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises less than 1% of cases.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.




Diagnoses of HIV Infection among Men Who Have Sex with Men, by Age at Diagnosis, 2010–2015—United States and 6 Dependent Areas



Note: Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2010–2015—United States and 6 Dependent Areas




Note: Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Young (Aged 13–24 Years) Men Who Have Sex with Men, by Race/Ethnicity, 2010–2015—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adults and Adolescents, 2010–2015— United States and 6 Dependent Areas

Of the 249,812 diagnoses of HIV infection among adults and adolescents during 2010 through 2015:

- 80% of diagnosed HIV infections in males were attributed to male-to-male sexual contact
- 24% of diagnosed HIV infections in males were among those aged 13–24 years
 - 91% of diagnosed HIV infections in males aged 13–24 years were attributed to male-to-male sexual contact

In 2015, 82% of HIV infections diagnosed among male adults and adolescents were attributed to male-to-male sexual contact



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

Men Who Have Sex with Men Living with Diagnosed HIV Infection, by Race/Ethnicity, Year-end 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	1,458	0.3
Asian ^a	8,732	1.6
Black/African American	162,264	30.6
Hispanic/Latino ^b	125,101	23.6
Native Hawaiian/other Pacific Islander	646	0.1
White	212,190	40.0
Multiple races	20,102	3.8
Total^{c, d}	530,970	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^d Includes persons whose race/ethnicity is unknown.



Deaths among Males with Diagnosed HIV Infection Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	20	0.3
Asian ^a	53	0.8
Black/African American	2,037	31.2
Hispanic/Latino ^b	1,120	17.2
Native Hawaiian/other Pacific Islander	5	0.1
White	2,951	45.2
Multiple races	342	5.2
Total^{c, d}	6,531	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Deaths of persons with a diagnosed HIV infection may be due to any cause.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^d Includes persons whose race/ethnicity is unknown.




Stage 3 (AIDS) Classifications among Men who Have Sex with Men, 1985–2015— United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Race/Ethnicity and Year of Diagnosis, 1985–2015—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications and Deaths among Men Who Have Sex with Men with HIV Infection Ever Classified as Stage 3 (AIDS), 1985–2015

United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Deaths of persons with diagnosed HIV infection may be due to any cause.



Men Who Have Sex with Men Living with HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	693	0.3
Asian ^a	4,042	1.5
Black/African American	77,217	29.0
Hispanic/Latino ^b	64,607	27.3
Native Hawaiian/other Pacific Islander	330	0.1
White	108,234	40.7
Multiple races	10,827	4.1
Total^{c, d}	265,977	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^d Includes persons whose race/ethnicity is unknown.



Diagnoses of HIV Infection among Male Adults and Adolescents, by Transmission Category 2016—United States and 6 Dependent Areas

Transmission category	No.	%
Male-to-male sexual contact	26,844	82.4
Injection drug use (IDU)	1,333	4.1
Male-to-male sexual contact and IDU	1,210	3.7
Heterosexual contact ^a	3,138	9.6
Other ^b	38	0.1
Total^c	32,563	100

Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Age Group and Race/Ethnicity, 2016—United States and 6 Dependent Areas



Note. Data for the year 2016 are preliminary and based on 6 months reporting delay.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2016—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	152	0.6
Asian	740	2.8
Black/African American	10,226	38.1
Hispanic/Latino ^a	7,689	28.6
Native Hawaiian/other Pacific Islander	35	0.1
White	7,392	27.5
Multiple races	610	2.3
Total^b	26,844	100

Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use


^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2016—United States and 6 Dependent Areas

N = 26,844



Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Young (Aged 13–24 Years) Men Who Have Sex with Men, by Race/Ethnicity, 2016—United States and 6 Dependent Areas

N = 6,916




Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Region of Residence and Race/Ethnicity, 2016—United States and 6 Dependent Areas



Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Numbers less than 12, and trends based on these numbers, should be interpreted with caution.

^a Hispanics/Latinos can be of any race



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Race/Ethnicity, 2016—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	57	0.6
Asian ^a	232	2.3
Black/African American	3,922	38.6
Hispanic/Latino ^b	2,692	26.5
Native Hawaiian/other Pacific Islander	11	0.1
White	2,910	28.7
Multiple races	333	3.3
Total^c	10,157	100

Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Region of Residence and Race/Ethnicity, 2016—United States and 6 Dependent Areas



Note. Data for the year 2016 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race

