




HIV Surveillance – Men Who Have Sex with Men (MSM) through 2017

Diagnoses of HIV Infection among Male Adults and Adolescents, by Transmission Category, 2010–2016—United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises less than 1% of cases.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.




Diagnoses of HIV Infection among Men Who Have Sex with Men by Age at Diagnosis, 2010–2016—United States and 6 Dependent Areas



Note: Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity 2010–2016—United States and 6 Dependent Areas




Note: Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Young (Aged 13–24 Years) Men Who Have Sex with Men, by Race/Ethnicity, 2010–2016—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adults and Adolescents 2010–2016—United States and 6 Dependent Areas

Of the 289,416 diagnoses of HIV infection among adults and adolescents during 2010 through 2016:

- 80% of diagnosed HIV infections in males were attributed to male-to-male sexual contact
- 24% of diagnosed HIV infections in males were among those aged 13–24 years
 - 91% of diagnosed HIV infections in males aged 13–24 years were attributed to male-to-male sexual contact

In 2016, 82% of HIV infections diagnosed among male adults and adolescents were attributed to male-to-male sexual contact



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

Men Who Have Sex with Men Living with Diagnosed HIV Infection, by Race/Ethnicity Year-end 2016—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	1,511	0.3
Asian ^a	9,253	1.7
Black/African American	168,460	30.8
Hispanic/Latino ^b	128,628	23.5
Native Hawaiian/other Pacific Islander	620	0.1
White	214,866	39.3
Multiple races	22,962	4.2
Total^{c, d}	546,730	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^d Includes persons whose race/ethnicity is unknown.



Deaths among Males with Diagnosed HIV Infection Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 2016—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	21	0.3
Asian ^a	58	0.9
Black/African American	2,004	30.5
Hispanic/Latino ^b	1,077	16.4
Native Hawaiian/other Pacific Islander	9	0.1
White	3,031	46.1
Multiple races	378	5.7
Total^{c, d}	6,578	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Deaths of persons with a diagnosed HIV infection may be due to any cause.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^d Includes persons whose race/ethnicity is unknown.




Stage 3 (AIDS) Classifications among Men who Have Sex with Men 1985–2016—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Race/Ethnicity and Year of Diagnosis, 1985–2016—United States and 6 Dependent Areas




Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications and Deaths among Men Who Have Sex with Men with HIV Infection Ever Classified as Stage 3 (AIDS) 1985–2016—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Deaths of persons with diagnosed HIV infection may be due to any cause.



Men Who Have Sex with Men Living with HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2016—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	694	0.3
Asian ^a	4,138	1.5
Black/African American	78,889	29.3
Hispanic/Latino ^b	64,627	24.0
Native Hawaiian/other Pacific Islander	304	0.1
White	108,552	40.3
Multiple races	12,233	4.5
Total^{c, d}	269,461	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^d Includes persons whose race/ethnicity is unknown.



Diagnoses of HIV Infection among Male Adults and Adolescents, by Transmission Category, 2017—United States and 6 Dependent Areas

Transmission category	No.	%
Male-to-male sexual contact	25,748	82.4
Injection drug use (IDU)	1,373	4.4
Male-to-male sexual contact and IDU	1,252	4.0
Heterosexual contact ^a	2,829	9.1
Other ^b	37	0.1
Total^c	31,239	100

Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Age Group and Race/Ethnicity, 2017—United States and 6 Dependent Areas



Note. Data for the year 2017 are preliminary and based on 6 months reporting delay.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity 2017—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	125	0.5
Asian	750	2.9
Black/African American	9,807	38.1
Hispanic/Latino ^a	7,436	28.9
Native Hawaiian/other Pacific Islander	39	0.2
White	6,982	27.1
Multiple races	609	2.4
Total^b	25,748	100

Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use


^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2017—United States and 6 Dependent Areas

N = 25,748



Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Young (Aged 13–24 Years) Men Who Have Sex with Men, by Race/Ethnicity, 2017—United States and 6 Dependent Areas

N = 6,592




Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Region of Residence and Race/Ethnicity, 2017—United States and 6 Dependent Areas



Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

Numbers less than 12, and trends based on these numbers, should be interpreted with caution.

^a Hispanics/Latinos can be of any race



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Race/Ethnicity, 2017—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	30	0.3
Asian ^a	240	2.5
Black/African American	3,829	39.1
Hispanic/Latino ^b	2,564	26.2
Native Hawaiian/other Pacific Islander	18	0.2
White	2,757	28.2
Multiple races	343	3.5
Total^c	9,781	100

Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Region of Residence and Race/Ethnicity, 2017—United States and 6 Dependent Areas



Note. Data for the year 2017 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race

