




HIV Surveillance – Men Who Have Sex with Men (MSM) through 2015

Diagnoses of HIV Infection among Male Adults and Adolescents, by Transmission Category, 2010–2014—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. “Other” transmission category not displayed as it comprises less than 1% of cases.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.




Diagnoses of HIV Infection among Men Who Have Sex with Men, by Age at Diagnosis, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2010–2014—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Young (Aged 13–24 Years) Men Who Have Sex with Men, by Race/Ethnicity, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnosis of HIV Infection among Male Adults and Adolescents, 2010–2014—United States and 6 Dependent Areas

Of the 209,838 diagnoses of HIV infection among male adults and adolescents during 2010 through 2014:

- **80%** were in males
 - 80% of diagnosed HIV infections in males were attributed to male-to-male sexual contact
 - 24% of diagnosed HIV infections in males were among those aged 13–24 years
 - 91% of diagnosed HIV infections in males aged 13–24 years were attributed to male-to-male sexual contact

In 2014, **82%** of HIV infections diagnosed among male adults and adolescents were attributed to male-to-male sexual contact

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.



Men Who Have Sex with Men Living with Diagnosed HIV Infection, by Race/Ethnicity, Year-end 2014—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	1,434	0.3
Asian ^a	8,301	1.6
Black/African American	157,832	30.8
Hispanic/Latino ^b	114,064	22.2
Native Hawaiian/other Pacific Islander	615	0.1
White	212,607	41.4
Multiple races	18,191	3.5
Total^c	513,045	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Deaths among Males with Diagnosed HIV Infection Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 2014—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	15	0.2
Asian ^a	35	0.6
Black/African American	1,939	31.3
Hispanic/Latino ^b	1,080	17.4
Native Hawaiian/other Pacific Islander	4	0.1
White	2,790	45.0
Multiple races	331	5.3
Total^c	6,195	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Deaths of persons with a diagnosed HIV infection may be due to any cause.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.




Stage 3 (AIDS) Classifications among Men who Have Sex with Men, 1985–2014— United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Race/Ethnicity and Year of Diagnosis, 1985–2014—United States and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact and injection drug use.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications and Deaths among Men Who Have Sex with Men with HIV Infection Ever Classified as Stage 3 (AIDS), 1985–2014

US and 6 Dependent Areas



Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Deaths of persons with diagnosed HIV infection may be due to any cause.



Men Who Have Sex with Men Living with HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2014—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	705	0.3
Asian ^a	4,001	1.5
Black/African American	76,876	29.2
Hispanic/Latino ^b	60,487	23.0
Native Hawaiian/other Pacific Islander	327	0.1
White	110,465	42.0
Multiple races	10,062	3.8
Total^c	262,924	100

Note. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Male Adults and Adolescents, by Transmission Category 2015—United States and 6 Dependent Areas

Transmission category	No.	%
Male-to-male sexual contact	26,646	82.2
Injection drug use (IDU)	1,474	4.5
Male-to-male sexual contact and IDU	1,215	3.7
Heterosexual contact ^a	3,034	9.4
Other ^b	53	0.2
Total^c	32,422	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Age Group and Race/Ethnicity, 2015—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	120	0.5
Asian	729	2.7
Black/African American	10,318	38.7
Hispanic/Latino ^a	7,271	27.3
Native Hawaiian/other Pacific Islander	64	0.2
White	7,572	28.4
Multiple races	570	2.1
Total^b	26,646	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use


^a Hispanics/Latinos can be of any race.

^b Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Race/Ethnicity, 2015—United States and 6 Dependent Areas

N = 26,646



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Young (Aged 13–24 Years) Men Who Have Sex with Men, by Race/Ethnicity, 2015—United States and 6 Dependent Areas

N = 7,159




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Men Who Have Sex with Men, by Region of Residence and Race/Ethnicity, 2015—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use. Numbers less than 12, and trends based on these numbers, should be interpreted with caution.

^a Hispanics/Latinos can be of any race



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Race/Ethnicity, 2015—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	36	0.4
Asian ^a	230	2.3
Black/African American	3,930	38.8
Hispanic/Latino ^b	2,510	24.8
Native Hawaiian/other Pacific Islander	16	0.2
White	3,097	30.6
Multiple races	314	3.1
Total^c	10,133	100

Note. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Because column totals for numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Stage 3 (AIDS) Classifications among Men Who Have Sex with Men, by Region of Residence and Race/Ethnicity, 2015—United States and 6 Dependent Areas



Note. Data for the year 2015 are preliminary and based on 6 months reporting delay. Data have been statistically adjusted to account for missing transmission category. Data on men who have sex with men do not include men with HIV infection attributed to male-to-male sexual contact *and* injection drug use.

^a Hispanics/Latinos can be of any race

