



An Individual-Level Single-Session Behavioral Intervention for Young African American Men Who Have Sex with Women

The Research

The Science Behind the Package

FOF is a peer-delivered, clinic-based, individual-level, single-session behavioral intervention (Crosby et. al., 2009). **FOF** aims to educate and motivate clients to use condoms correctly and consistently in order to reduce the spread of HIV and other STDs. It is intended for the participant to receive the intervention one time. **FOF** is based on the Information, Motivation, and Behavioral Skills (IMB) model, which states that people need information, motivation, and practice with a behavior in order to properly learn that behavior. It is also based on Bandura's Social Learning Theory, which states that people learn new behaviors through observational learning, imitation, and modeling.

Target Population

African American men who have sex with women (MSW), ages 18-29, who are seeking care in an STD clinic due to reported symptoms of an STD or who have received an STD diagnosis, who have used a condom in the last three months, and who are not knowingly HIV-positive.

Intervention

FOF is delivered in an STD clinic by a Peer Health Advisor (PHA) who is a trained peer from the community where the intervention is being delivered. Clients are recruited to meet with the PHA for a single 45-60 minute conversation. During this conversation the PHA establishes a rapport with the client, learns about the client's barriers to condom use, discusses how to address errors when using condoms, asks the client to practice correct condom and lubricant use 3+ times on a model, and allows the client to learn about and take home with him a wide variety of high-end condoms and water-based lubricants.

Research Results

A two-year study was conducted to evaluate whether or not the intervention reduced specific risk behaviors and subsequent STDs among eligible men who participated in the intervention at an STD clinic in Kentucky. Over the 6 months following the intervention, men receiving the intervention were significantly:

- Less likely to acquire subsequent STDs
- More likely to report using condoms during last sexual intercourse

Also, men in the intervention group reported significantly:

- Fewer female sex partners
- Fewer acts of unprotected sex

Clients who received **FOF** also had higher proficiency scores for condom application skills.

For Details on the Research Design

Crosby, R., DiClemente, R., et. al. (2009) A Brief, Clinic-Based, Safer Sex Intervention for Heterosexual African American Men Newly Diagnosed With an STD: A Randomized Controlled Trial, *American Journal of Public Health*, 99(1), S96-S103.

The Intervention

A Package Developed from Science

Replicating Effective Programs (REP) is a CDC-initiated project that supports the translation of evidence-based HIV/AIDS prevention interventions into everyday practice, by working with the original researchers in developing a user-friendly package of materials designed for prevention providers. **FOF** is one of the REP interventions and is the product of extensive collaboration among researchers, training and materials developers, STD prevention clinics, and the CDC. The intervention has been field tested in four STD clinics by non-research staff.

Core Elements

Core elements are intervention components that must be maintained without alteration to ensure program effectiveness. The core elements of **FOF** are:

- The trained Peer Health Advisor (PHA) conducts a customized one-to-one counseling session with the client for 45-60 minutes.
- The PHA establishes rapport and a trusting relationship with the client at the beginning of the session.
- The PHA teaches correct condom use skills for clients.
- The PHA and client discuss condom negotiation skills.
- The PHA provides clients with 25+ foils of water-based lubricants and 25+ condoms of their choice from a broad selection of high-end and popular brands.
- The PHA clearly communicates the importance of the client protecting his and his community's future by using condoms correctly and consistently with his partner(s).
- The PHA shows unconditional respect for men and maintains a non-judgmental environment for the client concerning any risk behaviors disclosed.
- The intervention is delivered at a point when the client is feeling vulnerable and is highly concerned about his STD infection status. This may be while he is in the clinic waiting, after a presumptive diagnosis, or after a confirmed lab result.

Package Contents

FOF Implementation Manual that includes the Facilitator's Guide and Monitoring & Evaluation Guide, Technical Assistance Guide, and Promotional & Marketing Materials.

For More Information on the *FOF* Intervention Package

To find out more about package availability and future trainings, please visit <http://www.effectiveinterventions.org>