

PARTNERSHIP FOR HEALTH (Loss-Frame Intervention)

Evidence-Based Structural Intervention

Good Evidence – Risk Reduction

INTERVENTION DESCRIPTION

Goal of Intervention

- Eliminate or reduce unprotected anal or vaginal sex

Target Population

- HIV-positive clinic patients

Brief Description

The *Partnership for Health (PfH)* Loss-frame intervention is a one-on-one, brief provider-administered safer sex intervention for HIV-positive persons in care. The intervention emphasizes the importance of the patient-provider relationship to promote patients' healthful behavior. HIV clinic providers and staff receive a 4-hour training on communication skills building, behavior change theories, delivering a brief counseling session, and role-playing safer-sex counseling. At each patient clinic visit, the provider delivers a brief counseling session with the loss-framed messages that emphasize the negative consequences of unsafe sex, and focus on self-protection, partner protection, and disclosure. The provider also uses the brochures, informational flyers and posters with the loss-framed messages to facilitate counseling and work with the patient to identify goals for the patient to work on.

Theoretical Basis

- Message Framing Theory
- Mutual Participation
- Stages of Changes

Intervention Duration

- A 3- to 5-minute counseling session at every clinic visit over 10 to 11 months

Intervention Setting

- HIV clinics

Deliverer

- Medical providers at the clinics (e.g., physician, physician assistant, nurse, nurse practitioner)

Delivery Methods

- Counseling
- Goal setting
- Printed materials

Structural Mechanism of Change

- Capacity Building – provider training
 - Trained HIV clinic providers deliver safer-sex counseling during patient visits
- Institutional policy/procedure
 - Implementation of provider-delivered safer-sex counseling in HIV clinics for all patients

INTERVENTION PACKAGE INFORMATION

An intervention package was developed with funding from CDC's Replicating Effective Programs (REP) Project. **The intervention package and training are available through CDC's High Impact Prevention Project (HIP): [Partnership for Health \(PfH\)](#).**

EVALUATION STUDY AND RESULTS

Study Location Information

The original evaluation was conducted in Chicago, Illinois between 1999 and 2001.

Key Intervention Effect

- Reduced unprotected sex

Study Sample

The analytic study sample of 585 HIV-positive clinic patients is characterized by the following:

- 41% white, 37% Hispanic/Latino, 16% black or African American, 6% other
- 86% male, 14% female
- 75% homosexual/bisexual, 25% heterosexual
- Mean age of 38 years
- 46% completed high school education or less

Recruitment Settings

Six HIV clinics

Eligibility Criteria

HIV-positive patients were eligible if they were sexually active in the previous 3 months, at least 18 years old, fluent in English or Spanish, and intended to obtain care at the recruitment clinic for the next year.

Assignment Method

Six HIV clinics were randomly assigned to 1 of 3 groups: Loss-frame intervention (n = 2 clinics; 214 patients assessed), Gain-frame intervention (n = 2 clinics; 175 patients assessed), or Medication Adherence comparison (n = 2 clinics; 196 patients assessed). In each clinic, all patients were offered the intervention and sub-samples of clinic patients were systematically selected for assessment.

Comparison Group

The Medication Adherence attention-control comparison group received a one-on-one, brief provider-administered counseling session (3-5 minutes) to promote medication adherence at each clinic visit.

Relevant Outcomes Measured and Follow-up Time

- Sex behaviors during past 3 months (including anal, vaginal and oral sex with or without using a condom with main partners, casual partners, or exchange partners) were measured at 1 to 7 months after the intervention.

Participant Retention

- Loss-Frame Intervention
 - 72% retained at 1 to 7 months after intervention
- Gain-Frame Intervention
 - 58% retained at 1 to 7 months after intervention
- Medication Adherence Comparison
 - 68% retained at 1 to 7 months after intervention

Significant Findings on Relevant Outcomes

- Among HIV-positive patients who had 2 or more sex partners at baseline, those assigned to the Loss-frame intervention were significantly less likely to report unprotected anal/vaginal intercourse than those in the comparison group at 1 to 7 months after intervention ($p = .03$). This intervention effect was also found to be significant among men who have sex with men with 2 or more sex partners at baseline ($p = .04$).
- Among HIV-positive patients who had any casual/exchange partners at baseline, the Loss-frame intervention participants were significantly less likely to report unprotected anal/vaginal intercourse than the comparison participants at 1 to 7 months after intervention ($p = .04$).

Considerations

- The Partnership for Health Loss-frame intervention fails to meet the best-evidence criteria due to a short follow-up time and low retention rates.
- The overall intervention effects for unprotected anal/vaginal sex among HIV-positive patients who had unprotected anal/vaginal sex at baseline were not statistically significant.
- Although the Loss-frame intervention reduced unprotected anal/vaginal sex among HIV-positive patients with 2 or more sex partners, patients with one sexual partner at baseline were unaffected by the intervention.
- The Gain-frame intervention, which has the same format as the loss-frame intervention but emphasizes the benefits or positive consequences of protective behavior, fails to meet the GOOD-EVIDENCE criteria due to no statistically significant intervention effects on sex risk behaviors at the follow-up and low retention rates.

Funding

National Institute of Mental Health grant # RO1 MH57208

California University-wide AIDS Research Program (CC99-SD-003 and ISO2-SD-701)

REFERENCES AND CONTACT INFORMATION

Richardson, J. L., Milam, J., McCutchan, A., Stoyanoff, S., Bolan, R., Weiss, J., . . . Marks, G. (2004). [Effect of brief safer-sex counseling by medical providers to HIV-1 seropositive patients: A multi-clinic assessment.](#) *AIDS*, *18*, 1179-1186.

Researcher: Jean L. Richardson, PhD

University of Southern California
Institute for Prevention Research (IPR)
2001 N Soto Street
3rd Floor, MC 9239, SSB 312E
Los Angeles, California 90089
Email: jeanr@usc.edu

