



HIV in the United States: At a Glance

Total estimated new HIV infections in 2015: 38,500*



About 70% of annual HIV infections are among **gay and bisexual men**.

New HIV Diagnoses in the United States Among the Most-Affected Groups, 2016



39,782
DIAGNOSES IN 2016

26,570
GAY AND BISEXUAL MEN

9,578
HETEROSEXUALS

3,425
PEOPLE WHO INJECT DRUGS (PWID)

From 2011 to 2015, HIV diagnoses:




* New HIV infections are the estimated number of people who get HIV during a year, which is different from the number of people diagnosed with HIV during a year.



African Americans make up **12%** of the US population, but **44%** of new HIV diagnoses.

Hispanics/Latinos make up **18%** of the population, but **25%** of new HIV diagnoses.

HIV Diagnoses by Age, 2016



In 2015, 6,465 people died from HIV disease.

1,122,900
PEOPLE ARE
LIVING WITH HIV.

1 in 7
DON'T KNOW IT.

FOR EVERY 100 PEOPLE
LIVING WITH HIV IN 2014:



** People living with HIV who take HIV medicine as prescribed and get and stay virally suppressed have effectively no risk of sexually transmitting HIV to HIV-negative partners. About one-third of people in HIV care don't get and stay virally suppressed throughout the year.

Reduce Your Risk



Not having sex




Using condoms



Not sharing syringes



Taking medicine to prevent or treat HIV



HIV IS A VIRUS THAT ATTACKS THE BODY'S IMMUNE SYSTEM.

It is usually spread by anal or vaginal sex or sharing syringes with a person who has HIV. The only way to know you have HIV is to be tested. Everyone aged 13-64 should be tested at least once, and people at high risk should be tested at least once a year. Ask your doctor, or visit gettested.cdc.gov to find a testing site. Without treatment, HIV can make a person very sick or may even cause death. If you have HIV, start treatment as soon as possible to stay healthy and help protect your partners.

For More Information

Call 1-800-CDC-INFO (232-4636)
Visit www.cdc.gov/hiv