


HIV Surveillance — Adolescents and Young Adults

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection Among Adolescents and Young Adults Aged 13-24 Years, by Race/Ethnicity, 2009–2013

United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection Among Adolescents and Young Adults Aged 13-24 Years, by Transmission Category, 2009–2013 United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection and Population Among Adolescents Aged 13–19 Years, by Race/Ethnicity 2013—United States

Diagnoses of HIV Infection

N=1,908




American Indian/Alaska Native
Asian
Black/African American

Multiple races

U.S. Population aged 13-19 years

N = 29,524,367




Hispanic/Latino^a
Native Hawaiian/Other Pacific Islander
White

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection and Population Among Young Adults Aged 20–24 Years, by Race/Ethnicity 2013—United States

Diagnoses of HIV Infection
N=8,053



U.S. Population aged 20-24 years
N = 22,795,438



■ American Indian/Alaska Native
■ Asian
■ Black/African American

■ Hispanic/Latino^a
■ Native Hawaiian/Other Pacific Islander
■ White


■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection Among Persons Aged 13 Years and Older, by Sex and Age Group, 2013—United States and 6 Dependent Areas


13–19 years
N=1,931



20–24 years
N=8,144



≥ 25 years
N=37,881



■ **Males** ■ **Females**

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of HIV Infection Among Adolescent and Young Adult Males, by Age Group and Transmission Category 2013—United States and 6 Dependent Areas

Transmission category	13–19 years		20–24 years	
	No.	%	No.	%
Male-to-male sexual contact	1,441	92.6	6,620	91.7
Injection drug use (IDU)	21	1.4	117	1.6
Male-to-male sexual contact and IDU	28	1.8	218	3.0
Heterosexual contact ^a	51	3.3	251	3.5
Other ^b	15	0.9	16	0.2
Total	1,556	100	7,221	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection Among Adolescent and Young Adult Females, by Age Group and Transmission Category 2013—United States and 6 Dependent Areas

Transmission category	13–19 years		20–24 years	
	No.	%	No.	%
Injection drug use	23	6.1	90	9.8
Heterosexual contact ^a	316	84.0	810	87.7
Other ^b	37	9.9	23	2.5
Total	376	100	923	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Rates of Diagnoses of HIV Infection Among Adolescents Aged 13–19 Years, 2013—United States and 6 Dependent Areas

N = 1,931

Total Rate = 6.5




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates of Diagnoses of HIV Infection Among Young Adults Aged 20–24 Years, 2013—United States and 6 Dependent Areas



N = 8,144

Total Rate = 35.3



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Adolescents and Young Adults Aged 13–24 Years Living with a Diagnosis of HIV Infection, by Sex and Race/Ethnicity, Year-end 2012—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

Adolescents and Young Adults Aged 13–24 Years Living with a Diagnosis of HIV Infection, by Sex and Transmission Category, Year-end 2012—United States and 6 Dependent Areas



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU



- Heterosexual contact^a
- Perinatal exposure
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Prevalence of diagnosed HIV Infection Among Adolescents Aged 13–19, Year-end 2012—United States and Dependent Areas

N = 7,300

Total Rate = 24.3




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Prevalence of diagnosed HIV Infection Among Young Adults Aged 20–24 Years, Year-end 2012—United States and 6 Dependent Areas

N = 33,334


Total Rate = 145.7



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.




Stage 3 (AIDS) Classifications Among Adolescents Aged 13–19 Years, by Sex, 1985–2013—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.




Stage 3 (AIDS) Classifications Among Young Adults Aged 20–24 Years, by Sex, 1985–2013—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Stage 3 (AIDS) Classifications Among Persons Aged 13 Years and Older with HIV Infection, by Race/Ethnicity and Age Group, 2013—United States and 6 U.S. Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

Rates of Stage 3(AIDS) Classifications among Adolescents Aged 13–19 Years with HIV Infection, 2013—United States and 6 Dependent Areas

N = 470

Total Rate = 1.6




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Stage 3(AIDS) Classifications among Young Adults Aged 20–24 Years with HIV Infection, 2013—United States and 6 Dependent Areas

N = 2,266


Total Rate = 9.8



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates of Stage 3 (AIDS) Classifications Among Adolescents Aged 13–19 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2012—United States and 6 Dependent Areas

N = 2,203 Total Rate = 7.3




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Stage 3 (AIDS) Classifications Among Young Adults Aged 20–24 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2012—United States and 6 Dependent Areas

N = 8,539 Total Rate = 37.3



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.