


Pediatric HIV Surveillance

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention




Diagnoses of HIV Infection among Children Aged <13 Years, by Age at Diagnosis, 2008–2011 United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of HIV Infection among Children Aged <13 Years, by Race/Ethnicity, 2008–2011—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection and Population in Children Aged <13 Years, by Race/Ethnicity, 2011—United States

Diagnoses of HIV infection


N = 192



American Indian/Alaska Native
Asian
Black/African American

Population, United States

N = 52,950,108



Hispanic/Latino^a
Native Hawaiian/Other Pacific Islander
White

Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Time of Antiretroviral (ARV) Treatment in HIV-Infected Pregnant Women or Perinatally Exposed Infants Birth Years 2008–2011—United States

Time of ARV treatment	Year of birth							
	2008		2009		2010		2011	
	No.	%	No.	%	No.	%	No.	%
During Pregnancy (DP) only	29	1.0	18	0.6	16	0.6	3	0.1
During Labor and Delivery (L&D) only	36	1.2	58	2.0	42	1.6	14	0.7
Infant received ARV after birth (Infant ARV) only	275	9.2	239	8.2	250	9.5	164	7.6
DP and L&D	66	2.2	62	2.1	41	1.6	34	1.6
DP and Infant ARV	137	4.6	129	4.4	109	4.1	81	3.8
L&D and Infant ARV	673	22.5	704	24.1	734	27.9	615	28.7
DP and L&D and Infant ARV	1,467	49.1	1,520	52.0	1,258	47.8	1,049	48.9
No known treatment	303	10.1	195	6.7	181	6.9	186	8.7
Total	2,986	100	2,925	100	2,631	100	2,146	100

Note: Exposure data from 43 areas.



HIV Diagnostic Outcomes among Infants, by Time of Antiretroviral (ARV) Treatment Birth Years 2008–2011—United States


Time of ARV treatment	Infected		Exposed but not infected	Total
	No.	% of Row Total	No.	No.
During Pregnancy (DP) only	4	6.1	62	66
During Labor and Delivery (L&D) only	2	1.3	148	150
Infant received ARV after birth (Infant ARV) only	53	5.7	875	928
DP and L&D	6	3.0	197	203
DP and Infant ARV	11	2.4	445	456
L&D and Infant ARV	77	2.8	2,649	2,726
DP and L&D and Infant ARV	66	1.2	5,228	5,294
No known treatment	115	13.3	750	865
Total	334	3.1	10,354	10,688

Note: Exposure data from 43 areas.



Time of Maternal HIV Testing among Infants with Perinatally Acquired HIV Infection Birth Years 2008–2011—United States


N = 327



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Perinatally Acquired HIV Infections in Children Born During 2011—United States and 6 Dependent Areas

N = 53




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Persons Living with Perinatally Acquired HIV Infection Year-end 2010—United States and 6 Dependent Areas


N = 10,798



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.




Stage 3 (AIDS) Classifications among Perinatally Infected Persons, 1985–2011—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Stage 3 (AIDS) Classifications among Children Aged <13 Years, by Race/Ethnicity, 1985–2011—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Persons Living with Perinatally Acquired HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2010 United States and 6 Dependent Areas

N = 5,134



Note: All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

