


# HIV Surveillance in Women

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention

Division of HIV/AIDS Prevention



# Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2008–2011— United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity 2011— United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	51	5.5
Asian	153	2.3
Black/African American	6,595	40.0
Hispanic/Latino <sup>a</sup>	1,530	7.9
Native Hawaiian/Other Pacific Islander	8	3.9
White	1,776	2.0
Multiple races	144	7.5
<b>Total</b>	<b>10,257</b>	<b>7.7</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.


<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity 2011—United States


## Diagnoses of HIV Infection

N=10,257



## Female Population

N = 132,402,857



■ American Indian/Alaska Native  
■ Asian  
■ Black/African American

■ Hispanic/Latino<sup>a</sup>  
■ Native Hawaiian/Other Pacific Islander  
■ White

■ Multiple races


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.




# Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity and Transmission Category 2011—United States and 6 Dependent Areas



**Black/African American**  
N=6,602



**Hispanic/Latino<sup>a</sup>**  
N=1,776



**White**  
N=1,776



*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Diagnoses of HIV Infection among Adult and Adolescent Females, by Transmission Category and Age at Diagnosis 2011—United States and 6 Dependent Areas

## Age at Diagnosis (in years)

Transmission category	13–19	20–24	25–34	35–44	≥45
	N=523 %	N=1,142 %	N=2,649 %	N=2,581 %	N=3,618 %
Injection drug use	7.0	8.8	12.7	14.2	17.4
Heterosexual contact <sup>a</sup>	92.7	91.2	87.2	85.8	82.2
Other <sup>b</sup>	0.3	0.0	0.0	0.1	0.3
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Rates of Diagnoses of HIV Infection among Adult and Adolescent Females, 2011—United States and 6 Dependent Areas


**N = 10,512**      **Total rate = 7.8**



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Diagnoses of HIV Infection among Adult and Adolescent Females, by Region and Race/Ethnicity 2011—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Deaths of Adult and Adolescent Females with Diagnosed HIV Infection, by Race/Ethnicity 2010—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	24	2.6
Asian <sup>a</sup>	23	0.3
Black/African American	3,241	19.9
Hispanic/Latino <sup>b</sup>	628	3.3
Native Hawaiian/Other Pacific Islander	6	2.8
White	902	1.0
Multiple races	214	11.5
<b>Total<sup>c</sup></b>	<b>5,039</b>	<b>3.8</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.


<sup>c</sup> Includes persons of unknown race/ethnicity.



# Rates of Adult and Adolescent Females Living with Diagnosed HIV Infection, Year-end 2010—United States and 6 Dependent Areas

N = 223,045


Total rate = 167.5



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.




# Stage 3 (AIDS) Classifications among Adult and Adolescent Females, 1985–2011—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Race/Ethnicity and Year of Diagnosis, 1985–2011—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Includes Asian/Pacific Islander legacy cases.



# Percentages of Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Race/Ethnicity and Year of Diagnosis 1985–2011—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Includes Asian/Pacific Islander legacy cases.



## Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Race/Ethnicity, 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	38	4.0
Asian <sup>a</sup>	86	1.3
Black/African American	5,311	32.2
Hispanic/Latino <sup>b</sup>	1,130	5.9
Native Hawaiian/Other Pacific Islander	7	3.4
White	1,202	1.4
Multiple races	176	9.2
<b>Total</b>	<b>7,949</b>	<b>6.0</b>

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



## Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Transmission Category and Age at Diagnosis 2011—United States and 6 Dependent Areas

Transmission category	Age at Diagnosis (in years)				
	13–19	20–24	25–34	35–44	≥45
	N=187	N=414	N=1,777	N=2,352	N=3,372
	%	%	%	%	%
Injection drug use	5.7	11.9	14.1	18.5	26.6
Heterosexual contact <sup>a</sup>	51.8	82.3	85.5	81.4	72.9
Other <sup>b</sup>	42.5	5.8	0.4	0.1	0.5
<b>Total</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>

Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Females with HIV Infection, 2011—United States and 6 Dependent Areas


**N=8,102**      **Total Rate = 6.0**



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Stage 3 (AIDS) Classifications among Adult and Adolescent Females, by Region and Race/Ethnicity 2011—United States




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Adult and Adolescent Females Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS) Year-end 2010—United States and 6 Dependent Areas

N=118,538 Total rate = 89.0



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

