


HIV Surveillance by Race/Ethnicity

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2008–2012 United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity 2008–2012—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.



HIV Infection in Blacks/African Americans United States and 6 Dependent Areas

Of the 238,139 diagnoses of HIV infection from 2008–2012,
blacks/African Americans accounted for:

47% of total

64% of women

65% of infections attributed to heterosexual contact^a

68% of children aged <13 years

In 2012, 46% of diagnoses of HIV infection among adults and
adolescents were in blacks/African Americans.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



HIV Infection in Hispanics/Latinos United States and 6 Dependent Areas

Of the 238,139 diagnoses of HIV infection from 2008–2012, Hispanics/Latinos^a accounted for:

- 22% of total

- 16% of women

- 18% of infections attributed to heterosexual contact^b

- 13% of children aged <13 years

In 2012, 22% of diagnoses of HIV infection among adults and adolescents were in Hispanics/Latinos.

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting


^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.




Diagnoses of HIV Infection and Population by Race/Ethnicity, 2012—United States

Diagnoses of HIV infection
N = 47,989



■ American Indian/Alaska Native
■ Asian
■ Black/African American

Population, United States
N = 313,914,040



■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White

■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection and Population among Adult and Adolescent Males, by Race/Ethnicity 2012—United States


Diagnoses of HIV Infection

N = 38,160



Male Population, United States

N = 127,481,432




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity 2012—United States

Diagnoses of HIV Infection




Female Population, United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, by Sex and Transmission Category 2012—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Transmission Category 2012—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adult and Adolescent Whites, by Sex and Transmission Category 2012—United States and 6 Dependent Areas



■ Male-to-male sexual contact
■ Injection drug use
■ Male-to-male sexual contact and IDU



■ Heterosexual contact^a
■ Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, by Sex and Place of Birth 2012—United States and 6 Dependent Areas

Males
N = 9,168



Females
N = 1,526



■ United States
■ Central America
■ South America
■ Cuba
■ Mexico
■ Puerto Rico
■ Other
■ Unknown

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2012—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	183	20.2
Asian	801	12.9
Black/African American	16,112	108.9
Hispanic/Latino ^a	8,516	42.0
Native Hawaiian/other Pacific Islander	67	31.8
White	11,638	14.0
Multiple races	842	46.0
Total	38,160	29.9

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2012—United States


Race/ethnicity	No.	Rate
American Indian/Alaska Native	45	4.7
Asian	146	2.1
Black/African American	6,293	37.8
Hispanic/Latino ^a	1,289	6.5
Native Hawaiian/Other Pacific Islander	11	5.1
White	1,620	1.9
Multiple races	184	9.3
Total	9,586	7.2

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adults and Adolescents, by Region and Race/Ethnicity 2012—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, 2012—United States

N = 22,405

Total Rate = 71.2




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates of Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos^a, 2012—United States

N = 9,805

Total Rate = 24.5



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.



Rates of Diagnoses of HIV Infection among Adult and Adolescent Whites, 2012—United States

N = 13,258

Total Rate = 7.8



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Deaths of Persons with Diagnosed HIV Infection by Race/Ethnicity, 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	64	2.8
Asian ^a	96	0.6
Black/African American	8,329	21.7
Hispanic/Latino ^b	2,806	5.4
Native Hawaiian/other Pacific Islander	13	2.5
White	5,257	2.7
Multiple races	775	13.3
Total^c	17,339	5.6

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.



Adults and Adolescents Living with Diagnosed HIV Infection by Race/Ethnicity, Year-end 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	2,787	152.3
Asian ^a	10,278	80.3
Black/African American	378,268	1,217.2
Hispanic/Latino ^b	171,958	440.9
Native Hawaiian/other Pacific Islander	812	198.8
White	288,471	169.8
Multiple races	25,254	685.2
Total^c	877,828	339.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.




Stage 3 (AIDS) Classifications among Racial/Ethnic Minorities with Diagnosed HIV Infection, 1985–2012 United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Includes American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, and Native Hawaiians/other Pacific Islanders. Hispanics/Latinos can be of any race.

Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity, 1985–2012 United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Rates of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity 2000–2012—United States





Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications and Population, by Race/Ethnicity, 2012—United States



■ American Indian/Alaska Native
■ Asian^b
■ Black/African American

■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White

■ Multiple races

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.

Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Blacks/African Americans with Diagnosed HIV Infection, 2012—United States

N = 14,094

Total Rate = 44.8




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.



Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Hispanics/Latinos^a with Diagnosed HIV Infection, 2012—United States

N = 5,418

Total Rate = 13.5



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.


^aHispanics/Latinos can be of any race.



Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Whites with Diagnosed HIV Infection, 2012—United States

N = 6,932


Total Rate = 4.1



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.



Survival after Classification of Stage 3 (AIDS) during 1998–2008, by Months Survived and Race/Ethnicity United States and 6 Dependent Areas



Note. Data exclude persons whose month of diagnosis or month of death is unknown.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2011—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	1,513	82.7
Asian ^a	5,639	44.1
Black/African American	208,171	669.8
Hispanic/Latino ^b	103,563	265.5
Native Hawaiian/Other Pacific Islander	443	108.3
White	159,065	93.6
Multiple races	15,851	430.1
Total^c	494,245	191.1

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.

^c Includes persons of unknown race/ethnicity.

