


# HIV Surveillance by Race/Ethnicity

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention  
Division of HIV/AIDS Prevention



# Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2009–2013—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity 2009–2013—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.



# HIV Infection in Blacks/African Americans United States and 6 Dependent Areas

Total 233,100 diagnoses of HIV infection in 2009–2013

- Blacks/African Americans: 46% of total
- Black/African American women: 63% of women
- Blacks/African Americans: 64% of infections attributed to heterosexual contact<sup>a</sup>
- Black/African American children: 67% of children aged <13 years

In 2013, 45% of diagnoses of HIV infection among adults and adolescents were in blacks/African Americans.

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



# HIV Infection in Hispanics/Latinos United States and 6 Dependent Areas

Total 233,100 diagnoses of HIV infection from 2009–2013

- Hispanics/Latinos: 22% of total
- Hispanic/Latino women: 17% of women
- Hispanics/Latinos : 18% of infections attributed to heterosexual contact<sup>a</sup>
- Hispanic/Latino children: 12% of children aged <13 years

In 2013, 23% of diagnoses of HIV infection among adults and adolescents were in Hispanics/Latinos.

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting


<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.




# Diagnoses of HIV Infection and Population by Race/Ethnicity, 2013—United States

**Diagnoses of HIV infection**  
N = 47,352



**Population, United States**  
N = 316,128,839



■ American Indian/Alaska Native  
■ Asian  
■ Black/African American  
■ Multiple races

■ Hispanic/Latino<sup>a</sup>  
■ Native Hawaiian/other Pacific Islander  
■ White

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection and Population among Adult and Adolescent Males, by Race/Ethnicity 2013—United States


**Diagnoses of HIV Infection**

N = 37,887



**Male Population, United States**

N = 128,717,938



American Indian/Alaska Native

Asian

Black/African American

Multiple races

Hispanic/Latino<sup>a</sup>

Native Hawaiian/Other Pacific Islander

White


Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.




# Diagnoses of HIV Infection and Population among Adult and Adolescent Females, by Race/Ethnicity 2013—United States

**Diagnoses of HIV Infection**



**Female Population, United States**



■ American Indian/Alaska Native  
■ Asian  
■ Black/African American

■ Hispanic/Latino<sup>a</sup>  
■ Native Hawaiian/other Pacific Islander  
■ White


■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, by Sex and Transmission Category 2013—United States and 6 Dependent Areas




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos<sup>a</sup>, by Sex and Transmission Category 2013—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Diagnoses of HIV Infection among Adult and Adolescent Whites, by Sex and Transmission Category 2013—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




# Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos<sup>a</sup>, by Sex and Place of Birth 2013—United States and 6 Dependent Areas

**Males**  
N = 9,266



**Females**  
N = 1,610



- United States
- Central America
- South America
- Cuba
- Mexico
- Puerto Rico
- Other
- Unknown

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.



## Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	169	18.4
Asian	799	12.5
Black/African American	15,847	105.7
Hispanic/Latino <sup>a</sup>	8,686	41.8
Native Hawaiian/other Pacific Islander	52	24.1
White	11,495	13.8
Multiple races	839	44.1
<b>Total</b>	<b>37,887</b>	<b>29.4</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

<sup>a</sup> Hispanics/Latinos can be of any race.



## Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2013—United States


Race/ethnicity	No.	Rate
American Indian/Alaska Native	49	5.1
Asian	159	2.2
Black/African American	5,867	34.8
Hispanic/Latino <sup>a</sup>	1,419	7.0
Native Hawaiian/Other Pacific Islander	15	7.3
White	1,579	1.8
Multiple races	190	9.3
<b>Total</b>	<b>9,278</b>	<b>6.9</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

<sup>a</sup> Hispanics/Latinos can be of any race.



# Diagnoses of HIV Infection among Adults and Adolescents, by Region and Race/Ethnicity 2013—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Diagnoses of HIV Infection among Adult and Adolescent Blacks/African Americans, 2013—United States

N = 21,714

Total Rate = 68.2




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Rates of Diagnoses of HIV Infection among Adult and Adolescent Hispanics/Latinos<sup>a</sup>, 2013—United States

N = 10,104

Total Rate = 24.7



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Diagnoses of HIV Infection among Adult and Adolescent Whites, 2013—United States

N = 13,075

Total Rate = 7.7



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



# Deaths of Persons with Diagnosed HIV Infection by Race/Ethnicity, 2012—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	63	2.7
Asian <sup>a</sup>	73	0.5
Black/African American	8,171	21.1
Hispanic/Latino <sup>b</sup>	2,586	4.9
Native Hawaiian/other Pacific Islander	4	0.7
White	5,426	2.7
Multiple races	849	14.1
<b>Total<sup>c</sup></b>	<b>17,173</b>	<b>5.5</b>

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.

<sup>c</sup> Includes persons of unknown race/ethnicity.



# Adults and Adolescents Living with Diagnosed HIV Infection by Race/Ethnicity, Year-end 2012—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	2,860	154.3
Asian <sup>a</sup>	10,966	83.1
Black/African American	389,857	1,239.1
Hispanic/Latino <sup>b</sup>	183,856	460.2
Native Hawaiian/other Pacific Islander	862	206.5
White	294,691	173.0
Multiple races	29,215	765.1
<b>Total<sup>c</sup></b>	<b>912,308</b>	<b>349.5</b>

*Note.* Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.


<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.

<sup>c</sup> Includes persons of unknown race/ethnicity.




# Stage 3 (AIDS) Classifications among Racial/Ethnic Minorities with Diagnosed HIV Infection, 1985–2013 United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Includes American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, and Native Hawaiians/other Pacific Islanders. Hispanics/Latinos can be of any race.



# Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity, 1985–2013 United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.

<sup>b</sup> Includes Asian/Pacific Islander legacy cases.



# Rates of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity 2000–2013—United States



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

<sup>a</sup> Hispanics/Latinos can be of any race.


<sup>b</sup> Includes Asian/Pacific Islander legacy cases.



# Stage 3 (AIDS) Classifications and Population, by Race/Ethnicity, 2013—United States


## Stage 3 (AIDS)

N = 26,688



## U.S. Population

N = 316,128,839



■ American Indian/Alaska Native  
■ Asian<sup>b</sup>  
■ Black/African American

■ Hispanic/Latino<sup>a</sup>  
■ Native Hawaiian/other Pacific Islander  
■ White

■ Multiple races

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

<sup>a</sup> Hispanics/Latinos can be of any race.


<sup>b</sup> Includes Asian/Pacific Islander legacy cases.



# Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Blacks/African Americans with Diagnosed HIV Infection, 2013—United States

N = 13,168

Total Rate = 41.3




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.



# Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Hispanics/Latinos<sup>a</sup> with Diagnosed HIV Infection, 2013—United States

N = 5,334

Total Rate = 13.0



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.


<sup>a</sup> Hispanics/Latinos can be of any race.



# Rates of Stage 3 (AIDS) Classifications among Adult and Adolescent Whites with Diagnosed HIV Infection, 2013— United States

N = 6,759


Total Rate = 4.0



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are not provided for the U.S. dependent areas because U.S. census information for race/ethnicity is limited for these areas.



# Survival after Classification of Stage 3 (AIDS) during 1998–2009, by Months Survived and Race/Ethnicity United States and 6 Dependent Areas



Note. Data exclude persons whose month of diagnosis or month of death is unknown.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.



# Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, Year-end 2012—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	1,541	83.2
Asian <sup>a</sup>	5,853	44.3
Black/African American	213,170	677.5
Hispanic/Latino <sup>b</sup>	109,416	273.9
Native Hawaiian/Other Pacific Islander	461	110.4
White	160,108	94.0
Multiple races	17,990	471.1
<b>Total<sup>c</sup></b>	<b>508,539</b>	<b>194.8</b>

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum up to the total column.

<sup>a</sup> Includes Asian/Pacific Islander legacy cases.

<sup>b</sup> Hispanics/Latinos can be of any race.

<sup>c</sup> Includes persons of unknown race/ethnicity.

