

Table 1: Vaccine Requirements According to Applicant Age

Vaccines by applicant age	Birth– 1 month	2–11 months	12 months–6 years	7–10 years	11–17 years	18–64 years	≥ 65 years
DTP/DTaP/DT	NO	YES		No			
Td/Tdap	NO			Sometimes*	YES, substitute 1-time dose of Tdap for Td booster; then boost with Td or Tdap every 10 years		
Polio** (IPV/OPV)	NO	YES				NO	
Measles, Mumps, and Rubella	NO		YES, if born in 1957 or later			NO	
Rotavirus***	NO	YES 6 weeks through 8 months	NO				
Hib	NO	YES 2 through 59 months old		NO			
Hepatitis A	NO		YES 12 months through 18 years old			NO	
Hepatitis B	YES, through 59 years old					NO	
Meningococcal (MenACWY)	NO				Yes 11 through 18 years old	NO	
Varicella	NO		YES				
Pneumococcal	NO	YES, 2 through 59 months old (administer PCV)		NO			One dose PCV15 followed by PPSV23 or one dose PCV20
Influenza	NO, if less than six months old		YES, ≥ 6 months (annually when flu vaccine is available in country of exam)				
COVID-19	See the COVID-19 Technical Instructions for the details of this requirement.						

DTP=pediatric formulation diphtheria and tetanus toxoids and pertussis vaccine; DTaP=pediatric formulation diphtheria and tetanus toxoids and acellular pertussis vaccine; DT=pediatric formulation diphtheria and tetanus toxoids; Td=adult formulation tetanus and diphtheria toxoids; Tdap=adolescent and adult formulation tetanus and diphtheria toxoids and acellular pertussis vaccine (*Children 7-10 years old sometimes need a dose of Tdap depending on their vaccine history. See [Diphtheria, Tetanus and Pertussis-Containing Vaccines Catch-Up Guidance](#) on CDC’s website for additional information). IPV=inactivated poliovirus vaccine (killed); OPV=oral poliovirus vaccine (live); Hib=*Haemophilus influenzae* type b conjugate vaccine; MenACWY=quadravalent meningococcal conjugate vaccine; PCV=pneumococcal conjugate vaccine; PPSV=pneumococcal polysaccharide vaccine. ** Please see posted [Addendum to Technical Instructions for Panel Physicians for Vaccinations](#) on CDC’s

website for changing guidance about polio vaccine. ***Rotavirus vaccination should not be initiated for infants aged 15 weeks 0 days or older.

This table describes vaccine requirements for U.S. immigrant visa and status adjustment applicants and does not include recommendations for other clinical purposes. See the [Immunization Schedules](#) on CDC's website for number and spacing of doses for required vaccines.

