



MMWRTM

Morbidity and Mortality Weekly Report

www.cdc.gov/mmwr

Recommendations and Reports

April 18, 2008 / Vol. 57 / No. RR-2

Compendium of Animal Rabies Prevention and Control, 2008

**National Association of State Public
Health Veterinarians, Inc.**

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

Suggested Citation: Centers for Disease Control and Prevention. [Title]. *MMWR* 2008;57(No. RR-#):[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH
Director

Tanja Popovic, MD, PhD
Chief Science Officer

James W. Stephens, PhD
Associate Director for Science

Steven L. Solomon, MD
Director, Coordinating Center for Health Information and Service

Jay M. Bernhardt, PhD, MPH
Director, National Center for Health Marketing

Katherine L. Daniel, PhD
Deputy Director, National Center for Health Marketing

Editorial and Production Staff

Frederic E. Shaw, MD, JD
Editor, MMWR Series

Teresa F. Rutledge
(Acting) Managing Editor, MMWR Series

David C. Johnson
(Acting) Lead Technical Writer-Editor

Patricia A. McGee
Project Editor

Peter M. Jenkins
(Acting) Lead Visual Information Specialist

Lynda G. Cupell
Malbea A. LaPete
Visual Information Specialists

Quang M. Doan, MBA
Erica R. Shaver
Information Technology Specialists

Editorial Board

- William L. Roper, MD, MPH, Chapel Hill, NC, Chairman
- Virginia A. Caine, MD, Indianapolis, IN
- David W. Fleming, MD, Seattle, WA
- William E. Halperin, MD, DrPH, MPH, Newark, NJ
- Margaret A. Hamburg, MD, Washington, DC
- King K. Holmes, MD, PhD, Seattle, WA
- Deborah Holtzman, PhD, Atlanta, GA
- John K. Iglehart, Bethesda, MD
- Dennis G. Maki, MD, Madison, WI
- Sue Mallonee, MPH, Oklahoma City, OK
- Stanley A. Plotkin, MD, Doylestown, PA
- Patricia Quinlisk, MD, MPH, Des Moines, IA
- Patrick L. Remington, MD, MPH, Madison, WI
- Barbara K. Rimer, DrPH, Chapel Hill, NC
- John V. Rullan, MD, MPH, San Juan, PR
- Anne Schuchat, MD, Atlanta, GA
- Dixie E. Snider, MD, MPH, Atlanta, GA
- John W. Ward, MD, Atlanta, GA

CONTENTS

Part I: Rabies Prevention and Control 1

Part II: Recommendations for Parenteral Rabies
Vaccination Procedures 6

Part III: Rabies Vaccines Licensed and Marketed
in the United States, 2008 7

References 8

This report is being published as a courtesy to the National Association of State Public Health Veterinarians, Inc., and to the *MMWR* readership.

Compendium of Animal Rabies Prevention and Control, 2008*

National Association of State Public Health Veterinarians, Inc. (NASPHV)

Rabies is a fatal viral zoonosis and a serious public health problem (1). The disease is an acute, progressive encephalitis caused by a lyssavirus. Although the United States has been declared free of canine rabies virus variant transmission, multiple viral variants are maintained in wild mammal populations, and there is always a risk of reintroduction of canine rabies (2). All mammals are believed to be susceptible to the disease, and for purposes of this document, use of the term “animal” refers to mammals.

The recommendations in this compendium serve as a basis for animal rabies-prevention and -control programs throughout the United States and facilitate standardization of procedures among jurisdictions, thereby contributing to an effective national rabies-control program. This document is reviewed annually and revised as necessary. The most current version replaces all previous versions. These recommendations do not supersede state and local laws or requirements. Principles of rabies-prevention and -control are detailed in Part I; recommendations for parenteral vaccination procedures are presented in Part II; and all animal rabies vaccines licensed by the U. S. Department of Agriculture (USDA) and marketed in the United States are listed in Part III.

Part I: Rabies Prevention and Control

A. Principles of Rabies Prevention and Control

- 1. Rabies Exposure.** Rabies is transmitted only when the virus is introduced into bite wounds, open cuts in skin, or onto mucous membranes from saliva or other potentially infectious material such as neural tissue (3). Questions regarding possible exposures should be directed promptly to state or local public health authorities.
- 2. Public Health Education.** Essential components of rabies prevention and control include ongoing public education, responsible pet ownership, routine veterinary care, and professional continuing education. The majority of animal and human exposures to rabies can be prevented by raising awareness concerning: rabies transmission routes, avoiding contact with wildlife, and following appropriate veterinary care. Prompt recognition and reporting of possible

exposures to medical professionals and local public health authorities is critical.

- 3. Human Rabies Prevention.** Rabies in humans can be prevented either by eliminating exposures to rabid animals or by providing exposed persons with prompt local treatment of wounds combined with the administration of human rabies immune globulin and vaccine. The rationale for recommending preexposure and postexposure rabies prophylaxis and details of their administration can be found in the current recommendations of the Advisory Committee on Immunization Practices (ACIP) (3). These recommendations, along with information concerning the current local and regional epidemiology of animal rabies and the availability of human rabies biologics, are available from state health departments.
- 4. Domestic Animals.** Local governments should initiate and maintain effective programs to ensure vaccination of all dogs, cats, and ferrets and to remove strays and unwanted animals. Such procedures in the United States have reduced laboratory-confirmed cases of rabies in dogs from 6,949 in 1947 to 71 in 2006 (2). Because more rabies cases are reported annually involving cats (247 in 2006) than dogs, vaccination of cats should be required (2). Animal shelters and animal-control authorities should establish policies to ensure that adopted animals are vaccinated against rabies. The recommended vaccination procedures and the licensed animal vaccines are specified in Parts II and III of this compendium, respectively.
- 5. Rabies in Vaccinated Animals.** Rabies is rare in vaccinated animals (4). If such an event is suspected, it should be reported to state public health officials; the vaccine manufacturer; and USDA, Animal and Plant Health Inspection Service, Center for Veterinary

* **The NASPHV Committee:** Ben Sun, DVM, MPVM, Chair; Michael Auslander, DVM, MSPH; Catherine M. Brown, DVM, MSc, MPH; Lisa Conti, DVM, MPH; Paul Ettestad, DVM, MS; Mira J. Leslie, DVM, MPH; Faye E. Sorhage, VMD, MPH.

Consultants to the Committee: Keith Friendshuh, DVM, American Veterinary Medical Association (AVMA); Donna M. Gatewood, DVM, MS, United States Department of Agriculture (USDA), Center for Veterinary Biologics; Suzanne R. Jenkins, VMD, MPH; Lorraine Moule, National Animal Control Association (NACA); Barbara Nay, Animal Health Institute; Raoult Ratard, MD, MS, MPH, Council of State and Territorial Epidemiologists (CSTE); Charles E. Rupprecht, VMD, MS, PhD, CDC; Dennis Slate, PhD, USDA Wildlife Services; Charles V. Trimarchi, MS, American Public Health Laboratory Association (APHL); Burton Wilcke, Jr., PhD, American Public Health Association (APHA). This compendium has been endorsed by APHA, APHL, AVMA, CDC, CSTE, and NACA.

Corresponding Preparer: Ben Sun, DVM, MPVM, State Public Health Veterinarian, California Department of Public Health, Veterinary Public Health Section, MS 7308, P.O. Box 997377, Sacramento, CA 95899-7377. Telephone: 916-552-9740; Fax: 916-552-9725; E-mail: Ben.Sun@cdph.ca.gov.

Biologics (Internet: http://www.aphis.usda.gov/animal_health/vet_biologics/vb_adverse_event.shtml; telephone: 800-752-6255; or e-mail: CVB@usda.gov). The laboratory diagnosis should be confirmed and the virus variant should be characterized by a rabies reference laboratory. A thorough epidemiologic investigation should be conducted.

6. **Rabies in Wildlife.** The control of rabies among wildlife reservoirs is difficult (5). Vaccination of free-ranging wildlife or selective population reduction might be useful in some situations, but the success of such procedures depends on the circumstances surrounding each rabies outbreak (see Part I. C.). Because of the risk of rabies in wild animals (especially raccoons, skunks, coyotes, foxes, and bats), the AVMA, CSTE, NACA, and NASPHV strongly recommend the enactment and enforcement of state laws prohibiting their importation, distribution, translocation, and private ownership.
7. **Rabies Surveillance.** Laboratory-based rabies surveillance and variant typing are essential components of rabies-prevention and -control programs. Accurate and timely information is necessary to guide human postexposure prophylaxis decisions, determine the management of potentially exposed animals, aid in emerging pathogen discovery, describe the epidemiology of the disease, and assess the need for and effectiveness of vaccination programs for wildlife.
8. **Rabies Diagnosis.** Rabies testing should be performed in accordance with the established national standardized protocol for rabies testing (http://www.cdc.gov/rabies/docs/standard_dfa_protocol_rabies.pdf) by a qualified laboratory that has been designated by the local or state health department (6,7). Euthanasia (8) should be accomplished in such a way as to maintain the integrity of the brain so that the laboratory can recognize the anatomical parts. Except in the case of very small animals, such as bats, only the head or brain (including brain stem) should be submitted to the laboratory. To facilitate laboratory processing and prevent a delay in testing, any animal or animal specimen being submitted for testing should preferably be stored and shipped under refrigeration and not be frozen. Chemical fixation of tissues should be avoided to prevent significant testing delays and because it might preclude reliable testing. Questions about testing of fixed tissues should be directed to the local rabies laboratory or public health department.

9. **Rabies Serology.** Some “rabies-free” jurisdictions might require evidence of vaccination and rabies virus antibodies for animal importation purposes. Rabies virus antibody titers are indicative of a response to vaccine or infection. Titers do not directly correlate with protection because other immunologic factors also play a role in preventing rabies, and our abilities to measure and interpret those other factors are not well-developed. Therefore, evidence of circulating rabies virus antibodies should not be used as a substitute for current vaccination in managing rabies exposures or determining the need for booster vaccinations in animals (9–11).

10. **Rabies Research.** Information derived from well-designed studies is essential for the development of science-based recommendations. Data are needed in several areas, including viral shedding periods for livestock and lagomorphs, potential shedding of virus in milk, earliest age at which rabies vaccination is effective, postexposure prophylaxis for domestic animals, extra label vaccine use in domestic animals and wildlife rabies reservoirs, and the ecology of wildlife rabies reservoir species, especially in relation to the use of oral rabies vaccines.

B. Prevention and Control Methods in Domestic and Confined Animals.

1. **Preexposure Vaccination and Management.** Parenteral animal rabies vaccines should be administered only by or under the direct supervision of a veterinarian. Rabies vaccinations may also be administered under the supervision of a veterinarian to animals held in animal-control shelters before release. Any veterinarian signing a rabies certificate must ensure that the person administering vaccine is identified on the certificate and is appropriately trained in vaccine storage, handling, administration, and in the management of adverse events. This practice assures that a qualified and responsible person can be held accountable for properly vaccinating the animal.

Within 28 days after initial vaccination, a peak rabies virus antibody titer is reached, and the animal can be considered immunized (12). An animal is currently vaccinated and is considered immunized if the initial vaccination was administered at least 28 days previously or booster vaccinations have been administered in accordance with this compendium.

Regardless of the age of the animal at initial vaccination, a booster vaccination should be administered 1 year later (see Parts II and III for vaccines and proce-

dures). No laboratory or epidemiologic data exist to support the annual or biennial administration of 3- or 4-year vaccines after the initial series. Because a rapid anamnestic response is expected, an animal is considered currently vaccinated immediately after a booster vaccination (13).

a. Dogs, Cats, and Ferrets. All dogs, cats, and ferrets should be vaccinated against rabies and revaccinated in accordance with Part III of this compendium. If a previously vaccinated animal is overdue for a booster, it should be revaccinated. Immediately after the booster, the animal is considered currently vaccinated and should be placed on a schedule, depending on the labeled duration of the vaccine used.

b. Livestock. Consideration should be given to vaccinating livestock that are particularly valuable. Animals that have frequent contact with humans (e.g., in petting zoos, fairs, and other public exhibitions) and horses traveling interstate should be currently vaccinated against rabies (14,15).

c. Confined Animals.

1.) Wild. No parenteral rabies vaccines are licensed for use in wild animals or hybrids (the offspring of wild animals crossbred to domestic animals). Wild animals or hybrids should not be kept as pets (16–19).

2.) Maintained in Exhibits and in Zoological Parks. Captive mammals that are not completely excluded from all contact with rabies vectors can become infected. Moreover, wild animals might be incubating rabies when initially captured; therefore, wild-caught animals susceptible to rabies should be quarantined for a minimum of 6 months. Employees who work with animals at such facilities should receive preexposure rabies vaccination. The use of pre- or postexposure rabies vaccinations for handlers who work with animals at such facilities might reduce the need for euthanasia of captive animals that expose handlers. Carnivores and bats should be housed in a manner that precludes direct contact with the public (14).

2. Stray Animals. Stray dogs, cats, and ferrets should be removed from the community. Local health departments and animal-control officials can enforce the removal of strays more effectively if owned animals have identification and are confined or kept on leash. Strays should be impounded for at least 3 business days to determine if human exposure has occurred and to give owners sufficient time to reclaim animals.

3. Importation and Interstate Movement of Animals.

a. International. CDC regulates the importation of dogs and cats into the United States. Importers of dogs must comply with rabies vaccination requirements (42 CFR, Part 71.51[c] [<http://www.cdc.gov/ncidod/dq/animal.htm>]) and complete CDC form 75.37 (http://www.cdc.gov/ncidod/dq/pdf/animal/dog_quarantine_notice_08-04-06-cdc7537.pdf). The appropriate health official of the state of destination should be notified within 72 hours of the arrival of any imported dog required to be placed in confinement under the CDC regulation. Failure of the owner to comply with these confinement requirements should be promptly reported to the Division of Global Migration and Quarantine, CDC (telephone: 404-639-3441).

Federal regulations alone are insufficient to prevent the introduction of rabid animals into the United States (20,21). All imported dogs and cats are subject to state and local laws governing rabies and should be currently vaccinated against rabies in accordance with this compendium. Failure of the owner to comply with state or local requirements should be referred to the appropriate state or local official.

b. Interstate. Before interstate movement (including commonwealths and territories), dogs, cats, ferrets, and horses should be currently vaccinated against rabies in accordance with this compendium's recommendations (see Part I. B.1.). Animals in transit should be accompanied by a currently valid NASPHV Form 51, Rabies Vaccination Certificate (<http://www.nasphv.org/Documents/RabiesVacCert.pdf>). When an interstate health certificate or certificate of veterinary inspection is required, it should contain the same rabies vaccination information as Form 51.

c. Areas with Dog-to-Dog Rabies Transmission. Canine rabies virus variants have been eliminated in the United States (2). Rabid dogs have been introduced into the continental United States from areas with dog-to-dog rabies transmission (20,21). This practice poses the risk of introducing canine-transmitted rabies to areas where it does not currently exist. The movement of dogs for the purposes of adoption or sale from areas with dog-to-dog rabies transmission should be prohibited.

4. Adjunct Procedures. Methods or procedures that enhance rabies control include the following:

a. Identification. Dogs, cats, and ferrets should be identified (e.g., metal or plastic tags or microchips) to allow for verification of rabies vaccination status.

- b. Licensure.** Registration or licensure of all dogs, cats, and ferrets is an integral component of an effective rabies-control program. A fee is frequently charged for such licensure, and revenues collected are used to maintain rabies- or animal-control activities. Evidence of current vaccination should be an essential prerequisite to licensure.
 - c. Canvassing.** House-to-house canvassing by animal-control officials facilitates enforcement of vaccination and licensure requirements.
 - d. Citations.** Citations are legal summonses issued to owners for violations, including the failure to vaccinate or license their animals. The authority for officers to issue citations should be an integral part of each animal-control program.
 - e. Animal Control.** All local jurisdictions should incorporate stray animal control, leash laws, animal bite prevention, and training of personnel in their programs.
 - f. Public Education.** All local jurisdictions should incorporate education covering responsible pet ownership, bite prevention, and appropriate veterinary care in their programs.
- 5. Postexposure Management.** This section refers to any animal exposed (see Part I.A.1.) to a confirmed or suspected rabid animal. Wild mammalian carnivores or bats that are not available or suitable for testing should be regarded as rabid animals.
- a. Dogs, Cats, and Ferrets.** Unvaccinated dogs, cats, and ferrets exposed to a rabid animal should be euthanized immediately. If the owner is unwilling to have this done, the animal should be placed in strict isolation for 6 months. Isolation in this context refers to confinement in an enclosure that precludes direct contact with people and other animals. Rabies vaccine should be administered upon entry into isolation or 1 month before release to comply with preexposure vaccination recommendations (see Part I.B.1.a.). There are currently no USDA licensed biologics for postexposure prophylaxis of previously unvaccinated domestic animals, and there is evidence that the use of vaccine alone will not reliably prevent the disease in these animals (22). Animals overdue for a booster vaccination need to be evaluated on a case-by-case basis (e.g., severity of exposure, time elapsed since last vaccination, number of previous vaccinations, current health status, and local rabies epidemiology). Dogs, cats, and ferrets that are currently vaccinated should be revaccinated immediately, kept under the owner's control, and observed for

45 days. Any illness in an isolated or confined animal should be reported immediately to the local health department. If signs suggestive of rabies develop, the animal should be euthanized and the head shipped for testing as described in Part I.A.8.

- b. Livestock.** All species of livestock are susceptible to rabies; cattle and horses are the most frequently reported infected species (2). Livestock exposed to a rabid animal and currently vaccinated with a vaccine approved by USDA for that species should be revaccinated immediately and observed for 45 days. Unvaccinated livestock should be euthanized immediately. If the animal is not euthanized it should be kept under close observation for 6 months. Any illness in an animal under observation should be reported immediately to the local health department. If signs suggestive of rabies develop, the animal should be euthanized and the head shipped for testing as described in Part I.A.8.

Handling and consumption of tissues from exposed animals might carry a risk for rabies transmission. Risk factors depend in part on the site(s) of exposure, amount of virus present, severity of wounds, and whether sufficient contaminated tissue has been excised. If an exposed animal is to be slaughtered for consumption, it should be done immediately after exposure, and all tissues should be cooked thoroughly. Persons handling exposed animals, carcasses, and tissues should use barrierprecautions. Historically, federal guidelines for meat inspectors required that any animal known to have been exposed to rabies within 8 months be rejected for slaughter. USDA Food and Inspection Service (FSIS) meat inspectors should be notified if such exposures occur in food animals before slaughter.

Rabies virus might be widely distributed in tissues of infected animals (23). Tissues and products from a rabid animal should not be used for human or animal consumption (24). Pasteurization temperatures will inactivate rabies virus; therefore, inadvertently drinking pasteurized milk or eating thoroughly cooked animal products does not constitute a rabies exposure.

Multiple rabid animals in a herd or herbivore-to-herbivore transmission are uncommon; therefore, restricting the rest of the herd if a single animal has been exposed to or infected by rabies is usually not necessary.

- c. Other Animals.** Other mammals exposed to a rabid animal should be euthanized immediately. Animals

maintained in USDA-licensed research facilities or accredited zoological parks should be evaluated on a case-by-case basis.

6. Management of Animals that Bite Humans.

a. Dogs, Cats, and Ferrets. Rabies virus might be excreted in the saliva of infected dogs, cats, and ferrets during illness and/or for only a few days before illness or death (25–27). A healthy dog, cat, or ferret that bites a person should be confined and observed daily for 10 days (28); administration of rabies vaccine to the animal is not recommended during the observation period to avoid confusing signs of rabies with possible side effects of vaccine administration. Any illness in the animal should be reported immediately to the local health department. Such animals should be evaluated by a veterinarian at the first sign of illness during confinement. If signs suggestive of rabies develop, the animal should be euthanized and the head submitted for testing as described in Part I.A.8. Any stray or unwanted dog, cat, or ferret that bites a person may be euthanized immediately and the head submitted for rabies examination.

b. Other Biting Animals. Other biting animals that might have exposed a person to rabies should be reported immediately to the local health department. Management of animals other than dogs, cats, and ferrets depends on the species, the circumstances of the bite, the epidemiology of rabies in the area, the biting animal's history, current health status, and the animal's potential for exposure to rabies. Previous vaccination of these animals might not preclude the necessity for euthanasia and testing.

7. Outbreak Prevention and Control. The emergence of new rabies virus variants or the introduction of non-indigenous viruses poses a significant risk to humans, domestic animals, and wildlife (29–36). A rapid and comprehensive response includes the following measures:

- a. Characterize the virus at a national or regional reference laboratory.
- b. Identify and control the source of the introduction.
- c. Enhance laboratory-based surveillance in wild and domestic animals.
- d. Increase animal rabies vaccination rates.
- e. Restrict the movement of animals.
- f. Evaluate the need for vector population reduction.
- g. Coordinate a multiagency response.
- h. Provide public and professional outreach and education.

8. Disaster Response. Animals might be displaced during and after man-made or natural disasters and require

emergency sheltering (<http://www.bt.cdc.gov/disasters/petshelters.asp> and <http://www.avma.org/disaster/default.asp>) (37). Animal rabies vaccination and exposure histories often are not available for displaced animals. Disaster response creates situations where animal caretakers might lack appropriate training and preexposure vaccination. In such situations, it is critical to implement and coordinate rabies-prevention and -control measures to reduce the risk of rabies transmission and the need for human postexposure prophylaxis. Such measures include:

- a. Coordinate relief efforts of individuals and organizations with the local emergency operations center before deployment.
- b. Examine each animal at a triage site for signs of rabies.
- c. Isolate animals exhibiting signs of rabies, pending evaluation by a veterinarian.
- d. Ensure that all animals have a unique identifier.
- e. Administer a rabies vaccination to all dogs, cats and ferrets unless reliable proof of vaccination exists.
- f. Adopt minimum standards for animal caretakers that include personal protective equipment, previous rabies vaccination, and appropriate training in animal handling (see Part I.C.).
- g. Maintain documentation of animal disposition and location (e.g., returned to owner, died or euthanized, adopted, relocated to another shelter, and address of new location).
- h. Provide facilities to confine and observe animals involved in exposures (see Part I.A.1.).
- i. Report human exposures to appropriate public health authorities (see Part I.B.6.).

C. Prevention and Control Methods Related to Wildlife.

The public should be warned not to handle or feed wild mammals. Wild mammals and hybrids that bite or otherwise expose persons, pets, or livestock should be considered for euthanasia and rabies examination. A person bitten by any wild mammal should immediately report the incident to a health-care provider who, in consultation with public health authorities, can evaluate the need for postexposure prophylaxis (3).

Translocation of infected wildlife has contributed to the spread of rabies (30–34); therefore, the translocation of known terrestrial rabies reservoir species should be prohibited. Whereas state-regulated wildlife rehabilitators and nuisance wildlife-control operators might play a role in a comprehensive rabies-control program, minimum standards for persons who handle wild mammals should include rabies vaccination, appropriate training, and continuing education.

1. **Carnivores.** The use of licensed oral vaccines for the mass vaccination of free-ranging wildlife should be considered in selected situations, with the approval of the state agency responsible for animal-rabies control (5,38). The distribution of oral rabies vaccine should be based on scientific assessments of the target species and followed by timely and appropriate analysis of surveillance data; such results should be provided to all stakeholders. In addition, parenteral vaccination (trap-vaccinate-release) of wildlife rabies reservoirs may be integrated into coordinated oral rabies vaccination programs to enhance their effectiveness. Continuous and persistent programs for trapping or poisoning wildlife are not effective in reducing wildlife rabies reservoirs on a state-wide basis. However, limited population control in high-contact areas (e.g., picnic grounds, camps, and suburban areas) might be indicated for the removal of selected high-risk species of wildlife (5). State agriculture, public health, and wildlife agencies should be consulted for planning, coordination, and evaluation of vaccination or population-reduction programs.
2. **Bats.** Indigenous rabid bats have been reported from every state except Hawaii and have caused rabies in at least 40 humans in the United States (39–46). Bats should be excluded from houses, public buildings, and adjacent structures to prevent direct association with humans (47,48). Such structures should then be made bat-proof by sealing entrances used by bats. Controlling rabies in bats through programs designed to reduce bat populations is neither feasible nor desirable.

Part II: Recommendations for Parenteral Rabies Vaccination Procedures

- A. **Vaccine Administration.** All animal rabies vaccines should be restricted to use by or under the direct supervision of a veterinarian (49), except as recommended in Part I.B.1. All vaccines must be administered in accordance with the specifications of the product label or package insert.
- B. **Vaccine Selection.** Part III lists all vaccines licensed by USDA and marketed in the United States at the time of publication. New vaccine approvals or changes in label specifications made subsequent to publication should be considered as part of this list. Any of the listed vaccines can be used for revaccination, even if the product is not the same as previously administered. Vaccines used in state and local rabies-control programs should have at least a 3-year duration of immunity. This constitutes the most effective method of increasing the proportion of immunized dogs and cats in any population (50). No laboratory or epidemiologic data exist to support the annual or biennial administration of 3- or 4-year vaccines following the initial series.
- C. **Adverse Events.** Currently, no epidemiologic association exists between a particular licensed vaccine product and adverse events (51,52). Adverse events, including rabies in a previously vaccinated animal, should be reported to the vaccine manufacturer and to USDA, Animal and Plant Health Inspection Service, Center for Veterinary Biologics (Internet: http://www.aphis.usda.gov/animal_health/vet_biologics/vb_adverse_event.shtml; telephone: 800-752-6255; or e-mail: cvb@usda.gov).
- D. **Wildlife and Hybrid Animal Vaccination.** The safety and efficacy of parenteral rabies vaccination of wildlife and hybrids have not been established, and no rabies vaccines are licensed for these animals. Parenteral vaccination (trap-vaccinate-release) of wildlife rabies reservoirs can be integrated into coordinated oral rabies vaccination programs as described in Part I.C.1. to enhance their effectiveness. Zoos or research institutions may establish vaccination programs to attempt to protect valuable animals, but these should not replace appropriate public health activities that protect humans (see Part I.B.1.c.2) (9).
- E. **Accidental Human Exposure to Vaccine.** Human exposure to parenteral animal rabies vaccines listed in Part III does not constitute a risk for rabies virus infection. Human exposure to vaccinia-vectored oral rabies vaccines should be reported to state health officials (53).
- F. **Rabies Certificate.** All agencies and veterinarians should use NASPHV Form 51 (revised 2007), Rabies Vaccination Certificate, or an equivalent. This form can be obtained from vaccine manufacturers, NASPHV (<http://www.nasphv.org/Documents/RabiesVacCert.pdf>), or CDC (<http://www.cdc.gov/rabies>). The form must be completed in full and signed by the administering or supervising veterinarian. Computer-generated forms containing the same information are also acceptable.

Part III: Rabies vaccines licensed and marketed in the United States, 2008

Product name	Produced by	License no.	Marketed by	For use in	Dosage (mL)	Age at primary vaccination*	Booster recommended	Route of inoculation
A) MONOVALENT (inactivated)								
RABVAC 1	Fort Dodge Animal Health	112	Fort Dodge Animal Health	Dogs	1	3 mos [†]	Annually	IM [§] or SC [¶]
RABVAC 3	Fort Dodge Animal Health	112	Fort Dodge Animal Health	Cats	1	3 mos	Annually	IM or SC
				Dogs	1	3 mos	1 year later and triennially	IM or SC
				Cats	1	3 mos	1 year later and triennially	IM or SC
RABVAC 3 TF	Fort Dodge Animal Health	112	Fort Dodge Animal Health	Horses	2	3 mos	Annually	IM
				Dogs	1	3 mos	1 year later and triennially	IM or SC
				Cats	1	3 mos	1 year later and triennially	IM or SC
DEFENSOR 1	Pfizer, Inc.	189	Pfizer, Inc.	Horses	2	3 mos	Annually	IM
				Dogs	1	3 mos	Annually	IM or SC
				Cats	1	3 mos	Annually	SC
DEFENSOR 3	Pfizer, Inc.	189	Pfizer, Inc.	Dogs	1	3 mos	1 year later and triennially	IM or SC
				Cats	1	3 mos	1 year later and triennially	SC
				Sheep	2	3 mos	Annually	IM
RABDOMUN	Pfizer, Inc.	189	Schering-Plough	Cattle	2	3 mos	Annually	IM
				Dogs	1	3 mos	1 year later and triennially	IM or SC
				Cats	1	3 mos	1 year later and triennially	SC
RABDOMUN 1	Pfizer, Inc.	189	Schering-Plough	Sheep	2	3 mos	Annually	IM
				Cattle	2	3 mos	Annually	IM
				Dogs	1	3 mos	Annually	IM or SC
CONTINUUM RABIES	Intervet, Inc.	286	Intervet, Inc.	Cats	1	3 mos	Annually	SC
				Dogs	1	3 mos	1 year later and triennially	SC
				Cats	1	3 mos	1 year later and quadrennially	SC
PRORAB-1	Intervet, Inc.	286	Intervet, Inc.	Dogs	1	3 mos	Annually	IM or SC
				Cats	1	3 mos	Annually	IM or SC
				Sheep	2	3 mos	Annually	IM
IMRAB 1	Merial, Inc.	298	Merial, Inc.	Dogs	1	3 mos	Annually	SC
				Cats	1	3 mos	Annually	SC
				Dogs	1	3 mos	Annually	SC
IMRAB 1 TF	Merial, Inc.	298	Merial, Inc.	Cats	1	3 mos	Annually	SC
				Dogs	1	3 mos	Annually	SC
				Cats	1	3 mos	Annually	SC
IMRAB 3	Merial, Inc.	298	Merial, Inc.	Dogs	1	3 mos	1 year later and triennially	IM or SC
				Cats	1	3 mos	1 year later and triennially	IM or SC
				Sheep	2	3 mos	1 year later and triennially	IM or SC
IMRAB 3 TF	Merial, Inc.	298	Merial, Inc.	Cattle	2	3 mos	Annually	IM or SC
				Horses	2	3 mos	Annually	IM or SC
				Ferrets	1	3 mos	Annually	SC
IMRAB Large Animal	Merial, Inc.	298	Merial, Inc.	Dogs	1	3 mos	1 year later and triennially	IM or SC
				Cats	1	3 mos	1 year later and triennially	IM or SC
				Ferrets	1	3 mos	Annually	SC
PUREVAX Feline Rabies	Merial, Inc.	298	Merial, Inc.	Cattle	2	3 mos	Annually	IM or SC
				Horses	2	3 mos	Annually	IM or SC
				Sheep	2	3 mos	1 year later and triennially	IM or SC
B) MONOVALENT (Rabies glycoprotein, live canary pox vector)								
PUREVAX Feline Rabies	Merial, Inc.	298	Merial, Inc.	Cats	1	8 wks	Annually	SC
C) COMBINATION (Inactivated rabies)								
CONTINUUM DAP-R	Intervet, Inc.	286	Intervet, Inc.	Dogs	1	3 mos	1 year later and triennially	SC
CONTINUUM Feline HCP-R	Intervet, Inc.	286	Intervet, Inc.	Cats	1	3 mos	1 year later and quadrennially**	SC
Equine POTOMAVAC + IMRAB	Merial, Inc.	298	Merial, Inc.	Horses	1	3 mos	Annually	IM
D) COMBINATION (Rabies glycoprotein, live canary pox vector)								
PUREVAX Feline 3/Rabies	Merial, Inc.	298	Merial, Inc.	Cats	1	8 wks	Annually	SC
PUREVAX Feline 4/Rabies	Merial, Inc.	298	Merial, Inc.	Cats	1	8 wks	Annually	SC
E) ORAL (Rabies glycoprotein, live vaccinia vector) — RESTRICTED TO USE IN STATE AND FEDERAL RABIES-CONTROL PROGRAMS								
RABORAL V-RG	Merial, Inc.	298	Merial, Inc.	Coyotes Raccoons	N/A ^{††}	N/A	As determined by local authorities	Oral

* Minimum age (or older) and revaccinated 1 year later.

† One month = 28 days.

§ Intramuscularly.

¶ Subcutaneously.

** Nonrabies fractions have a 3-year duration (see label).

†† Not applicable.

Rabies vaccine manufacturer contact information

Manufacturer	Phone number	Internet address
Fort Dodge Animal Health	800-533-8536	http://www.wyeth.com/divisions/fort_dodge.asp
Intervet, Inc.	800-835-0541	http://www.intervetusa.com
Merial, Inc.	888-637-4251	http://us.merial.com
Pfizer, Inc.	800-366-5288	http://www.pfizerah.com
Schering-Plough Animal Health	800-521-5767	http://www.spah.com/usa

Note: ADVERSE EVENTS: Adverse events should be reported to the vaccine manufacturer and to U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Center for Veterinary Biologics (Internet: http://www.aphis.usda.gov/animal_health/vet_biologics/vb_adverse_event.shtml); telephone: 800-752-6255; or e-mail: cvb@usda.gov).

References

- Rabies. In: Heymann D, ed. Control of communicable diseases manual. 18th ed. Washington, DC: American Public Health Association; 2004:438–47.
- Blanton JD, Hanlon CA, Rupprecht CE. Rabies surveillance in the United States during 2006. *J Am Vet Med Assoc* 2007;231:540–56. Available at <http://www.cdc.gov/rabies/publications>.
- CDC. Human rabies prevention—United States, 1999. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1999;48:(No. RR-1). Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/00056176.htm>.
- McQuiston J, Yager PA, Smith JS, Rupprecht CE. Epidemiologic characteristics of rabies virus variants in dogs and cats in the United States, 1999. *J Am Vet Med Assoc* 2001;218:1939–42.
- Hanlon CA, Childs JE, Nettles VF, et al. Recommendations of the Working Group on Rabies. Article III: rabies in wildlife. *J Am Vet Med Assoc* 1999;215:1612–8.
- Hanlon CA, Smith JS, Anderson GR, et al. Recommendations of the Working Group on Rabies. Article II: laboratory diagnosis of rabies. *J Am Vet Med Assoc* 1999;215:1444–6.
- Rudd RJ, Smith JS, Yager PA, et al. A need for standardized rabies-virus diagnostic procedures: effect of cover-glass mountant on the reliability of antigen detection by the fluorescent antibody test. *Virus Res* 2005;111:83–8.
- American Veterinary Medical Association. AVMA guidelines on euthanasia, June 2007. Schaumburg, IL: American Veterinary Medical Association; 2007. Available at http://www.avma.org/issues/animal_welfare/euthanasia.pdf.
- Tizard I, Ni Y. Use of serologic testing to assess immune status of companion animals. *J Am Vet Med Assoc* 1998;213:54–60.
- Greene CE, ed. Rabies and other lyssavirus infections. In: Infectious diseases of the dog and cat. 3rd ed. London, England: Saunders Elsevier; 2006:167–83.
- Rupprecht CE, Gilbert J, Pitts R, Marshall K, Koprowski H. Evaluation of an inactivated rabies virus vaccine in domestic ferrets. *J Am Vet Med Assoc* 1990;196:1614–6.
- Aubert MF. Practical significance of rabies antibodies in cats and dogs. *Rev Sci Tech* 1992;11:735–60.
- Cliquet F, Verdier Y, Sagné L, et al. Neutralising antibody titration in 25,000 sera of dogs and cats vaccinated against rabies in France, in the framework of the new regulations that offer an alternative to quarantine. *Rev Sci Tech* 2003;22:857–66.
- National Association of State Public Health Veterinarians. Compendium of measures to prevent disease and injury associated with animals in public settings, 2006. Available at <http://www.nasphv.org>.
- Bender J, Schulman S. Reports of zoonotic disease outbreaks associated with animal exhibits and availability of recommendations for preventing zoonotic disease transmission from animals to people in such settings. *J Am Vet Med Assoc* 2004;224:1105–9.
- American Veterinary Medical Association. Private ownership of wild animals. Schaumburg, IL: American Veterinary Medical Association. Available at http://www.avma.org/issues/policy/wild_animal_ownership.asp.
- Position on canine hybrids. In: Directory and resource manual. Schaumburg, IL: American Veterinary Medical Association; 2002:88–9.
- Siino BS. Crossing the line. *American Society for the Prevention of Cruelty to Animals, Animal Watch*; 2000:22–9.
- Jay MT, Reilly KF, DeBess EE, Haynes EH, Bader DR, Barrett LR. Rabies in a vaccinated wolf-dog hybrid. *J Am Vet Med Assoc* 1994;205:1729–32.
- CDC. An imported case of rabies in an immunized dog. *MMWR* 1987;36:94–6,101. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/00000874.htm>.
- CDC. Imported dog and cat rabies—New Hampshire, California. *MMWR* 1988;37:559–60. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/00001275.htm>.
- Hanlon CA, Niezgoda MN, Rupprecht CE. Postexposure prophylaxis for prevention of rabies in dogs. *Am J Vet Res* 2002;63:1096–100.
- Charlton, KM. The pathogenesis of rabies and other lyssaviral infections: recent studies. *Curr Top Microbiol Immunol* 1994;187:95–119.
- CDC. Mass treatment of humans who drank unpasteurized milk from rabid cows—Massachusetts, 1996–1998. *MMWR* 1999;48:228–9. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/00056759.htm>.
- Vaughn JB, Gerhardt P, Paterson J. Excretion of street rabies virus in saliva of cats. *J Am Med Assoc* 1963;184:705.
- Vaughn JB, Gerhardt P, Newell KW. Excretion of street rabies virus in saliva of dogs. *J Am Med Assoc* 1965;193:363–8.
- Niezgoda M, Briggs DJ, Shaddock J, Rupprecht CE. Viral excretion in domestic ferrets (*Mustela putorius furo*) inoculated with a raccoon rabies isolate. *Am J Vet Res* 1998;59:1629–32.
- Tepsumethanon V, Lumlerdacha B, Mitmoonpitak C, Sitprija V, Meslin FX, Wilde H. Survival of naturally infected rabid dogs and cats. *Clin Infect Dis* 2004;39:278–80.
- Jenkins SR, Perry BD, Winkler WG. Ecology and epidemiology of raccoon rabies. *Rev Infect Dis* 1988;10(Suppl 4):S620–5.
- CDC. Translocation of coyote rabies—Florida, 1994. *MMWR* 1995;44:580–7. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/00038451.htm>.
- Rupprecht CE, Smith JS, Fekadu M, Childs JE. The ascension of wildlife rabies: a cause for public health concern or intervention? *Emerg Infect Dis* 1995;1:107–14. Available at <http://www.cdc.gov/ncidod/eid/vol1no4/rupprech.htm#top>.
- Constantine DG. Geographic translocation of bats: known and potential problems. *Emerg Infect Dis* 2003;9:17–21. Available at <http://www.cdc.gov/ncidod/EID/vol9no1/02-0104.htm>.
- Krebs JW, Strine TW, Smith JS, Rupprecht CE, Childs JE. Rabies surveillance in the United States during 1993. *J Am Vet Med Assoc* 1994;205:1695–709.
- VF Nettles, JH Shaddock, RK Sikes, CR Reyes. Rabies in translocated raccoons. *Am J Public Health* 1979;69:601–2.
- RM Engeman, KL Christensen, MJ Pipas, DL Bergman. Population monitoring in support of a rabies vaccination program for skunks in Arizona. *J Wildl Dis* 2003;39:746–50.
- Leslie MJ, Messenger S, Rohde RE, et al. Bat-associated rabies virus in skunks. *Emerg Infect Dis* 2006;12:1274–7. Available at <http://www.cdc.gov/ncidod/EID/vol12no08/05-1526.htm>.
- The Humane Society of the United States. HSUS Disaster Services. Available at http://www.hsus.org/hsus_field/hsus_disaster_center.
- Slate D, Rupprecht CE, Rooney JA, Donovan D, Lein DH, Chipman RB. Status of oral rabies vaccination in wild carnivores in the United States. *Virus Res* 2005;111:68–76.
- Messenger SL, Smith JS, Rupprecht CE. Emerging epidemiology of bat-associated cryptic cases of rabies in humans in the United States. *Clin Infect Dis* 2002;35:738–47.

40. CDC. Human rabies—California, 2002. *MMWR* 2002;51:686–8. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5131a4.htm>.
41. CDC. Human rabies—Tennessee, 2002. *MMWR* 2002;51:828–9. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5137a2.htm>.
42. CDC. Human rabies—Iowa, 2002. *MMWR* 2003;52:47–8. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5203a3.htm>.
43. CDC. Human death associated with bat rabies—California, 2003. *MMWR* 2004;53:33–5. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5302a4.htm>.
44. CDC. Recovery of a patient from clinical rabies, Wisconsin, 2004. *MMWR* 2004;53:1171–3. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5350a1.htm>.
45. CDC. Human rabies—Mississippi, 2005. *MMWR* 2006;55:207–8. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5508a4.htm>.
46. CDC. Human rabies—Indiana and California, 2006. *MMWR* 2007;56:361–5. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5615a1.htm>.
47. Frantz SC, Trimarchi CV. Bats in human dwellings: health concerns and management. In: Decker DF, ed. *Proceedings of the 1st Eastern Wildlife Damage Control Conference*. Ithaca, NY: Cornell University Press; 1983:299–308.
48. Greenhall AM. House bat management. US Fish and Wildlife Service, Resource Publication 143;1982.
49. American Veterinary Medical Association. Model rabies control ordinance. Schaumburg, IL: American Veterinary Medical Association. Available at http://www.avma.org/issues/policy/rabies_control.asp.
50. Bunn TO. Canine and feline vaccines, past and present. In Baer GM, ed. *The natural history of rabies*. 2nd ed. Boca Raton, FL: CRC Press; 1991:415–25.
51. Gobar GM, Kass PH. World wide web-based survey of vaccination practices, postvaccinal reactions, and vaccine site-associated sarcomas in cats. *J Am Vet Med Assoc* 2002;220:1477–82.
52. Macy DW, Hendrick MJ. The potential role of inflammation in the development of postvaccinal sarcomas in cats. *Vet Clin North Am Small Anim Pract* 1996;26:103–9.
53. Rupprecht CE, Blass L, Smith K, et al. Human infection due to recombinant vaccinia-rabies glycoprotein virus. *N Engl J Med* 2001;345:582–6.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's Internet server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. Paper copy subscriptions are available through the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Data are compiled in the National Center for Public Health Informatics, Division of Integrated Surveillance Systems and Services. Address all inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333 or to www.mmwrq@cdc.gov.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.