

Notifiable Diseases and Mortality Tables

TABLE I. Provisional cases of selected* infrequently reported notifiable diseases (<1,000 cases reported during the preceding year) — United States, week ending December 19, 2015 (50th week)[†] (Export data)

Disease	Current week	Cum 2015	5-year weekly average [§]	Total cases reported for previous years					States reporting cases during current week (No.)
				2014	2013	2012	2011	2010	
Anthrax	—	—	—	—	—	—	1	—	
Arboviral diseases** ^{††} :									
California serogroup virus disease	—	56	—	96	95	81	137	75	
Chikungunya virus disease [¶]	NN	NN	—	NN	NN	NN	NN	NN	
Eastern equine encephalitis virus disease	—	3	—	8	8	15	4	10	
Powassan virus disease	—	5	0	8	12	7	16	8	
St. Louis encephalitis virus disease	—	19	0	10	1	3	6	10	
Western equine encephalitis virus disease	—	—	—	—	—	—	—	—	
Botulism, total	3	174	4	161	152	168	153	112	
foodborne	—	40	1	15	4	27	24	7	
infant	3	115	3	127	136	123	97	80	NYC (1), OH (2)
other (wound and unspecified)	—	19	0	19	12	18	32	25	
Brucellosis	1	112	2	92	99	114	79	115	FL (1)
Chancroid	1	17	0	6	—	15	8	24	WY (1)
Cholera	—	1	1	5	14	17	40	13	
Cyclosporiasis ^{††}	—	605	2	388	784	123	151	179	
Diphtheria	—	—	—	1	—	1	—	—	
<i>Haemophilus influenzae</i> invasive disease (age <5 yrs) ^{§§} :									
serotype b	—	21	1	40	31	30	14	23	
nonsertotype b	4	219	5	228	222	205	145	200	OH (1), SC (1), CO (1), AZ (1)
unknown serotype	1	200	5	204	185	210	226	223	AZ (1)
Hansen disease ^{††}	1	63	2	88	81	82	82	98	FL (1)
Hantavirus infections ^{††} :									
Hantavirus infection (non-HPS) ^{¶¶}	NN	NN	—	NN	NN	NN	NN	NN	
Hantavirus Pulmonary Syndrome (HPS)	—	10	0	32	21	30	23	20	
Hemolytic uremic syndrome, postdiarrheal ^{††}	—	224	4	250	329	274	290	266	
Hepatitis B, virus infection perinatal	—	24	1	47	48	40	NP	NP	
Influenza-associated pediatric mortality ^{††,¶¶}	1	118	2	141	160	52	118	61	TN (1)
Leptospirosis ^{††}	—	27	0	38	NN	NN	NN	NN	
Listeriosis	4	649	15	769	735	727	870	821	PA (1), AL (1), OK (1), TX (1)
Measles ^{***}	—	188	1	667	187	55	220	63	
Meningococcal disease, invasive ^{†††} :									
serogroup ACWY	—	76	5	123	142	161	257	280	
serogroup B	—	79	3	89	99	110	159	135	
other serogroup	—	18	1	25	17	20	20	12	
unknown serogroup	2	164	7	196	298	260	323	406	FL (1), OR (1)
Novel influenza A virus infections ^{§§§}	—	6	1	3	21	313	14	4	
Plague	—	9	—	10	4	4	3	2	
Poliomyelitis, paralytic	—	—	—	—	1	—	—	—	
Polio virus infection, nonparalytic ^{††}	—	—	—	—	—	—	—	—	
Psittacosis ^{††}	—	7	0	8	6	2	2	4	
Q fever, total ^{††}	2	127	4	168	170	135	134	131	
acute	2	109	3	132	137	113	110	106	PA (1), TN (1)
chronic	—	18	1	36	33	22	24	25	
Rabies, human	—	1	0	1	2	1	6	2	
SARS-CoV ^{††}	—	—	—	—	—	—	—	—	
Smallpox ^{††}	—	—	—	—	—	—	—	—	
Streptococcal toxic-shock syndrome ^{††}	2	239	4	259	224	194	168	142	NY (1), OH (1)
Syphilis, congenital (age <1 yr) ^{††,¶¶¶}	—	333	7	458	348	322	360	377	
Toxic-shock syndrome (staphylococcal) ^{††}	—	98	1	59	71	65	78	82	
Trichinellosis ^{††}	—	7	0	14	22	18	15	7	
Tularemia	—	240	2	180	203	149	166	124	
Typhoid fever	4	305	7	349	338	354	390	467	NYC (3), MD (1)
Vancomycin-intermediate <i>Staphylococcus aureus</i> ^{††}	2	152	4	212	248	134	82	91	NY (1), LA (1)
Vancomycin-resistant <i>Staphylococcus aureus</i> ^{††}	—	1	—	—	—	2	—	2	

See Table I footnotes on next page.

