

Elevated Blood Lead Levels Among Employed Adults — United States, 1994–2013

Walter A. Alarcon, MD¹

State Adult Blood Lead Epidemiology and Surveillance (ABLES) Program Investigators

¹National Institute for Occupational Safety and Health, CDC

Preface

CDC's National Institute for Occupational Safety and Health (NIOSH) and state health departments collect data on laboratory-reported adult blood lead levels (BLLs). This report presents data on elevated BLLs among employed adults (defined as persons aged ≥ 16 years) in the United States for 1994–2013. This report is a part of the *Summary of Notifiable Noninfectious Conditions and Disease Outbreaks — United States*, which encompasses various surveillance years but is being published in 2016 (1). The *Summary of Notifiable Noninfectious Conditions and Disease Outbreaks* appears in the same volume of the Morbidity Mortality Weekly Report (MMWR) as the annual *Summary of Notifiable Infectious Diseases* (2).

Background

Since 1987, NIOSH and state health departments have maintained the Adult Blood Lead Epidemiology and Surveillance (ABLES) Program, a state-based surveillance program of laboratory-reported adult BLLs (3). The BLL is an often-used estimate of recent external exposure to lead (4,5). This report summarizes data on elevated BLLs among employed adults during January 1, 1994–December 31, 2013.

Information is provided by geographic division and reporting state, for “all cases” reported by a state (these include cases among adult residents in the reporting state plus cases identified by the reporting state but occurring among persons who reside in another state) and “state-residents” only, by exposure source, for BLLs ≥ 10 $\mu\text{g}/\text{dL}$ (definition of elevated BLL from 2009 until 2014) (3,6–8), and for BLLs ≥ 25 $\mu\text{g}/\text{dL}$ (previous definition of elevated BLL) (9). The current case definition (BLL ≥ 5 $\mu\text{g}/\text{dL}$) was adopted in 2015 and became effective in 2016, on the basis of mounting evidence for adverse health outcomes among adults with BLLs between 5 $\mu\text{g}/\text{dL}$ and 25 $\mu\text{g}/\text{dL}$ (4,5). State prevalence rates of elevated BLLs (≥ 10 $\mu\text{g}/\text{dL}$) for 2013 are categorized into two groups (above

or below the national prevalence rate) (Figure 1). Trends of national prevalence rates of BLLs ≥ 10 $\mu\text{g}/\text{dL}$ and BLLs ≥ 25 $\mu\text{g}/\text{dL}$ from 1994 to 2013 are provided (Figure 2).


ABLES is the only program conducting nationwide adult lead exposure surveillance. It has provided the occupational safety and health community with essential information for setting research and intervention priorities. ABLES' impact is achieved through its longstanding strategic partnerships with state ABLES programs, federal agencies, and worker-affiliated organizations. For example, in 2008, the Occupational Safety and Health Administration (OSHA) updated its National Lead Emphasis Program to reduce occupational lead exposure by targeting unsafe conditions and high-hazard industries (10). To accomplish this objective, OSHA used national ABLES data to identify industries whose employees exhibit high BLLs. OSHA has agreements with state ABLES programs to use their lead exposure data to target workplace inspections.

Although federal funding for state ABLES programs was discontinued in September 2013, a total of 30 states continue to collaborate with NIOSH (down from a peak of 41 states) to provide data. In August 2015, funding to support adult BLL surveillance was resumed at a reduced level. To sustain lead exposure surveillance and prevention activities, state ABLES programs share resources with two other CDC programs: the Childhood Lead Poisoning Prevention Program and the Environmental Public Health Tracking Program. Since September 2013, NIOSH has continued to provide technical assistance to states with adult blood lead surveillance programs and maintains the ABLES website for reporting ABLES findings.

The BLL is a direct index of a worker's exposure to lead as well as an indication of the potential for adverse effects from that exposure (4,5). The half-life of lead in blood is approximately 40 days in males (11), so the BLL is an estimate primarily of recent exposure to lead. Because lead accumulates in bone and BLL is in equilibrium with bone lead, the BLL might be elevated in some persons who have not had recent exposure to lead. Because this equilibrium can lead to persistent BLL elevations, the public health burden of elevated BLLs in adults is measured as prevalence. In contrast, the public health burden of elevated BLLs in children aged < 3 years is measured as incidence because these young children have little lead

Corresponding author: Walter Alarcon, National Institute for Occupational Safety and Health, CDC. Telephone: 513-841-4451; E-mail: wda7@cdc.gov.

FIGURE 2. National prevalence rate* of reported cases of elevated blood lead levels,[†] by year — State Adult Blood Epidemiology and Surveillance Programs, United States, 1994–2013[§]



Abbreviation: BLL = blood lead level.

* Per 100,000 employed adults aged ≥ 16 years. Denominator data extracted from 2015 U.S. Department of Labor, Bureau of Labor Statistics Local Area Unemployment Statistics (LAUS) program (<http://www.bls.gov/lau/staadata.txt>).

[†] Since 2009, the case definition for an elevated blood lead level is a BLL ≥ 10 $\mu\text{g/dL}$. For historical comparisons, prevalence rates at the previous case definition (BLL ≥ 25 $\mu\text{g/dL}$) are provided.

[§] A total of 30 states submitted data in 2013 (down from 41 states in 2012): Alabama, Alaska, Arizona, California, Colorado, Connecticut, Florida, Georgia, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Nebraska, New Jersey, New Mexico, New York, North Carolina, Oklahoma, Oregon, Pennsylvania, Vermont, Washington, Wisconsin, Wyoming. Massachusetts provided data for BLLs ≥ 25 $\mu\text{g/dL}$. For 2013, the first number is the number of states reporting BLLs ≥ 25 $\mu\text{g/dL}$ (i.e., 30 states in 2013), and the second number is the number of states reporting BLLs ≥ 10 $\mu\text{g/dL}$ (i.e., 29 states in 2013).

2010 forward. Prevalence rates of BLLs ≥ 25 $\mu\text{g/dL}$ are a subset of prevalence rates of BLLs ≥ 10 $\mu\text{g/dL}$. In the United States, most lead exposures among adults are occupational (9). A total of 29 states submitted work-relatedness information in 2013. Prevalence rate differences across states could reflect improved compliance with required OSHA monitoring in some states.

These counts and rates of elevated BLLs must be considered minimum estimates of the actual magnitude of the problem of lead exposures in the United States. This is for multiple reasons:

- not all states participate in the ABLES program;
- not all employers provide BLL testing to lead-exposed workers as required by OSHA regulations;
- not all nonoccupationally exposed adults are tested; and
- some laboratories might not report all tests as required by state laws or regulations.

For specific explanations, interpretation, and possible updates on data for any individual state, the state ABLES program investigator should be contacted directly. Contact information is available at <http://www.cdc.gov/niosh/topics/ABLES/state.html>.

Methods for Identifying Elevated BLLs Among Employed Adults

Beginning in 2016, a nationally reportable case of an employed adult with an elevated BLL is defined as a case in an employed person aged ≥ 16 years at the time of blood collection with a venous blood lead level ≥ 5 $\mu\text{g/dL}$ of whole blood. The standardized diagnostic test is the BLL test using a venous blood sample. All participating state health departments have a requirement for laboratories and/or health care providers

to report laboratory blood lead results to the state health department. However, this requirement varies among ABLES states, ranging from the reporting of all BLLs to reporting only BLLs ≥ 40 $\mu\text{g}/\text{dL}$ (3). The ABLES program ultimately aims to establish a national database for all BLL tests among adults and encourages all states to share information with NIOSH.

Publication Criteria

Cases meet the publication criteria if the employed adult (aged ≥ 16 years) had a venous BLL ≥ 25 $\mu\text{g}/\text{dL}$ during 1994–2013 or a venous BLL ≥ 10 $\mu\text{g}/\text{dL}$ during 2010–2013. When an adult had multiple blood lead tests in a given year, only the highest BLL for that adult in that year was counted. Prevalence rates of BLLs ≥ 25 $\mu\text{g}/\text{dL}$ are a subset of prevalence rates of BLLs ≥ 10 $\mu\text{g}/\text{dL}$ and are included for historic comparison.

Highlights

In 2013, the prevalence rate of BLLs ≥ 10 $\mu\text{g}/\text{dL}$ was 20.4 adults per 100,000 employed population, calculated from 29 reporting states. In 2013, a total of 30 states submitted data on 5,504 adults with BLLs ≥ 25 $\mu\text{g}/\text{dL}$, and 29 states submitted data on 20,880 adults with BLLs ≥ 10 $\mu\text{g}/\text{dL}$ (Table 1). A total of 23 states submitted individual level data, and seven states submitted count data only. Overall, the national prevalence rate of BLLs ≥ 10 $\mu\text{g}/\text{dL}$ declined from 26.6 adults per 100,000 employed in 2010 (among 37 states) to 20.4 in 2013 (among 29 reporting states). In 2013, of the 29 reporting states, 12 had prevalence rates of BLLs ≥ 10 $\mu\text{g}/\text{dL}$ equal to or above the national prevalence rate (20.4/100,000) (Figure 1). The national prevalence rate of BLLs ≥ 25 $\mu\text{g}/\text{dL}$ among state residents and nonresidents declined from 14.0 adults per

100,000 employed in 1994 (among 17 states) to 5.2 in 2013 (among 30 states).

Historically, in the United States, most lead exposures among adults have been occupational. In 2013, a total of 29 states submitted data on 5,491 adults with BLLs ≥ 25 $\mu\text{g}/\text{dL}$ of which 944 (17.2%) had no known exposure history (Table 2). Among the 4,547 adults with known exposure, 93.7% had occupational exposure, ranging from 42.9% to 100% among reporting states. Individual level data on 2,313 occupational cases with BLLs ≥ 25 $\mu\text{g}/\text{dL}$ were available from 22 states. The majority of these adults were employed in four main industry sectors: manufacturing (n = 1,227 [53.1%]), construction (n = 468 [20.2%]), services (n = 194 [8.4%]), and mining (n = 182 [7.9%]). Within manufacturing, the majority of cases (n = 878; 71.6%) were among workers employed in storage battery manufacturing (North American Industry Classification System [NAICS] 33591), alumina and aluminum production and processing (NAICS 33131), and nonferrous metal (except copper and aluminum) rolling, drawing, extruding, and alloying (NAICS 33149) industries. Within construction, the majority of cases (n = 329 [70.3%]) were among workers employed in painting and wall covering contractors (NAICS 23832); highway, street, and bridge construction (NAICS 23731); and residential building construction (NAICS 23611) industries. Within the services sector, the majority of cases (n = 128 [66%]) were among workers employed in remediation services (NAICS 56291); all other amusement and recreation industries (NAICS 71399); automotive, mechanical, and electrical repair and maintenance (NAICS 81111); and fitness and recreational sports centers (NAICS 71394). Copper, nickel, lead, and zinc mining (NAICS 21223) accounted for 98.9% of the mining cases.

TABLE 1. Reported numbers of cases and prevalence rates of adults* with blood lead levels $\geq 10 \mu\text{g}/\text{dL}$ and blood lead levels $\geq 25 \mu\text{g}/\text{dL}$, by geographic division and area — state Adult Blood Lead Epidemiology and Surveillance programs, United States, 2013[†]

Division/State	No. of employed state-resident adults (in 1,000s)	Blood lead levels $\geq 10 \mu\text{g}/\text{dL}$				Blood lead levels $\geq 25 \mu\text{g}/\text{dL}$ [§]			
		All cases [¶]		State residents ^{**}		All cases		State residents	
		No.	(Rate)	No.	(Rate)	No.	(Rate)	No.	(Rate)
Total	105,474	20,880	(20.4)	19,603	(19.2)	5,504	(5.2)	5,183	(4.9)
New England									
Connecticut	1,724	331	(19.2)	313	(18.2)	62	(3.6)	61	(3.5)
Massachusetts	3,272	— ^{††}	(—)	—	(—)	126	(3.9)	105	(3.2)
Vermont	336	47	(14.0)	47	(14.0)	12	(3.6)	12	(3.6)
Mid Atlantic									
New Jersey	4,164	832	(20.0)	832	(20.0)	158	(3.8)	158	(3.8)
New York	8,891	1,873	(21.1)	1,731	(19.5)	295	(3.3)	270	(3.0)
Pennsylvania	5,964	2,928	(49.1)	2,915	(48.9)	1,533	(25.7)	1,527	(25.6)
East North Central									
Illinois	5,961	1,279	(21.5)	1,253	(21.0)	283	(4.7)	279	(4.7)
Indiana	2,947	596	(20.2)	596	(20.2)	113	(3.8)	113	(3.8)
Michigan	4,306	596	(13.8)	595	(13.8)	108	(2.5)	108	(2.5)
Wisconsin	2,877	687	(23.9)	686	(23.8)	105	(3.7)	105	(3.7)
West North Central									
Iowa	1,594	856	(53.7)	856	(53.7)	202	(12.7)	202	(12.7)
Minnesota	2,819	598	(21.2)	598	(21.2)	107	(3.8)	107	(3.8)
Missouri	2,814	3,145	(111.8)	2,835	(100.8)	690	(24.5)	613	(21.8)
Nebraska	983	195	(19.8)	195	(19.8)	32	(3.3)	32	(3.3)
South Atlantic									
Florida	8,783	888	(10.1)	863	(9.8)	270	(3.1)	266	(3.0)
Georgia	4,368	898	(20.6)	897	(20.5)	237	(5.4)	237	(5.4)
Maryland	2,917	275	(9.4)	234	(8.0)	75	(2.6)	62	(2.1)
North Carolina	4,310	219	(5.1)	218	(5.1)	99	(2.3)	99	(2.3)
East South Central									
Alabama	2,012	928	(46.1)	548	(27.2)	433	(21.5)	299	(14.9)
Kentucky	1,892	478	(25.3)	468	(24.7)	94	(5.0)	92	(4.9)
West South Central									
Louisiana	1,965	380	(19.3)	380	(19.3)	92	(4.7)	92	(4.7)
Oklahoma	1,707	144	(8.4)	121	(7.1)	29	(1.7)	27	(1.6)
Mountain									
Arizona ^{§§}	2,804	178	(6.3)	178	(6.3)	20	(0.7)	20	(0.7)
Colorado	2,591	103	(4.0)	41	(1.6)	29	(1.1)	15	(0.6)
New Mexico	859	48	(5.6)	48	(5.6)	13	(1.5)	13	(1.5)
Wyoming	292	66	(22.6)	66	(22.6)	12	(4.1)	12	(4.1)
Pacific									
Alaska	340	123	(36.1)	62	(18.2)	8	(2.4)	6	(1.8)
California	17,003	1,825	(10.7)	1,790	(10.5)	192	(1.1)	191	(1.1)
Oregon	1,761	92	(5.2)	79	(4.5)	12	(0.7)	9	(0.5)
Washington	3,217	272	(8.5)	158	(4.9)	63	(2.0)	51	(1.6)

* An employed person aged ≥ 16 years at the time of blood collection. When an adult had multiple blood lead tests in a given year, only the highest blood lead level for that adult in that year was counted. Rate per 100,000 employed adults. Data from the Adult Blood Epidemiology and Surveillance (ABLES) Program, National Institute for Occupational Safety and Health (NIOSH/CDC). Denominators extracted from 2015 U.S. Department of Labor, Bureau of Labor Statistics, Local Area Unemployment Statistics (LAUS) program (<http://www.bls.gov/lau/staadata.txt>).

[†] A total of 30 states participated in the ABLES Program in 2013.

[§] The numbers and rates of adults with BLLs $\geq 25 \mu\text{g}/\text{dL}$ are subsets of the numbers and rates of adults with BLLs $\geq 10 \mu\text{g}/\text{dL}$.

[¶] All cases reported by a state. These include cases among adult residents in the reporting state plus cases identified by the reporting state but who reside in another state.

^{**} Adults residing in the reporting state.

^{††} 10–15 $\mu\text{g}/\text{dL}$ BLL data were not available.

^{§§} Data from Arizona were available only for January to August 2013.

TABLE 2. Reported numbers of adults* with blood lead levels $\geq 25 \mu\text{g}/\text{dL}$, by exposure source and area — state Adult Blood Lead Epidemiology and Surveillance programs, United States, 2013[†]

Division/State	Occupational [§]		Nonoccupational		Unknown		Total
	No.	(%)	No.	(%)	No.	(%)	No.
Total	4,262	(77.6)	285	(5.2)	944	(17.2)	5,491
New England							
Connecticut	37	(59.7)	23	(37.1)	2	(3.2)	62
Massachusetts	71	(56.3)	24	(19.0)	31	(24.6)	126
Vermont	3	(25.0)	4	(33.3)	5	(41.7)	12
Mid Atlantic							
New Jersey	105	(66.5)	— [¶]	(—)	53	(33.5)	158
New York	191	(64.7)	78	(26.4)	26	(8.8)	295
Pennsylvania	1,449	(94.5)	—	(—)	84	(5.5)	1,533
East North Central							
Illinois	177	(62.5)	14	(4.9)	92	(32.5)	283
Indiana	67	(59.3)	—	(—)	46	(40.7)	113
Michigan	70	(64.8)	28	(25.9)	10	(9.3)	108
Wisconsin	88	(83.8)	9	(8.6)	8	(7.6)	105
West North Central							
Iowa	200	(99.0)	2	(1.0)	—	(—)	202
Minnesota	92	(86.0)	3	(2.8)	12	(11.2)	107
Missouri	682	(98.8)	8	(1.2)	—	(—)	690
Nebraska	25	(78.1)	2	(6.3)	5	(15.6)	32
South Atlantic							
Florida	82	(30.4)	11	(4.1)	177	(65.6)	270
Georgia	100	(42.2)	—	(—)	137	(57.8)	237
Maryland	57	(76.0)	4	(5.3)	14	(18.7)	75
North Carolina	89	(89.9)	8	(8.1)	2	(2.0)	99
East South Central							
Alabama	353	(81.5)	—	(—)	80	(18.5)	433
Kentucky	—	(—)	—	(—)	94	(100.0)	94
West South Central							
Louisiana	78	(92.9)	5	(6.0)	1	(1.2)	84
Oklahoma	7	(24.1)	2	(6.9)	20	(69.0)	29
Mountain							
Arizona	12	(80.0)	3	(20.0)	—	(—)	15
Colorado	4	(13.8)	3	(10.3)	22	(75.9)	29
New Mexico	4	(30.8)	3	(23.1)	6	(46.2)	13
Wyoming	12	(100.0)	—	(—)	—	(—)	12
Pacific							
Alaska	5	(62.5)	—	(—)	3	(37.5)	8
California	146	(76.0)	45	(23.4)	1	(0.5)	192
Oregon	7	(58.3)	1	(8.3)	4	(33.3)	12
Washington	49	(77.8)	5	(7.9)	9	(14.3)	63

* An employed person aged ≥ 16 years at the time of blood collection. When an adult had multiple blood lead tests in a given year, only the highest blood lead level for that adult in that year was counted.

[†] Among the 30 reporting states, 29 states submitted data on exposure source in 2013. These data include adult residents in the state and residents of other states reported by the state ABLES programs.

[§] Includes 23 cases coded with both occupational and nonoccupational exposure source.

[¶] No cases were reported.

References

1. CDC. Summary of notifiable noninfectious conditions and disease outbreaks—United States. *MMWR Morb Mortal Wkly Rep* 2014;63(55).
2. CDC. Summary of notifiable infectious diseases and conditions—United States, 2014. *MMWR Morb Mortal Wkly Rep* 2014;63(54).
3. CDC. Adult Blood Lead Epidemiology and Surveillance (ABLES) Program. Cincinnati, OH: US Department of Health and Human Services, CDC, National Institute for Occupational Safety and Health; 2015. <http://www.cdc.gov/niosh/topics/ables/description.html>
4. Association of Occupational and Environmental Clinics. Medical management guidelines for lead-exposed adults. Washington, DC: Association of Occupational and Environmental Clinics; 2013. http://www.aococ.org/documents/positions/mmg_revision_with_cste_2013.pdf
5. Kosnett MJ, Wedeen RP, Rothenberg SJ, et al. Recommendations for medical management of adult lead exposure. *Environ Health Perspect* 2007;115:463–71. <http://dx.doi.org/10.1289/ehp.9784>
6. CDC. National Notifiable Diseases Surveillance System. Lead, elevated blood levels. 2016 Case definition; 2015. Atlanta, GA: US Department of Health and Human Services, CDC; 2015. <https://wwwn.cdc.gov/nndss/conditions/lead-elevated-blood-levels/case-definition/2016/>
7. Council of State and Territorial Epidemiologists. Position Statement 09-OH-02. Public health reporting and national notification for elevated blood: lead levels. Atlanta, GA: Council of State and Territorial Epidemiologists; 2009. <http://c.ymcdn.com/sites/www.cste.org/resource/resmgr/PS/09-OH-02.pdf>
8. US Department of Health and Human Services. Healthy people 2020: occupational safety and health objectives. Washington, DC: US Department of Health and Human Services; 2013. <https://www.healthypeople.gov/2020/topics-objectives/topic/occupational-safety-and-health/objectives>
9. CDC. Adult blood lead epidemiology and surveillance—United States, 2005–2007. *MMWR Morb Mortal Wkly Rep* 2009;58:365–9.
10. Occupational Safety and Health Administration. Directive number: CPL 03-00-009. OSHA instruction: National Emphasis Program on Lead. Washington, DC: US Department of Labor, Occupational Safety and Health Administration; 2008. https://www.osha.gov/OshDoc/Directive_pdf/CPL_03-00-0009.pdf
11. Barbosa F Jr, Tanus-Santos JE, Gerlach RF, Parsons PJ. A critical review of biomarkers used for monitoring human exposure to lead: advantages, limitations, and future needs. *Environ Health Perspect* 2005;113:1669–74. <http://dx.doi.org/10.1289/ehp.7917>
12. CDC. Fourth national report on human exposure to environmental chemicals. Updated tables, February 2015. Atlanta, GA: US Department of Health and Human Services, CDC; 2015. http://www.cdc.gov/biomonitoring/pdf/FourthReport_UpdatedTables_Feb2015.pdf
13. National Toxicology Program. Health effects of low-level lead evaluation. Research Triangle Park, NC: US Department of Health and Human Services, National Toxicology Program; 2013. <http://ntp.niehs.nih.gov/pubhealth/hat/noms/lead/index.html>

State Adult Blood Lead Epidemiology and Surveillance (ABLES) Program Investigators

Sherri Davidson, MPH, Alabama Department of Public Health; Brigitte Dufour, Matthew Roach, MPH, Kaleb Tsang, MS, Arizona Department of Health Services; Susan F. Payne, MA, California Department of Public Health; Amanda M. DeLoreto, MPH, Thomas St. Louis, MSPH, Connecticut Department of Public Health; Sudha Rajagopalan, MPH, Sharon Watkins, PhD, Juanita Chalmers, MPH, Florida Department of Health; Tiefu Shen, Illinois Department of Public Health; Jeffery M. Turner, Indiana State Department of Health; Kathy Leinenkugel, MPA, Iowa Department of Public Health; MaAdwoa Asamoah, MPH, Kentucky Department for Public Health; Jocelyn Lewis, PhD, Louisiana Department of Health and Hospitals; Ezattollah Keyvan, MD, Maryland Department of the Environment; Kenneth Roseman MD, Joanna Kica, MPA, Michigan State University; Stephanie Yendell, DVM, Minnesota Department of Health; Carol R. Braun, Missouri Department of Health and Senior Services; Derry Stover, MPH, Nebraska Department of Health and Human Services; Marija Borjan, PhD, Margaret E. Lumia, PhD, Devendra Singh, New Jersey Department of Health; Edward O. Irobi, PhD, Heidi Krapfl, New Mexico Department of Health; Alicia M. Fletcher, MPH, New York State Department of Health; Sheila Higgins, MPH, North Carolina Division of Public Health; Susan J. Quigley, Christin T. Benner, MPH, Oklahoma State Health Department; David Dreher, Oregon Health Authority; Tanecia Richardson, Pennsylvania Department of Health; Mike Sullivan, MBA, Vermont Department of Health; Carrie Tomasallo, PhD, Wisconsin Department of Health Services; Steve Melia, MSPH, Wyoming Department of Health; and ABLES Programs coordinators in Alaska Department of Health and Social Services; Colorado Department of Public Health and Environment; Georgia Department of Public Health; Massachusetts Department of Labor Standards; Washington Department of Labor and Industries.