

U.S. Centers for Disease Control and Prevention

MMWR

Morbidity and Mortality Weekly Report

Surveillance Summaries / Vol. 75 / No. 4

July 2, 2026

Surveillance for *Candida auris* — United States, 2022–2024



**U.S. DEPARTMENT OF
HEALTH AND HUMAN SERVICES**
CENTERS FOR DISEASE
CONTROL AND PREVENTION

CONTENTS

Introduction	2
Methods.....	2
Results	3
Discussion	4
Limitations	5
Future Directions	5
Conclusion	6
References.....	7

The *MMWR* series of publications is published by the Office of Science, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30329-4027.

Suggested citation: [Author names; first three, then et al., if more than six.] [Title]. *MMWR Surveill Summ* 2026;75(SS-#):[inclusive page numbers].

Centers for Disease Control and Prevention

Jay Bhattacharya, MD, PhD, *Acting Director*
Althea Grant-Lenzy, PhD, *Director, Office of Science*

MMWR Editorial and Production Staff (Serials)

Leonard Jack, Jr, PhD, MSc, *Acting Editor in Chief*

Michael Lynch, MD, MPH, *Acting Editor*
Mary Dott, MD, MPH, *Online Editor*
Terisa F. Rutledge, *Managing Editor*
Catherine B. Lansdowne, MS,
Acting Lead Technical Writer-Editor
Jackie Kelly, MS, *Project Editor*

Terraye M. Starr,
Acting Lead Health Communication Specialist
Alexander J. Gottardy,
Maureen A. Leahy,
Visual Information Specialists
Quang M. Doan, MBA,
Phyllis H. King, Moua Yang,
Information Technology Specialists

Lowery Johnson,
Health Communication Specialist
Will Yang, MA,
Visual Information Specialist

MMWR Editorial Board

Matthew L. Boulton, MD, MPH
Carolyn Brooks, ScD, MA
Virginia A. Caine, MD
Jonathan E. Fielding, MD, MPH, MBA

Timothy F. Jones, MD, *Chairman*
David W. Fleming, MD
William E. Halperin, MD, DrPH, MPH
Jewel Mullen, MD, MPH, MPA
Jeff Niederdeppe, PhD
Patricia Quinlisk, MD, MPH

Patrick L. Remington, MD, MPH
Carlos Roig, MS, MA
William Schaffner, MD
Morgan Bobb Swanson, MD, PhD

Surveillance for *Candida auris* — United States, 2022–2024

Jeremy A. W. Gold, MD¹; Anna D. Baker, MPH¹; Kaitlin Benedict, MPH¹; Kaitlin Forsberg, MPH¹; Jessica E. Laury, MPH¹; Ria R. Ghai, PhD¹; Danielle A. Rankin, PhD²; Maroia Spalding Walters, PhD²; Shawn R. Lockhart, PhD¹; Sophie Jones, PhD¹; Meghan Lyman, MD¹; *Candida auris* surveillance team members

¹Division of Foodborne, Waterborne, and Environmental Diseases, National Center for Emerging and Zoonotic Infectious Diseases, CDC; ²Division of Healthcare Quality Promotion, National Center for Emerging and Zoonotic Infectious Diseases, CDC

Abstract

Problem/Condition: *Candida auris* is an emerging yeast that is frequently resistant to antifungal drugs. *C. auris* can cause invasive infections associated with high mortality and can colonize patients asymptotically, which facilitates transmission in health care settings. Since it was first reported in the United States in 2016, *C. auris* has been identified in multiple states, with increasing numbers of cases reported annually. Monitoring national trends in cases identified through clinical testing and screening for colonization is critical to guide infection prevention and control efforts.

Period Covered: 2022–2024.

Description of System: State and jurisdictional health departments voluntarily report clinical and screening *C. auris* cases to CDC using standardized case definitions of the Council of State and Territorial Epidemiologists. Clinical cases are defined as detection of *C. auris* from specimens collected for diagnostic purposes; screening cases are defined as detection from colonization screening swabs. Cases were reported to CDC through the Research Electronic Data Capture (REDCap) or Data Collation and Integration for Public Health Event Response (DCIPHER) platforms. Data included patient age and sex, case type, specimen type (for clinical cases), health care facility type, Antimicrobial Resistance Laboratory Network geographic region, and specimen collection date. Analyses were descriptive and limited to cases with specimens collected during 2022–2024.

Results: During 2022–2024, a total of 13,507 clinical *C. auris* cases were reported to CDC, increasing from 2,882 in 2022 to 4,428 in 2023 and 6,197 in 2024, with smaller annual percentage increases over time (53.7% from 2022 to 2023 and 39.9% from 2023 to 2024). Most clinical cases occurred among adults aged ≥ 45 years (87.8%) and among males (61.0%). The most common specimen types among all clinical cases were urine (31.5%) and blood (30.2%); by year, the proportion of blood as the specimen type was 34.4% in 2022, 30.2% in 2023, and 25.6% in 2024. Most clinical cases were identified through specimens collected in acute care hospitals (76.6%) and long-term acute care hospitals (17.8%).

During the same period, a total of 27,853 screening cases were reported to CDC, increasing from 6,226 in 2022 to 9,195 in 2023 and 12,432 in 2024. Screening cases most frequently occurred among adults aged ≥ 45 years (90.0%) and males (57.9%). Among cases with known facility type, the proportion of specimens collected in acute care hospitals increased from 24.7% in 2022 to 50.7% in 2024, whereas the proportion of specimens collected in long-term acute care hospitals decreased from 56.1% to 35.7% during the same period.

Interpretation: The number of clinical and screening *C. auris* cases reported to CDC increased during 2022–2024, indicating ongoing transmission in U.S. health care settings. Although annual percentage increases in clinical cases declined over time, absolute case counts reported to CDC continued to rise. The increasing proportion of screening cases with specimens collected in acute care hospitals might reflect increased use of screening in acute care hospitals, including screening at admission.

Public Health Action: Because of increases in the number of reported *C. auris* cases, sustained infection prevention and control efforts in health care facilities, including adherence to transmission-based precautions, environmental disinfection with agents effective against *C. auris*, and communication of *C. auris* status during patient transfers remain essential to preventing clinical infections and colonization. Because this pathogen is frequently resistant to antifungal drugs, continued investment in laboratory capacity and surveillance, including antifungal susceptibility testing and screening of patients at high risk for *C. auris* infection,

can support timely detection and guide prevention strategies. Ongoing public health coordination at federal, state, and local levels is critical to limit further spread and to address emerging antifungal drug resistance.

Corresponding author: Jeremy A. W. Gold, Division of Foodborne, Waterborne, and Environmental Diseases, National Center for Emerging and Zoonotic Infectious Diseases, CDC. Telephone: 404-718-3650; Email: jgold@cdc.gov.

Introduction

Candida auris is an emerging, frequently drug-resistant yeast that can cause invasive infections with high mortality rates (1–4). It can colonize patients asymptotically and persist on surfaces, facilitating rapid spread in health care settings, especially in acute care hospitals (ACHs), long-term acute care hospitals (LTACHs), and skilled nursing facilities (SNFs) equipped with a ventilator (5–7). Symptoms of *C. auris* infection are nonspecific and depend on the affected body site and infection severity, ranging from superficial skin infections to invasive forms such as bloodstream infections (i.e., candidemia) (7). Because most *C. auris* isolates are resistant to the azole antifungal drug fluconazole, a common treatment for candidal infections, echinocandin antifungals are first-line treatments (8). However, echinocandin-resistant and multidrug-resistant strains (resistant to azoles, echinocandins, and polyenes) have been detected in the United States; although echinocandin resistance remains uncommon, these strains can limit available treatment options (9–11).

Risk factors for *C. auris* colonization (i.e., presence on skin or mucosal surfaces without causing infection) and for invasive infection include frequent or prolonged stays in health care facilities, complex medical care involving invasive medical devices (e.g., mechanical ventilation and central venous catheters), and recent antimicrobial use (12). Screening for *C. auris* colonization in patients at high risk is a critical component of strategies to prevent spread in health care facilities because identification of colonized patients enables facilities to implement recommended infection prevention and control measures (13). Deciding which patients to screen might be based on multiple factors, including local *C. auris* epidemiology, epidemiologic linkages to other cases (e.g., those on the same unit), specific patient risk factors, and the purpose of screening (14). Screening can be performed using a broad approach by conducting a point prevalence survey (e.g., screening all patients or residents of a unit or facility) or a more targeted approach (e.g., screening certain patients on the basis of risk factors, epidemiologic links, or clinical characteristics) (14).

The first *C. auris* cases were reported in the United States in 2016 (15). Early cases were usually associated with health care exposure abroad, whereas most of the recent cases were acquired in U.S. health care settings (1,16). The number of reported clinical *C. auris* cases increased from approximately 50 in 2016 to approximately 1,500 in 2021; the largest percentage increase (95%) occurred from 2020 to 2021 (1,16). In certain states and jurisdictions, *C. auris* has become endemic, leading to sustained high rates of transmission and increasing prevalence across health care facilities. Meanwhile, other states and jurisdictions are just beginning to report their first cases (16).

A comprehensive understanding of the epidemiology of *C. auris* in the United States is critical to guide public health response efforts. This report summarizes national *C. auris* case report data for clinical and screening cases identified during 2022–2024. Public health officials and policymakers can use these findings to strengthen surveillance and public health responses, guide resource allocation, and focus initiatives aimed at enhancing patient safety in health care facilities.

Methods

Surveillance Case Definitions

Consistent with the [Council of State and Territorial Epidemiologists](#) (CSTE) 2023 *C. auris* case definitions (17), a clinical case was defined as the detection of *C. auris* from a patient specimen tested to determine the cause of a suspected infection. The specimen could be taken from blood, which typically indicates an invasive infection, or from other diagnostic specimen types (e.g., urine) that might represent an infection or colonization detected during clinical testing. A screening case was defined as the detection of *C. auris* from a swab, most often a composite 1) axilla and groin or 2) nares, axilla, and groin swab collected to test for colonization. A patient could be classified as having up to two cases: one screening case followed by one clinical case; analyses were conducted at the case level, and multiple positive specimens within a single case type were not counted as separate cases.

Data Sources

Reporting requirements for *C. auris* vary by state and jurisdiction. State and jurisdictional health departments can voluntarily report clinical and screening *C. auris* cases to CDC. *C. auris* cases from states and jurisdictions were transmitted to CDC using either the secure Research Electronic Data Capture (REDCap) database (18,19) or Data Collation and Integration for Public Health Event Response (DCIPHER), a cloud-based CDC data platform. The data collected included information on patient age, sex, case type (i.e., clinical or screening), specimen type for clinical cases (i.e., blood, wound, urine, respiratory, or other), date of specimen collection, facility type, and Antimicrobial Resistance Laboratory Network (AR Laboratory Network) region of the facility where the specimen was collected. The AR Laboratory Network regions are *Central* (Arkansas, Iowa, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Oklahoma, and South Dakota); *Mid-Atlantic* (Delaware, District of Columbia, Maryland, North Carolina, Pennsylvania, South Carolina, Virginia, and West Virginia); *Midwest* (Illinois, Indiana, Kentucky, Michigan, Ohio, and Wisconsin); *Mountain* (Arizona, Colorado, Idaho, Montana, New Mexico, Texas, Utah,

and Wyoming); *Northeast* (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, and Vermont); *Southeast* (Alabama, Florida, Georgia, Louisiana, Mississippi, Puerto Rico, and Tennessee); and *West* (Alaska, California, Guam, Hawaii, Nevada, Oregon, and Washington).

In the United States, *C. auris* clinical cases have been nationally notifiable since 2019, and *C. auris* screening cases have been nationally notifiable since 2023. Nationally notifiable conditions are typically reported to CDC through the [National Notifiable Diseases Surveillance System](#) (NNDSS). However, data analyzed for this report were case-level data from CDC's case-based surveillance system, which includes variables not available in NNDSS (e.g., facility type). Data included in these analyses were current as of February 20, 2026 (the date on which the analytic dataset for this report was finalized). CDC web-based surveillance summaries might display different case totals because of ongoing case reporting, data updates, and reconciliation.

Analyses

Total *C. auris* case counts for specimens collected during 2022–2024 and annual counts by case type, age group, sex, AR Laboratory Network region, specimen type for clinical cases (i.e., blood, wound, urine, respiratory, or other), and facility type (i.e., LTACH, ACH, SNF, ventilator-equipped SNF, or other) at the time of specimen collection were examined. For each patient, only the first specimen (by date collected) per case type (i.e., screening or clinical) was included in analyses. *Candida* isolates from blood are usually identified to the species level, whereas species identification for nonblood isolates is dependent on local laboratory practices (e.g., implementation of enhanced surveillance to identify all yeast from urine); therefore, year-specific analyses of specimen type focused on blood specimens.

Analyses for this report were descriptive; statistical testing was not performed because denominator data were unavailable and testing and screening practices varied over time and across jurisdictions. The primary measure for this analysis was the number of reported *C. auris* cases by year of specimen collection. These counts are influenced by multiple factors, including changes in testing capacity, screening practices, and reporting across jurisdictions. Therefore, findings should be interpreted as descriptions of trends in reported cases within the surveillance system and not representative of population-based incidence. This activity was reviewed by CDC, deemed not research, and conducted consistent with applicable federal law and CDC policy.*

*45 C.F.R. part 46.102(l)(2), 21 C.F.R. part 56; 42 U.S.C. Sect. 241(d); 5 U.S.C. Sect. 552a; 44 U.S.C. Sect. 3501 et seq.

Results

Clinical *C. auris* Cases

A total of 13,507 clinical *C. auris* cases from specimens collected during 2022–2024 were reported to CDC (Table 1). The number of clinical *C. auris* cases reported to CDC increased from 2,882 in 2022 to 4,428 in 2023 to 6,197 in 2024. Overall, most clinical cases occurred among adults aged 45–64 years (32.2%) or 65–74 years (29.5%); 19.6% of cases occurred among those aged 75–84 years, 12.0% among those aged 18–44 years, 6.5% among adults aged ≥85 years, and 0.2% among those aged 0–17 years. Most cases were among males (61.0%). The most common specimen type was urine (31.5%), followed by blood (30.2%), wound (16.7%), and respiratory (13.1%). The majority of specimens were collected from the West AR Laboratory Network region (28.5%), the Midwest region (21.3%), or the Southeast region (20.2%). The most common facility type for specimen collection was ACH (76.6%), followed by LTACH (17.8%) and SNF (1.7%). Overall, data completeness varied across variables; the highest proportion of missing data was for facility type (missing for 13.1% of clinical cases and 12.5% of screening cases).

Of the specimen types among clinical cases, blood accounted for 34.4% (991 of 2,882) in 2022, 30.2% (1,337 of 4,428) in 2023, and 25.6% (1,586 of 6,197) in 2024 (Tables 1 and 2). Among clinical cases with blood specimens, age, sex, and facility type distributions were similar across years. Regional percentage distributions were also comparable across years, but changes in percentages were observed from 2022 to 2024, including in the Northeast (21.8% of cases in 2022 to 13.7% in 2024).

Screening *C. auris* Cases

A total of 27,853 screening cases with specimens collected during 2022–2024 were reported to CDC (Table 3). The number of screening cases reported to CDC increased from 6,226 in 2022 to 9,195 in 2023 to 12,432 in 2024. Overall, most screening cases occurred among adults aged 45–64 years (30.6%) or 65–74 years (29.5%); 21.3% of cases occurred among adults aged 75–84 years, 10.0% among those aged 18–44 years, 8.5% among those aged ≥85 years, and <0.1% among those aged 0–17 years. Most (57.9%) screening cases were among males, and the largest proportion of screening specimens were collected from the West AR Laboratory Network region (36.0%), followed by the Southeast (18.8%) and Midwest regions (18.8%). The most common facility type for specimen collection was LTACH (42.1%), followed by ACH (41.8%) and ventilator-equipped SNF (10.3%).

From 2022 to 2024, the distribution patterns for age, sex, and region were comparable. Among screening cases with known facility type (n = 24,373 [87.5%]; facility type was unknown for 12.5% of cases), the percentage of cases from ACHs changed from 24.7% (n = 1,345 of 5,438) in 2022 to 41.0% (n = 3,235 of 7,894) in 2023 to 50.7% (n = 5,598 of 11,041) in 2024, whereas the percentage from LTACHs changed from 56.1% (n = 3,048 of 5,438) in 2022 to 41.5% (n = 3,278 of 7,894) in 2023 to 35.7% (n = 3,937 of 11,041) in 2024.

Discussion

This analysis of U.S. national surveillance data revealed that 2,882 clinical cases of *C. auris* were reported to CDC in 2022, representing an approximate 95.9% increase compared with 2021 (1). Case counts continued to increase in 2023, with 4,428 cases reported, and in 2024, with 6,197 cases, but the annual changes were smaller over time: 95.9% (2021 to 2022), 53.7% (2022 to 2023), and 39.9% (2023 to 2024). The pronounced increase in clinical case counts from 2021 to 2022 might have been influenced by multiple factors, including strains on health care systems caused by the COVID-19 pandemic, which resulted in shortages of supplies and personnel, as well as overcrowding (20). In addition, more patients might have been at risk for *C. auris* colonization and infection during this period because of the widespread use of broad-spectrum antibiotics to treat secondary bacterial infections associated with COVID-19 and the increased population of critically ill patients receiving mechanical ventilation and having prolonged stays at health care facilities (21). The smaller percentage increases observed after 2022 might reflect, in part, a renewed focus on standard protocols aimed at mitigating the spread of *C. auris* and other health care–associated infections, particularly after the resolution of personal protective equipment shortages and staffing challenges experienced during the earlier phases of the COVID-19 pandemic. Improved testing capacity might also have contributed to greater detection of clinical *C. auris* cases over time as laboratories have increasingly adopted species-level identification or updated laboratory equipment (e.g., testing libraries) to properly identify *C. auris*. Increased screening, particularly in ACHs, and changes in testing and reporting practices over time might also have contributed (22,23). However, the relative contribution of these factors to the observed increases cannot be quantified with the available data.

The continued increase in clinical *C. auris* case counts reported to CDC during 2022–2024 underscores the establishment and ongoing transmission of this pathogen. Clinical cases most frequently involved specimens collected

from patients in ACHs (76.6%) and LTACHs (17.8%), although the data do not indicate where transmission might have occurred. This finding highlights the importance of strict adherence to infection prevention and control measures in health care facilities, which is bolstered by sustained support from state and local health departments, as well as federal agencies, through guidance, technical assistance, and site visits (24). CDC recommendations involve several core components of infection prevention and control in health care settings, including rigorous hand hygiene practices, the use of [disinfectants effective against *C. auris*](#) in facilities caring for patients with *C. auris*, and the implementation of transmission-based precautions, such as [contact precautions](#) in ACHs and LTACHs and either contact precautions or enhanced barrier precautions in nursing homes and skilled nursing facilities, depending on the setting and jurisdictional recommendations (24). CDC also recommends regular audits of infection control practices and measures to ensure effective communication among health care facilities of patient *C. auris* status during transfers (24). In general, facilities that are equipped to care for patients with other multidrug-resistant organisms or *Clostridioides difficile* can also effectively care for patients with *C. auris* (24).

Approximately one third of reported clinical cases (30.2%) were identified from blood specimens. Because isolating *Candida* from blood indicates an invasive infection, this result is concerning because of the pathogen's frequent antifungal drug resistance and high mortality (30%–72%) (25–27). Trends in nonblood specimen types should be interpreted with caution because changes in laboratory practices, such as expanded species-level identification of yeast from nonsterile sources (e.g., urine), might influence their relative distribution over time. These findings underscore the importance of efforts to prevent health care–associated *Candida* bloodstream infections. Consistent with previous data, clinical cases predominantly occurred in adults aged ≥45 years, with most cases identified in the West (28.5%), Midwest (21.3%), and Southeast (20.2%) regions of the AR Laboratory Network (1). Clinical *C. auris* cases were more common among males (61.0%); the reasons for this pattern are unclear and were not evaluated in this surveillance analysis.

The number of screening *C. auris* cases reported to CDC increased from 6,226 in 2022 to 12,432 in 2024, and by 2024, screening cases were most frequently detected in ACHs. Initially, *C. auris* screening programs and containment efforts focused on LTACHs and ventilator-equipped SNFs, because these were identified as settings at high risk for *C. auris* colonization and outbreaks (28,29). This risk is largely because of their patient populations, which often include patients receiving mechanical ventilation, experiencing prolonged hospital stays,

and with severe underlying conditions (13,28). The reason for the shift in screening case detection is uncertain; however, it might indicate increased use of screening in ACHs to identify colonized patients during admission, rather than reflecting actual transmission within ACHs (23). Consequently, ACHs might be identifying colonized patients at the time of admission through expanded in-house or admission screening protocols or policies, even when colonization acquisition occurred at another facility (1). Finally, screening cases became nationally notifiable in 2023, which might have improved completeness of reporting and contributed to increases observed after 2022.

Overall, because the number of *C. auris* screening cases is dependent on local screening practices, which vary among facilities and regions (30), determining the degree to which increased screening case counts reflect a true increase in disease prevalence rather than sampling bias is difficult. The increase in screening cases might be caused by heightened awareness and enhanced surveillance efforts as more institutions implement routine screening protocols. This increase might be linked to the adoption by public health agencies and health care facilities of CDC guidelines, which promote targeted and, when warranted, facilitywide screening for *C. auris*, with increased frequency among populations at high risk for infection on the basis of local epidemiology and transmission risk (14). In addition, advancements in diagnostic testing, particularly the increased use of polymerase chain reaction–based tests that are more rapid and sensitive than traditional culture-based methods, might also have contributed to increased screening case detection (31–33).

Limitations

The findings in this report are subject to several limitations. First, *C. auris* case counts in this report might differ from those reported publicly by individual state and jurisdictional health departments or other surveillance platforms, which might limit direct comparisons. Differences across data sources might reflect variations in surveillance systems (e.g., REDCap and NNDSS), surveillance case definitions, inclusion criteria (e.g., individual surveillance cases, which might represent more than one case per patient; laboratory specimens; or unique patients), assignment on the basis of patient residence versus location of specimen collection, and use of report date versus specimen collection date. Second, although data in CDC’s case-based surveillance system are reported using standardized definitions (i.e., CSTE surveillance case definitions), which supports consistency across jurisdictions, these standardized data are not intended to replicate jurisdiction-specific public reporting. Continued efforts to align reporting platforms and

required data elements across jurisdictions could enhance data completeness and standardization. Third, because clinical cases are defined on the basis of detection of *C. auris* from specimens collected for diagnostic purposes, rather than on the basis of clinical evidence of infection, certain specimen types, particularly urine and respiratory specimens, might represent colonization rather than true infection in the absence of detailed clinical information. Fourth, certain denominator data, including the number of patients tested or screened, the number of participating facilities, and facility- or jurisdiction-level census data, were unavailable; therefore, incidence and prevalence could not be estimated, and changes over time might reflect changes in ascertainment rather than true changes in disease frequency. Fifth, the prevalence data in this report are likely underestimates of the true prevalence of *C. auris* because colonized patients might not be screened, and certain clinical infections might not be detected because cultures are not obtained or isolates are not identified at the species level. Sixth, *C. auris* screening cases became nationally notifiable in 2023, which might have increased completeness of reporting in 2023 compared with 2022 and affected interpretation of temporal trends. Seventh, jurisdiction-level changes in reporting requirements, as well as changes in testing and screening practices over time, might have influenced case counts; therefore, described trends should be interpreted with caution and might not fully reflect true changes in incidence or prevalence. Finally, the surveillance system does not include information on negative test results, race, ethnicity, socioeconomic information, treatments, outcomes, or antifungal susceptibility testing, which would be valuable for guiding public health surveillance and treatment guidelines (1,4).

Future Directions

Sustained investment in laboratory testing capacity for *C. auris* identification and antifungal drug susceptibility testing, such as through the AR Laboratory Network and by other laboratories, could facilitate rapid case detection and evaluation of emerging trends such as changes in antimicrobial resistance patterns. Expanding testing capacity in clinical and commercial laboratories might further improve timely detection by bringing testing closer to health care facilities. Together, these investments might support enhanced surveillance efforts, which might help refine prevention strategies by identifying the most effective screening approaches for preventing transmission and understanding modifiable factors associated with the progression from colonization to infection (34). Enhanced surveillance might also direct clinical guidance by providing information on treatment practices, patient outcomes, and antifungal drug susceptibility testing patterns and results

(10,25). Research initiatives might focus on identifying effective decolonization or pathogen reduction strategies to reduce the risk for infection among colonized patients and to disrupt *C. auris* transmission. In addition, initiatives could focus on optimizing recommended antimicrobial use and developing novel antifungal treatment options to expand the range of therapies available against resistant strains (35–37).

Conclusion

C. auris remains a critical public health concern in the United States and continues to be most frequently identified among middle-aged and older adults, particularly in specimens collected from patients in ACHs and LTACHs. From 2022 to 2024, the number of clinical *C. auris* cases reported to CDC increased approximately twofold, from 2,882 to 6,197. Although the annual percent increase in reported cases decreased during the study period, the continued rise in cases underscores ongoing transmission in health care settings and the importance of infection prevention and control efforts, with continued support from federal, state, and local public health partners to prevent further spread.

Candida auris Surveillance Team Members

Katherine Draper, Melanie Roderick, Alabama Department of Public Health; Sarah Fitzhugh, Kelley Garner, Arkansas Department of Health; Diana Holden, Sam Horwich-Scholefield, California Department of Public Health; Hira Adil, Michelle Funk, Shane Zelencik, Chicago Department of Public Health; Christopher A. Czaja, Helen Johnston, Kelsey Orten, Colorado Department of Public Health and Environment; Meghan Maloney, Connecticut Department of Public Health; Marie Argyriou, Mefruz Haque, Charity Gaitirira, Candace Johnson, District of Columbia Department of Health; Argentina Charles, Jalysa Erskine, Mary Syla, Florida Department of Health; Michael Anderson, Teresa Fox, Ben Roberts, Georgia Department of Public Health; Deb Patterson Burdsall, Hektoen Institute of Medicine; Tracy Boyle, Dawn Chinn-Flournoy, Mai Vue, Illinois Department of Public Health; Marcie Bryant, Caleb Cox, Cayman Doran, Indiana Department of Health; Andrew Hennenfent, Michael Patten, Kenzie Teno, Iowa Department of Health and Human Services; Michael Curran, Andrea Flinchum, Honey (Mary) Issac, Cody Rocha, Kentucky Department of Public Health; Wendy Knight, Jennifer Nguyen, Kelsey OYong, Los Angeles County Department of Public Health; Natalie Christophe, Ashley Marin, Louisiana Department of Health; Serena Bailey, Siru Prasai, Karen Zabel, Maricopa County Department of Public Health Administration; Brittany Grace, Elisabeth Vaeth, Maryland Department of Health; Liore Klein, Maryland Department of Health Laboratories Administration; Joseph Gerth, Jessica Leaf, T. Valleroy, Massachusetts Department of Public Health; Brenda Brennan, Morgan Bulk, Nicole McGuire, Sara McNamara, Margaret Sturgis, Michigan Department of Health and Human Services; Krista Knowles, Laura Tourdot, Minnesota Department of Health; Jennifer Narcy, Brittany Fowler Whittington, Mississippi State Department of Health; Sarah Hanson, George Turabelidze, Brooklyn White, Missouri Department of Health and Senior Services; Erika Baldry, Victoria Doll, Montana Department of Public Health and Human Services; Muhammad Salman Ashraf, Lacey Pavlovsky, Nebraska Department of Health and Human Services; Divine Abbasszadeh, Shreya Batra, Richa Chaturvedi, Clarence Q.T. Collins, Christine Emmons, Pang Vang,

Nevada Office of State Epidemiology; Claire Burns-Lynch, Shivani Patel, Adrienne Sherman, New Jersey Department of Health; Melissa Judson, Marla M. Sievers, New Mexico Department of Health; Ernie Clement, Sarah Kogut, New York State Department of Health; Catherine Bryan, Lauren Pasutti, North Carolina Department of Health and Human Services; Faye Salzer, Jordan Taghon, North Dakota Department of Health and Human Services; Sarah Mitchell, Rachel Saadey, Ohio Department of Health; Jennifer Brown, Megan Crumpler, Mi Le, Orange County Health Care Agency; Evelyn Donahoe, Heather Hertzog, Alexia Y. Zhang, Oregon Health Authority; Alison Feathers, Jenna Sinkevitch, Pennsylvania Department of Health; Bonnie G. Chaoui, Michelle L. Gardner, Philadelphia Department of Public Health; Patricia McAuley, Nancy B. Persson, Rhode Island Department of Health; Ben Cohen, Erin Johnson, South Carolina Department of Public Health; Erin Christenson, Simone Godwin, Tennessee Department of Health; Thi Dang, Gabrielle Franco, Jenna Harlan, Jennifer E. Lee, Texas Department of State Health Services; Joshua Mongillo, Ashley Young, Utah Department of Health and Human Services; Paul Capcara, Jenna Cebelius, Allison G. Lafferty, Cedric Mubikayi, Vermont Department of Health; Shaina Bernard, Carolyn Kiefer, Allison O'Rourke, Virginia Department of Health; Emily Schneider, Mike Tran, Audrey Valentine, Washington State Department of Health; Sudha Chaturvedi, Anuradha Marathe, Mayuri Vaidya, Wadsworth Center, New York State Department of Health; Valerie Jividen, Emilee Melchior, West Virginia Department of Health; Greta Starr, Wisconsin Department of Health Services; Megan Lasure, Ann Valley, Wisconsin State Laboratory of Hygiene.

Acknowledgments

Cierra Jackson, Alabama Department of Public Health; Tisha Mitsunaga, California Department of Public Health; Justa Ewalu, Sonali Shakya, Florida Department of Health; Lynda Henderson, Renee Miller, Sue Milligan, Lynn Reynolds, Connie Stanfill, JoAnna Wagner, Karen Williams, Diane Yi, Georgia Department of Public Health; Tamara Bell, Kristen Crawford, Yvonne Overton, Hillary Spencer, Angela Tang, Nneka Ukegbu, Illinois Department of Public Health; Chris Barten, Tyra Goss, Abby Hough, DeeAnn Vaage, Lisa Vitale, Iowa Department of Health and Human Services; Carrie Welch, Kansas Department of Health and Environment; Julia Bowman, Teresa Curry, Monica Hernandez, Olivia Hunziker, Niki Marquardt, Bianca Munoz, Maricopa County Department of Public Health; Marcus Bernal, Jinshui Fan, Michael Gotesman, John Martinez, Tyler Maruca, Michele Plehn, Alessandra Rivera, Greg Steffensen, John Thoguru, Maryland Department of Health Laboratories Administration; Niki Mach, Michigan Department of Health and Human Services; Tammy Hale, Christine Lees, Ruth Lynfield, Tara Suhs, Jacy Walters, Minnesota Department of Health; Onyekachi Ahaiwe, Missouri Department of Health and Senior Services; Rebecca Greeley, New Jersey Department of Health; Trina Ryans, Kim Southerland, Oklahoma State Department of Health; Jelynn Alano, Victoria Buchanan, Tania Chiem, Stacy Dykman, Estephany Esqueda, Cherry Q. Fontela, Elizabeth Kryger, Denise Tirol, Angelica Torres, Gabrielle Villareal, Matthew Zahn, Orange County Health Care Agency; Adelina Mart, Dat Tran, Oregon Health Authority; Bobbie Rae Erickson, Monica Giacomucci, Julie Paoline, Pennsylvania Department of Health; Jane M. Gould, Tiina Peritz, Philadelphia Department of Public Health; Rhode Island State Health Laboratories (Special Pathogens & Biothreats Laboratory), Rhode Island Department of Health; Cherlly Bailey, Carolyn Stover, Tennessee Department of

Health; Devin Beard, Beatrice Jensen, Linda Rider, Angela Weil, Utah Department of Health and Human Services; Audrey Brezak, Marisa D'Angeli, Sophie Hun, Kelly Kauber, Washington State Department of Health.

Conflicts of Interest

All authors have completed and submitted the International Committee of Medical Journal Editors form for disclosure of potential conflicts of interest. No potential conflicts of interest were disclosed.

References

- Lyman M, Forsberg K, Sexton DJ, et al. Worsening spread of *Candida auris* in the United States, 2019 to 2021. *Ann Intern Med* 2023;176:489–95. PMID:36940442 <https://doi.org/10.7326/M22-3469>
- Chakrabarti A, Sood P, Rudramurthy SM, et al. Incidence, characteristics and outcome of ICU-acquired candidemia in India. *Intensive Care Med* 2015;41:285–95. PMID:25510301 <https://doi.org/10.1007/s00134-014-3603-2>
- Lockhart SR, Etienne KA, Vallabhaneni S, et al. Simultaneous emergence of multidrug-resistant *Candida auris* on 3 continents confirmed by whole-genome sequencing and epidemiological analyses. *Clin Infect Dis* 2017;64:134–40. PMID:27988485 <https://doi.org/10.1093/cid/ciw691>
- Meek HC, Konkle S, Ostrowsky B, et al. Characterization of patients with *Candida (Candidozyma) auris* before and during the COVID-19 pandemic in New York, 2017–2022. *Am J Infect Control* 2025;53:1034–42. PMID:40578439 <https://doi.org/10.1016/j.ajic.2025.06.016>
- Arenas SP, Persad PJ, Patel S, et al. Persistent colonization of *Candida auris* among inpatients rescreened as part of a weekly surveillance program. *Infect Control Hosp Epidemiol* 2024;45:762–5. PMID:38087651 <https://doi.org/10.1017/ice.2023.251>
- Pacilli M, Kerins JL, Clegg WJ, et al. Regional emergence of *Candida auris* in Chicago and lessons learned from intensive follow-up at 1 ventilator-capable skilled nursing facility. *Clin Infect Dis* 2020;71:e718–25. PMID:32291441 <https://doi.org/10.1093/cid/ciaa435>
- Forsberg K, Woodworth K, Walters M, et al. *Candida auris*: the recent emergence of a multidrug-resistant fungal pathogen. *Med Mycol* 2019;57:1–12. PMID:30085270 <https://doi.org/10.1093/mmy/myy054>
- CDC. *Candida auris (C. auris)*: clinical treatment of *C. auris* infections. Atlanta, GA: US Department of Health and Human Services, CDC; 2024. <https://www.cdc.gov/candida-auris/hcp/clinical-care/index.html>
- Ostrowsky B, Greenko J, Adams E, et al.; *C. auris* Investigation Work Group. *Candida auris* isolates resistant to three classes of antifungal medications—New York, 2019. *MMWR Morb Mortal Wkly Rep* 2020;69:6–9. PMID:31917780 <https://doi.org/10.15585/mmwr.mm6901a2>
- Lyman M, Forsberg K, Reuben J, et al. Notes from the field: transmission of pan-resistant and echinocandin-resistant *Candida auris* in health care facilities—Texas and the District of Columbia, January–April 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:1022–3. PMID:34292928 <https://doi.org/10.15585/mmwr.mm7029a2>
- Laury JE, Forsberg K, Berkow EL, et al. *Candida auris* testing by the Antimicrobial Resistance Laboratory Network, United States, 2022–2023. *Emerg Infect Dis* 2026;32:308–10. PMID:41714865 <https://doi.org/10.3201/eid3202.251043>
- McKinnell JA, Singh RD, Miller LG, et al. The SHIELD Orange County project: multidrug-resistant organism prevalence in 21 nursing homes and long-term acute care facilities in Southern California. *Clin Infect Dis* 2019;69:1566–73. PMID:30753383 <https://doi.org/10.1093/cid/ciz119>
- Karmarkar EN, O'Donnell K, Prestel C, et al. Rapid assessment and containment of *Candida auris* transmission in postacute care settings—Orange County, California, 2019. *Ann Intern Med* 2021;174:1554–62. PMID:34487450 <https://doi.org/10.7326/M21-2013>
- CDC. *Candida auris (C. auris)*: screening recommendations for healthcare facilities. Atlanta, GA: US Department of Health and Human Services, CDC; 2025. <https://www.cdc.gov/candida-auris/hcp/screening-hcp/index.html>
- Vallabhaneni S, Kallen A, Tsay S, et al. Investigation of the first seven reported cases of *Candida auris*, a globally emerging invasive, multidrug-resistant fungus—United States, May 2013–August 2016. *MMWR Morb Mortal Wkly Rep* 2016;65:1234–7. PMID:27832049 <https://doi.org/10.15585/mmwr.mm6544e1>
- CDC. *Candida auris (C. auris)*: tracking *C. auris*. Atlanta, GA: US Department of Health and Human Services, CDC; 2026. <https://www.cdc.gov/candida-auris/tracking-c-auris/index.html>
- CDC. Case data implementation resources: *Candida auris* 2023 case definition. Atlanta, GA: US Department of Health and Human Services, CDC; 2023. <https://ndc.services.cdc.gov/case-definitions/candida-auris-2023/>
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81. PMID:18929686 <https://doi.org/10.1016/j.jbi.2008.08.010>
- Harris PA, Taylor R, Minor BL, et al.; REDCap Consortium. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208. PMID:31078660 <https://doi.org/10.1016/j.jbi.2019.103208>
- Ham DC, Li R, Mitsunaga T, et al. Clusters of emerging multidrug-resistant organisms in US health care facilities during the initial months of the SARS-CoV-2 pandemic. *Am J Infect Control* 2024;52:1390–6. PMID:39089494 <https://doi.org/10.1016/j.ajic.2024.07.013>
- Schaefer S, Walits E, Thaler K, Patel G. Impact of the COVID-19 pandemic on *Candida auris* infections: a retrospective analysis in an academic medical center in New York City. *Open Forum Infect Dis* 2024;11:ofae148. PMID:38887476 <https://doi.org/10.1093/ofid/ofae148>
- Keighley C, Garnham K, Harch SAJ, et al. *Candida auris*: diagnostic challenges and emerging opportunities for the clinical microbiology laboratory. *Curr Fungal Infect Rep* 2021;15:116–26. PMID:34178208 <https://doi.org/10.1007/s12281-021-00420-y>
- Gold JAW, Benedict K, Haas KA, et al. Yeast identification, antifungal susceptibility testing, and *Candida auris* screening practices at acute care hospitals and long-term acute care hospitals, United States. *J Clin Microbiol* 2026;64:e00091-26. <https://doi.org/10.1128/jcm.00091-26>
- CDC. *Candida auris (C. auris)*. Infection control guidance: *Candida auris*. Atlanta, GA: US Department of Health and Human Services, CDC; 2024. <https://www.cdc.gov/candida-auris/hcp/infection-control/index.html>
- Benedict K, Forsberg K, Gold JAW, Baggs J, Lyman M. *Candida auris*-associated hospitalizations, United States, 2017–2022. *Emerg Infect Dis* 2023;29:1485–7. PMID:37347923 <https://doi.org/10.3201/eid2907.230540>
- Osei Sekyere J. *Candida auris*: a systematic review and meta-analysis of current updates on an emerging multidrug-resistant pathogen. *MicrobiologyOpen* 2018;7:e00578. PMID:29345117 <https://doi.org/10.1002/mbo3.578>
- Zhu Y, O'Brien B, Leach L, et al. Laboratory analysis of an outbreak of *Candida auris* in New York from 2016 to 2018: impact and lessons learned. *J Clin Microbiol* 2020;58:e01503-19. PMID:31852764 <https://doi.org/10.1128/JCM.01503-19>
- Paul P, Forsberg K, Vallabhaneni S, et al. Transmissibility of *Candida auris* by type of inpatient healthcare facility. *Open Forum Infect Dis* 2018;5(suppl_1):S386–7. <https://doi.org/10.1093/ofid/ofy210.1101>
- Paul P, Slayton RB, Kallen AJ, Walters MS, Jernigan JA. Modeling regional transmission and containment of a healthcare-associated multidrug-resistant organism. *Clin Infect Dis* 2020;70:388–94. PMID:30919885 <https://doi.org/10.1093/cid/ciz248>

30. Hennessee I, Forsberg K, Beekmann SE, Polgreen P, Gold J, Lyman M. *Candida auris* screening practices at healthcare facilities in the United States: a survey of the Emerging Infections Network. *Antimicrob Steward Healthc Epidemiol* 2023;3(Suppl 2):s99. <https://doi.org/10.1017/ash.2023.371>
31. Leach L, Zhu Y, Chaturvedi S. Development and validation of a real-time PCR assay for rapid detection of *Candida auris* from surveillance samples. *J Clin Microbiol* 2018;56:e01223-17. PMID:29187562 <https://doi.org/10.1128/JCM.01223-17>
32. Rowlands J, Dufort E, Chaturvedi S, et al. *Candida auris* admission screening pilot in select units of New York City health care facilities, 2017–2019. *Am J Infect Control* 2023;51:866–70. PMID:36736380 <https://doi.org/10.1016/j.ajic.2023.01.012>
33. Nijhuis RHT, Rispens T, Godschalk PCR. Evaluation of five real-time PCR assays for the detection of *Candidozyma auris* in clinical specimens. *Med Mycol* 2025;63:myaf031. PMID:40185690 <https://doi.org/10.1093/mmy/myaf031>
34. Baker AD, Gold JAW, Forsberg K, Jones S, Lyman MM. Progression from *Candida auris* colonization screening to clinical case status, United States, 2016–2023. *Emerg Infect Dis* 2025;31:1613–7. PMID:40705382 <https://doi.org/10.3201/eid3108.250315>
35. Rosa R, Abbo LM, Jimenez A, et al. Effectiveness of a sodium hypochlorite isotonic solution in decolonization of patients with *Candida auris*: learnings from a county health care system. *Am J Infect Control* 2024;52:595–8. PMID:38007101 <https://doi.org/10.1016/j.ajic.2023.11.008>
36. Lockhart SR, Chowdhary A, Gold JAW. The rapid emergence of antifungal-resistant human-pathogenic fungi. *Nat Rev Microbiol* 2023;21:818–32. PMID:37648790 <https://doi.org/10.1038/s41579-023-00960-9>
37. Giacobbe DR, Magnasco L, Sepulcri C, et al. Recent advances and future perspectives in the pharmacological treatment of *Candida auris* infections. *Expert Rev Clin Pharmacol* 2021;14:1205–20. PMID:34176393 <https://doi.org/10.1080/17512433.2021.1949285>

TABLE 1. Demographic characteristics of persons with reported *Candida auris*, number of clinical cases, and facility type in which specimen was collected — United States, 2022–2024*

Characteristic	Clinical cases
	No. (%)
Year of specimen collection	13,507 (100)
2022	2,882 (21.3)
2023	4,428 (32.8)
2024	6,197 (45.9)
Age group, yrs	12,058 (100)
0–17	21 (0.2)
18–44	1,447 (12.0)
45–64	3,886 (32.2)
65–74	3,557 (29.5)
75–84	2,365 (19.6)
≥85	782 (6.5)
Sex	12,003 (100)
Female	4,683 (39.0)
Male	7,320 (61.0)
Specimen type among clinical cases	12,958 (100)
Blood [†]	3,914 (30.2)
Respiratory	1,693 (13.1)
Urine	4,086 (31.5)
Wound	2,168 (16.7)
Other	1,097 (8.5)
Antimicrobial Resistance Laboratory Network region[§]	13,507 (100)
West	3,845 (28.5)
Midwest	2,875 (21.3)
Southeast	2,733 (20.2)
Northeast	1,571 (11.6)
Mountain	1,448 (10.7)
Mid-Atlantic	900 (6.7)
Central	135 (1.0)
Facility type where specimen was collected[¶]	11,740 (100)
Acute care hospital	8,994 (76.6)
Long-term acute care hospital	2,084 (17.8)
Skilled nursing facility, not ventilator equipped	194 (1.7)
Skilled nursing facility, ventilator equipped	91 (0.8)
Other	377 (3.2)

* A clinical case was defined as the detection of *Candida auris* from a patient specimen tested to determine the cause of an infection rather than to detect colonization. Category totals might not sum to the overall number of clinical cases because of missing data. Percentages were calculated using the number of cases with complete information for each variable as the denominator. Antimicrobial Resistance Laboratory Network region had no missing data in any year.

[†] The following number of blood specimens were collected from clinical cases, by year: in 2022, a total of 991 (34.4%); in 2023, a total of 1,337 (30.2%); and in 2024, a total of 1,586 (25.6%).

[§] Antimicrobial Resistance Laboratory Network region of the facility where the specimen was collected. Jurisdictions within each network region are as follows: *Central* (Arkansas, Iowa, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Oklahoma, and South Dakota); *Mid-Atlantic* (Delaware, District of Columbia, Maryland, North Carolina, Pennsylvania, South Carolina, Virginia, and West Virginia); *Midwest* (Illinois, Indiana, Kentucky, Michigan, Ohio, and Wisconsin); *Mountain* (Arizona, Colorado, Idaho, Montana, New Mexico, Texas, Utah, and Wyoming); *Northeast* (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, and Vermont); *Southeast* (Alabama, Florida, Georgia, Louisiana, Mississippi, Puerto Rico, and Tennessee); and *West* (Alaska, California, Guam, Hawaii, Nevada, Oregon, and Washington). [Antimicrobial Resistance Laboratory Network | CDC](https://www.cdc.gov/antimicrobial-resistance-laboratory-network/)

[¶] Does not necessarily indicate where acquisition or transmission of *C. auris* occurred.

TABLE 2. Demographic characteristics of persons with reported *Candida auris*, number of clinical cases involving blood specimens, and facility type in which specimen was collected, by year — United States, 2022–2024*

Characteristic	2022	2023	2024	Total
	No. (%)	No. (%)	No. (%)	No. (%)
Age group, yrs	782 (100)	1,143 (100)	1,419 (100)	3,344 (100)
0–17	2 (0.3)	2 (0.2)	2 (0.1)	6 (0.2)
18–44	99 (12.7)	151 (13.2)	176 (12.4)	426 (12.7)
45–64	265 (33.9)	383 (33.5)	489 (34.5)	1,137 (34.0)
65–74	229 (29.3)	333 (29.1)	418 (29.5)	980 (29.3)
75–84	139 (17.8)	226 (19.8)	265 (18.7)	630 (18.8)
≥85	48 (6.1)	48 (4.2)	69 (4.9)	165 (4.9)
Sex	777 (100)	1,141 (100)	1,413 (100)	3,331 (100)
Female	353 (45.4)	499 (43.7)	649 (45.9)	1,501 (45.1)
Male	424 (54.6)	642 (56.3)	764 (54.1)	1,830 (54.9)
Antimicrobial Resistance Laboratory Network region[†]	991 (100)	1,337 (100)	1,586 (100)	3,914 (100)
West	269 (27.1)	341 (25.5)	386 (24.3)	996 (25.4)
Southeast	191 (19.3)	268 (20.0)	350 (22.1)	809 (20.7)
Midwest	188 (19.0)	254 (19.0)	282 (17.8)	724 (18.5)
Northeast	216 (21.8)	219 (16.4)	218 (13.7)	653 (16.7)
Mountain	60 (6.1)	166 (12.4)	191 (12.0)	417 (10.7)
Mid-Atlantic	67 (6.8)	80 (6.0)	134 (8.4)	281 (7.2)
Central	0 (—)	9 (0.7)	25 (1.6)	34 (0.9)
Facility type where specimen was collected[§]	839 (100)	1,167 (100)	1,391 (100)	3,397 (100)
Acute care hospital	644 (76.8)	933 (79.9)	1,072 (77.1)	2,649 (78.0)
Long-term acute care hospital	170 (20.3)	212 (18.2)	278 (20.0)	660 (19.4)
Skilled nursing facility, not ventilator equipped	7 (0.8)	5 (0.4)	12 (0.9)	24 (0.7)
Skilled nursing facility, ventilator equipped	10 (1.2)	9 (0.8)	11 (0.8)	30 (0.9)
Other	8 (1.0)	8 (0.7)	18 (1.3)	34 (1.0)

* A clinical case was defined as the detection of *Candida auris* from a patient specimen tested to determine the cause of an infection rather than to detect colonization. Category totals might not sum to the total values because of missing data. Percentages were calculated using the number of cases with complete information for each variable as the denominator. Antimicrobial Resistance Laboratory Network region had no missing data in any year.

[†] Antimicrobial Resistance Laboratory Network region of the facility where the specimen was collected. Jurisdictions within each network region are as follows: *Central* (Arkansas, Iowa, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Oklahoma, and South Dakota); *Mid-Atlantic* (Delaware, District of Columbia, Maryland, North Carolina, Pennsylvania, South Carolina, Virginia, and West Virginia); *Midwest* (Illinois, Indiana, Kentucky, Michigan, Ohio, and Wisconsin); *Mountain* (Arizona, Colorado, Idaho, Montana, New Mexico, Texas, Utah, and Wyoming); *Northeast* (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, and Vermont); *Southeast* (Alabama, Florida, Georgia, Louisiana, Mississippi, Puerto Rico, and Tennessee); and *West* (Alaska, California, Guam, Hawaii, Nevada, Oregon, and Washington). [Antimicrobial Resistance Laboratory Network | CDC](#)

[§] Does not necessarily indicate where acquisition or transmission of *C. auris* occurred.

TABLE 3. Demographic characteristics of persons with reported *Candida auris*, number of screening cases, and facility type in which specimen was collected, by year — United States, 2022–2024*

Characteristic	2022	2023	2024	Total
	No. (%)	No. (%)	No. (%)	No. (%)
Age group, yrs	5,390 (100)	8,569 (100)	11,821 (100)	25,780 (100)
0–17	2 (<0.1)	3 (<0.1)	4 (<0.1)	9 (<0.1)
18–44	497 (9.2)	926 (10.8)	1,156 (9.8)	2,579 (10.0)
45–64	1,688 (31.3)	2,644 (30.9)	3,568 (30.2)	7,900 (30.6)
65–74	1,617 (30.0)	2,480 (28.9)	3,500 (29.6)	7,597 (29.5)
75–84	1,110 (20.6)	1,823 (21.3)	2,567 (21.7)	5,500 (21.3)
≥85	476 (8.8)	693 (8.1)	1,026 (8.7)	2,195 (8.5)
Sex	5,268 (100)	8,445 (100)	11,681 (100)	25,394 (100)
Female	2,277 (43.2)	3,540 (41.9)	4,884 (41.8)	10,701 (42.1)
Male	2,991 (56.8)	4,905 (58.1)	6,797 (58.2)	14,693 (57.9)
Antimicrobial Resistance Laboratory Network region†	6,226 (100)	9,195 (100)	12,432 (100)	27,853 (100)
West	2,242 (36.0)	3,208 (34.9)	4,575 (36.8)	10,025 (36.0)
Southeast	1,167 (18.7)	2,018 (21.9)	2,054 (16.5)	5,239 (18.8)
Midwest	1,111 (17.8)	1,676 (18.2)	2,444 (19.7)	5,231 (18.8)
Northeast	824 (13.2)	952 (10.4)	1,274 (10.2)	3,050 (11.0)
Mountain	406 (6.5)	774 (8.4)	1,099 (8.8)	2,279 (8.2)
Mid-Atlantic	473 (7.6)	522 (5.7)	734 (5.9)	1,729 (6.2)
Central	3 (<0.1)	45 (0.5)	252 (2.0)	300 (1.1)
Facility type where specimen was collected§	5,438 (100)	7,894 (100)	11,041 (100)	24,373 (100)
Acute care hospital	1,345 (24.7)	3,235 (41.0)	5,598 (50.7)	10,178 (41.8)
Long-term acute care hospital	3,048 (56.1)	3,278 (41.5)	3,937 (35.7)	10,263 (42.1)
Skilled nursing facility, not ventilator equipped	229 (4.2)	343 (4.3)	466 (4.2)	1,038 (4.3)
Skilled nursing facility, ventilator equipped	722 (13.3)	914 (11.6)	877 (7.9)	2,513 (10.3)
Other	94 (1.7)	124 (1.6)	163 (1.5)	381 (1.6)

* A screening case was defined as the detection of *Candida auris* from a swab collected to test for colonization in a patient. A patient could be classified as having up to two cases: one screening case followed by one clinical case. Category totals might not sum to the overall number of screening cases because of missing data. Percentages were calculated using the number of cases with complete information for each variable as the denominator. Antimicrobial Resistance Laboratory Network region had no missing data in any year.

† Antimicrobial Resistance Laboratory Network region of the facility where the specimen was collected. Jurisdictions within each network region are as follows: *Central* (Arkansas, Iowa, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Oklahoma, and South Dakota); *Mid-Atlantic* (Delaware, District of Columbia, Maryland, North Carolina, Pennsylvania, South Carolina, Virginia, and West Virginia); *Midwest* (Illinois, Indiana, Kentucky, Michigan, Ohio, and Wisconsin); *Mountain* (Arizona, Colorado, Idaho, Montana, New Mexico, Texas, Utah, and Wyoming); *Northeast* (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, and Vermont); *Southeast* (Alabama, Florida, Georgia, Louisiana, Mississippi, Puerto Rico, and Tennessee); and *West* (Alaska, California, Guam, Hawaii, Nevada, Oregon, and Washington). [Antimicrobial Resistance Laboratory Network | CDC](#)

§ Does not necessarily indicate where acquisition or transmission of *C. auris* occurred.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, visit *MMWR* at <https://www.cdc.gov/mmwr/index.html>.

Readers who have difficulty accessing this PDF file may access the HTML file at https://www.cdc.gov/mmwr/volumes/75/ss/ss7504a1.htm?_cid=ss7504a1_w. Address all inquiries about the *MMWR* Series to Editor-in-Chief, *MMWR* Series, Mailstop V25-5, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30329-4027 or to mmwrq@cdc.gov.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

MMWR and *Morbidity and Mortality Weekly Report* are service marks of the U.S. Department of Health and Human Services.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.

ISSN: 1546-0738 (Print)